

03.12-01/19/95-01508

FINAL  
~~REDACTED~~

REMEDIAL INVESTIGATION REPORT  
OPERABLE UNIT NO. 7  
(SITES 1, 28, AND 30)

MARINE CORPS BASE  
CAMP LEJEUNE, NORTH CAROLINA

CONTRACT TASK ORDER 0231

JANUARY 19, 1995

*Prepared For:*

DEPARTMENT OF THE NAVY  
ATLANTIC DIVISION  
NAVAL FACILITIES  
ENGINEERING COMMAND

*Norfolk, Virginia*

*Under:*

LANTDIV CLEAN Program  
Contract N62470-89-D-4814

*Prepared by:*

BAKER ENVIRONMENTAL, INC.  
*Coraopolis, Pennsylvania*

## TABLE OF CONTENTS

### APPENDICES

Volume I - Appendix A through Appendix I

- A Test Boring Records
- B Test Boring and Well Construction Records
- C Chain-of-Custody Forms
- D Field Well Development Records
- E Investigation Derived Waste Summary and Recommendations
- F Field Duplicate Summaries
- G TCLP Results
- H Grain Size and Wet Chemistry Results
- I QA/QC Summary

Volume II - Appendix J through Appendix L

- J Sampling Summary
- K Data and Frequency Summaries
- L Statistical Summaries

Volume III - Appendix M through Appendix U

- M Base Background
- N White Oak River Basin Reference Station
- O Shower Model
- P Lead UBK Model
- Q Site Conceptual Model
- R CDI Calculations
- S Summary of Groundwater Data and Aquifer Characteristics at MCB Camp Lejeune
- T Sampling Station Characterization Data Sheet
- U Biota Population

**APPENDIX M**  
**BASE BACKGROUND**

**SOIL**

---

**BASE BACKGROUND**  
**SURFACE SOIL**  
**MCB CAMP LEJEUNE, NORTH CAROLINA**  
**TAL INORGANICS**

6-201N-SB11-00      6-201N-SB12-00      6-201C-SB38-00      6-201C-SB39-00      78-BB-SB-00      41-BB-SB01-00      41-BB-SB02-00

	6-201N-SB11-00	6-201N-SB12-00	6-201C-SB38-00	6-201C-SB39-00	78-BB-SB-00	41-BB-SB01-00	41-BB-SB02-00
Aluminum	1120	45.25	748	245	1490	528	1430
Antimony	4.7	4.8	1.4	1.3	0.33	2.07	0.865
Arsenic	0.28	0.29	0.91	0.28	0.22	0.356	0.317
Barium	2	2.05	16.5	3.5	8.6	1.525	4.06
Beryllium	0.095	0.1	0.03	0.03	0.11	0.1	0.09
Cadmium	0.285	0.295	0.58	0.175	0.55	0.392	0.349
Calcium	178	108	10700	402	941	18.3	54.6
Chromium	0.475	0.49	1.6	0.33	2.2	1.02	0.91
Cobalt	0.85	0.9	0.195	0.185	1.8	1.965	1.75
Copper	0.55	0.6	3.1	0.75	2	2	87.2
Iron	525	160	684	238	1020	83	970
Lead	2	3	62.9	25.1	20.4	2.59	10.9
Magnesium	11.65	10.1	200	26	118	8.85	39.1
Manganese	3.1	1	16	4.5	11.1	0.87	10.2
Mercury	0.01	0.01	0.05	0.06	0.05	0.0305	0.078
Nickel	1.6	1.65	0.8	0.75	2.2	3.55	3.15
Potassium	36.55	37.5	54.5	30.6	102	91.5	81.5
Selenium	0.47	0.485	0.5	0.465	0.31	0.311	0.277
Silver	0.95	1	0.195	0.185	0.33	0.1965	0.175
Sodium	19.65	15.85	14	4.7	67.5	44.1	39.3
Thallium	0.19	0.195	0.205	0.185	0.11	0.565	0.505
Vanadium	1.05	0.8	2.8	1.6	5.3	2.505	2.23
Zinc	0.55	0.8	23.1	4.6	28.3	2.66	6.11

Concentrations are in milligrams per kilogram (mg/kg).

Qualifiers have been removed per Baker's standards.

Qualifiers R, U, and UJ have been given one-half the detection value.

Qualifiers J, NJ, and B have been removed with no detection value change.

BASE BACKGROUND  
SURFACE SOIL  
MCB CAMP LEJEUNE, NORTH CAROLINA  
TAL INORGANICS

41-BB-SB03-00 41-BB-SB04-00 69-BB-SB01-00 69-BB-SB02-00 69-BB-SB03-00 69-BB-SB04-00 74-BB-SB01-00

	41-BB-SB03-00	41-BB-SB04-00	69-BB-SB01-00	69-BB-SB02-00	69-BB-SB03-00	69-BB-SB04-00	74-BB-SB01-00
Aluminum	2100	5370	1310	4150	9570	5360	3110
Antimony	0.87	0.94	0.85	0.95	0.95	0.95	0.905
Arsenic	0.3205	0.345	0.31	0.345	0.79	0.35	0.3325
Barium	4.53	13.4	5.6	15.4	19.6	20.8	11.1
Beryllium	0.09	0.095	0.14	0.155	0.155	0.155	0.148
Cadmium	0.3525	0.38	0.26	0.285	0.29	0.29	0.2695
Calcium	79.2	46.3	28.2	43.6	282	53	181
Chromium	2.64	3.24	0.75	4	12.5	5.8	0.84
Cobalt	1.77	1.905	2.1	2.3	2.35	2.35	2.225
Copper	1.8	1.94	1.75	1.9	1.95	1.95	4.56
Iron	1120	2160	425	1430	9640	3890	1740
Lead	9.98	6.61	2.8	6	5.3	5.6	5.19
Magnesium	74	144	37.3	91.8	610	247	70
Manganese	11.6	11.8	15.1	12.7	12.3	8.3	9.44
Mercury	0.057	0.08	0.015	0.06	0.045	0.025	0.04
Nickel	3.2	3.45	2.9	1.6	1.65	1.65	1.56
Potassium	190	177	32.25	35.5	361	106	87.5
Selenium	0.2795	0.301	0.27	0.295	0.3	0.3	0.29
Silver	0.177	0.1905	0.045	0.045	4.3	0.39	0.046
Sodium	39.65	42.75	20	22	22.4	22.3	70.4
Thallium	0.51	0.55	0.495	0.55	0.55	0.55	0.53
Vanadium	2.255	2.43	1.8	1.95	13.5	5.6	5.21
Zinc	5.97	7.15	3.1	5.2	10.8	7.9	1.27

Concentrations are in millograms per kilogram (mg/kg).

Qualifiers have been removed per Baker's standards.

Qualifiers R, U, and UJ have been given one-half the detection value.

Qualifiers J, NJ, and B have been removed with no detection value change.

**BASE BACKGROUND**  
**SURFACE SOIL**  
**MCB CAMP LEJEUNE, NORTH CAROLINA**  
**TAL INORGANICS**

74-BB-SB02-00 74-BB-SB03-00 74-BB-SB04-00 1-BB-SB38-00 1-BB-SB39-00 1-GW13-00 28-BB-SB37-00 28-BB-SB38-00

	74-BB-SB02-00	74-BB-SB03-00	74-BB-SB04-00	1-BB-SB38-00	1-BB-SB39-00	1-GW13-00	28-BB-SB37-00	28-BB-SB38-00
Aluminum	1730	1000	2100	3920	4930	1600	2840	379
Antimony	0.925	0.855	0.96	3.6	3.15	8.0	3.55	2.9
Arsenic	0.339	0.314	0.352	0.315	0.28	0.29	0.31	0.255
Barium	1.6	3.12	16	9.6	9.3	2.8	5.1	1.8
Beryllium	0.151	0.14	0.1565	0.105	0.10	0.095	0.105	0.085
Cadmium	0.275	0.2545	0.285	0.315	0.28	0.285	0.31	0.255
Calcium	46.9	43.9	377	538	353	248	114	13.10
Chromium	2.7	0.795	1.98	3.5	4.7	4.1	2.0	0.60
Cobalt	2.27	2.1	2.355	0.42	0.375	0.38	0.415	0.34
Copper	3.92	1.755	1.965	1.6	0.6	1.9	0.6	0.50
Iron	401	787	1640	2270	1470	1000	1210	444
Lead	3.79	1.14	142	5.9	4.5	4.2	2.8	1.7
Magnesium	37.5	16.1	52.5	152	183	47.2	68.8	12.9
Manganese	3.13	7.37	4.61	10.6	4.2	5.9	2.7	3.3
Mercury	0.048	0.0305	0.05	0.03	0.025	0.03	0.025	0.025
Nickel	1.59	1.475	1.65	0.8	0.65	0.65	0.750	0.6
Potassium	89	82.5	92.5	149	153	20.650	29.75	8.35
Selenium	0.296	0.274	0.307	0.42	0.375	0.38	0.415	0.34
Silver	0.047	0.0435	0.0485	0.5	0.465	0.475	0.5	0.425
Sodium	71.8	87.6	122	11.0	17.2	7.25	28.5	18.2
Thallium	0.54	0.4985	0.56	0.42	0.38	0.38	0.415	0.34
Vanadium	1.94	1.8	4.69	7.9	6.1	3.5	3.6	2.1
Zinc	1.15	1.97	2.87	7.2	4.0	1.4	0.9	0.71

Concentrations are in milligrams per kilogram (mg/kg).

Qualifiers have been removed per Baker's standards.

Qualifiers R, U, and UJ have been given one-half the detection value.

Qualifiers J, NJ, and B have been removed with no detection value change.

**BASE BACKGROUND**  
**SURFACE SOIL**  
**MCB CAMP LEJEUNE, NORTH CAROLINA**  
**TAL INORGANICS**

28-GW09DW-00 30-BB-SB12-00 30-BB-SB13-00 30-BB-SB14-00 30-BB-SB15-00 30-BB-SB16-00 30-GW03-00 35-SS01-00

Aluminum	5460	54.6	24.9	49.2	37.5	196	17.7	2220.0
Antimony	3.35	3.2	3.2	3.3	3.5	3.650	3.9	2.45
Arsenic	1.8	0.28	0.29	0.29	0.31	0.325	0.34	0.065
Barium	11.6	1.8	0.7	0.7	0.7	3.100	0.8	15.6
Beryllium	0.10	0.095	0.10	0.10	0.10	0.110	0.12	0.11
Cadmium	0.295	0.28	0.29	0.29	0.31	0.325	0.34	0.04
Calcium	368	11.45	4.3	9.9	9.0	172	5.2	605.0
Chromium	6.0	1.6	0.7	1.9	0.7	0.75	0.8	1.9
Cobalt	0.91	0.375	0.38	0.38	0.41	0.43	0.45	0.60
Copper	2.9	0.55	0.6	0.6	0.6	0.65	0.7	3.9
Iron	2250	276	102	218	69.7	167	80.4	1250.0
Lead	11.6	3.3	0.47	2.4	0.73	4.4	0.86	3.60
Magnesium	157	6.5	2.6	2.6	2.8	37.1	3.1	71.6
Manganese	4.1	11.9	4.4	9.5	1.3	2.5	2.3	5.5
Mercury	0.025	0.06	0.02	0.03	0.05	0.03	0.03	0.065
Nickel	1.9	0.65	0.7	0.7	1.7	0.9	0.8	1.3
Potassium	158	8.25	11.1	3.8	1.0	29.6	1.2	129.5
Selenium	0.94	0.375	0.38	0.38	0.41	0.43	0.45	0.075
Silver	0.49	0.47	0.47	0.48	0.5	0.6	0.6	0.16
Sodium	15.0	14.8	26.0	4.9	5.2	18.2	5.8	126.00
Thallium	0.395	0.375	0.38	0.38	0.41	0.43	0.45	0.06
Vanadium	8.3	1.7	0.75	1.7	0.31	0.76	0.34	3.60
Zinc	6.6	0.35	0.30	0.48	1.7	2.0	1.2	7.4

Concentrations are in millograms per kilogram (mg/kg).

Qualifiers have been removed per Baker's standards.

Qualifiers R, U, and UJ have been given one-half the detection value.

Qualifiers J, NJ, and B have been removed with no detection value change.

**BASE BACKGROUND  
SURFACE SOIL  
MCB CAMP LEJEUNE, NORTH CAROLINA  
TAL INORGANICS**

	MIN	MAX	AVG	2Xaverage
--	-----	-----	-----	-----------

Aluminum	17.7	9570	2104.505	4209.010
Antimony	0.33	8	2.406	4.811
Arsenic	0.065	1.8	0.386	0.772
Barium	0.65	20.8	7.096	14.192
Beryllium	0.03	0.1565	0.108	0.216
Cadmium	0.04	0.58	0.306	0.611
Calcium	4.25	10700	534.460	1068.920
Chromium	0.33	12.5	2.382	4.765
Cobalt	0.185	2.355	1.174	2.348
Copper	0.5	87.2	4.508	9.016
Iron	69.7	9640	1257.337	2514.673
Lead	0.47	142	12.059	24.117
Magnesium	2.55	610	84.698	169.397
Manganese	0.87	16	7.044	14.088
Mercury	0.01	0.08	0.039	0.078
Nickel	0.6	3.55	1.546	3.092
Potassium	1	361	79.682	159.363
Selenium	0.075	0.94	0.370	0.739
Silver	0.0435	4.3	0.480	0.960
Sodium	4.7	126	34.132	68.263
Thallium	0.06	0.565	0.403	0.806
Vanadium	0.305	13.5	3.271	6.541
Zinc	0.3	28.3	4.920	9.839

Concentrations are in milligrams per kilogram (mg/kg).

Qualifiers have been removed per Baker's standards.

Qualifiers R, U, and UJ have been given one-half the detection value.

Qualifiers J, NJ, and B have been removed with no detection value change.

**BASE BACKGROUND**  
**SUBSURFACE SOIL**  
**MCB CAMP LEJEUNE, NORTH CAROLINA**  
**TAL INORGANICS**

6-201N-SB11-07      6-201N-SB12-02    6-201C-SB38-01    6-201C-SB39-04    78-BB-SB-01    2-GW09-01    1-BB-SB38-05

Aluminum	672	857	3620	2970	10200	8520	4580.000
Antimony	4.7	4.85	1.4	1.25	0.355	1.6	4.200
Arsenic	0.31	0.315	0.033	0.305	0.24	0.47	1.100
Barium	2	2.05	7.6	6.5	10.9	6.6	7.500
Beryllium	0.095	0.1	0.03	0.025	0.12	0.23	0.125
Cadmium	0.285	0.295	0.57	0.17	0.6	1.2	0.370
Calcium	5.35	5.4	4410	12.1	81.3	10.6	35.600
Chromium	1.6	1.85	6	2.2	5.7	8.7	10.500
Cobalt	0.65	0.9	0.235	0.175	0.95	1.9	0.495
Copper	0.475	0.6	1.7	0.65	0.95	0.47	6.600
Iron	257	126	456	833	822	2840	4940.000
Lead	1.2	1.6	11.5	2.7	6.1	4.3	5.100
Magnesium	13.1	12.7	133	86.8	188	260	222.000
Manganese	0.475	0.395	7.5	2.6	2.4	5.2	4.100
Mercury	0.01	0.01	0.04	0.015	0.045	0.11	0.025
Nickel	1.6	1.7	0.8	0.7	2.4	4.7	0.850
Potassium	48.9	40.8	84.7	187	123	184	409.000
Selenium	0.5	0.5	0.55	0.5	0.29	0.115	0.495
Silver	0.95	1	0.195	0.175	0.355	0.7	0.600
Sodium	12.7	12.15	13.25	7.25	44.9	31.5	12.850
Thallium	0.205	0.21	0.22	0.2	0.12	0.23	0.495
Vanadium	0.75	1	3	4.7	7.4	13.4	12.200
Zinc	0.475	0.395	11.6	0.9	2.1	1.4	4.700
					0.285	0.57	

Concentrations are in milligrams per kilograms (mg/kg).

Qualifiers have been removed per Baker's standards.

Qualifiers R, U, and UJ have been given one-half the detection value.

Qualifiers J, NJ, and B have been removed with no detection value change.

**BASE BACKGROUND**  
**SUBSURFACE SOIL**  
**MCB CAMP LEJEUNE, NORTH CAROLINA**  
**TAL INORGANICS**

1-BB-SB39-04 1-BB-SB39-06 1-GW13-04 1-GW13-08 28-BB-SB37-03 28-BB-SB38-04 28-GW09DW-01 30-BB-SB12-03

Aluminum	6180.000	5980.000	4160.000	6600.000	5170.000	2830.000	5730.000	2970
Antimony	3.250	2.950	6.900	3.200	3.550	3.550	3.750	3.9
Arsenic	0.290	0.260	0.285	0.280	0.315	0.315	1.500	0.34
Barium	11.800	8.600	7.500	8.400	9.700	5.000	11.700	0.8
Beryllium	0.095	0.085	0.095	0.095	0.105	0.105	0.110	0.12
Cadmium	0.290	0.260	0.285	0.280	0.315	0.315	0.330	0.34
Calcium	12.250	19.700	52.400	92.600	23.450	6.850	441.000	7.0
Chromium	5.500	5.300	7.100	8.300	7.300	3.400	4.700	3.9
Cobalt	0.385	0.350	0.380	0.375	0.420	0.420	0.930	0.45
Copper	0.600	0.500	2.100	1.600	0.650	0.650	0.650	0.7
Iron	1510.000	1210.000	567.000	959.000	2090.000	749.000	2780.000	908
Lead	3.800	3.100	3.300	4.000	4.100	2.300	7.400	0.7
Magnesium	189.000	217.000	131.000	262.000	153.000	66.000	157.000	24.7
Manganese	4.900	5.400	2.000	4.500	3.200	1.500	5.300	1.7
Mercury	0.025	0.020	0.050	0.025	0.025	0.025	0.025	0.03
Nickel	2.300	0.600	0.650	0.650	0.750	0.750	1.000	0.8
Potassium	191.000	268.000	98.100	308.000	122.000	91.300	136.000	13.2
Selenium	0.385	0.350	0.380	0.375	0.420	0.420	0.440	0.45
Silver	0.480	0.435	0.475	0.470	0.500	0.550	0.550	0.6
Sodium	21.600	9.200	9.600	10.900	33.800	28.600	20.300	12.5
Thallium	0.385	0.350	0.380	0.375	0.420	0.420	0.440	0.45
Vanadium	6.500	6.100	3.500	10.100	6.400	2.800	8.500	6.2
Zinc	2.900	2.400	1.000	2.700	1.900	0.970	4.200	0.35

Concentrations are in milligrams per kilograms (mg/kg).

Qualifiers have been removed per Baker's standards.

Qualifiers R, U, and UJ have been given one-half the detection value.

Qualifiers J, NJ, and B have been removed with no detection value change.

**BASE BACKGROUND**  
**SUBSURFACE SOIL**  
**MCB CAMP LEJEUNE, NORTH CAROLINA**  
**TAL INORGANICS**

30-BB-SB13-01 30-BB-SB14-01 30-BB-SB15-01 30-BB-SB16-02 30-GW03-01 35-GWDS01-03

Aluminum	17.1	25.7	42.6	777	16.9	2910.0
Antimony	3.1	3.6	3.6	3.4	3.9	2.750
Arsenic	0.28	0.32	0.32	0.30	0.34	0.12
Barium	0.7	0.8	0.8	3.5	0.8	5.5
Beryllium	0.09	0.11	0.11	0.10	0.12	0.06
Cadmium	0.28	0.32	0.32	0.30	0.34	0.30
Calcium	6.9	4.8	6.3	116	6.6	456.0
Chromium	0.7	0.8	0.8	0.7	0.8	2.2
Cobalt	0.37	0.42	0.43	0.40	0.46	0.65
Copper	0.6	0.7	0.7	0.6	0.7	0.550
Iron	95.9	155	63.3	514	74.5	442.0
Lead	0.47	1.9	0.91	3.2	0.59	8.10
Magnesium	7.5	2.9	2.9	30.2	3.1	63.5
Manganese	4.3	6.7	1.1	3.7	1.7	5.6
Mercury	0.03	0.08	0.25	0.03	0.68	0.03
Nickel	0.7	0.8	2.2	1.7	0.8	1.050
Potassium	6.3	1.1	21.3	21.9	1.2	145.0
Selenium	0.37	0.42	0.43	0.40	0.46	0.085
Silver	0.46	0.6	0.6	0.50	0.6	0.39
Sodium	11.1	19.3	5.4	14.4	5.8	141.0
Thallium	0.37	0.42	0.43	0.40	0.46	0.06
Vanadium	0.73	1.0	0.84	1.6	0.34	3.00
Zinc	0.32	0.39	1.2	1.7	1.3	2.6

Concentrations are in milligrams per kilograms (mg/kg).

Qualifiers have been removed per Baker's standards.

Qualifiers R, U, and UJ have been given one-half the detection value.

Qualifiers J, NJ, and B have been removed with no detection value change.

**BASE BACKGROUND**  
**SUBSURFACE SOIL**  
**MCB CAMP LEJEUNE, NORTH CAROLINA**  
**TAL INORGANICS**

	MIN	MAX	AVG	2Xaverage
Aluminum	16.900	10200.000	3563.252	7126.505
Antimony	0.355	6.900	3.317	6.634
Arsenic	0.033	1.500	0.382	0.764
Barium	0.650	11.800	5.648	11.295
Beryllium	0.025	0.230	0.100	0.200
Cadmium	0.170	1.200	0.369	0.738
Calcium	4.750	4410.000	276.762	553.524
Chromium	0.650	10.500	4.186	8.371
Cobalt	0.175	1.900	0.559	1.117
Copper	0.470	6.600	1.076	2.152
Iron	63.300	4940.000	1066.271	2132.543
Lead	0.465	11.500	3.636	7.273
Magnesium	2.850	262.000	105.964	211.929
Manganese	0.395	7.500	3.537	7.073
Mercury	0.010	0.680	0.075	0.150
Nickel	0.600	4.700	1.305	2.610
Potassium	1.050	409.000	119.126	238.252
Selenium	0.085	0.550	0.396	0.792
Silver	0.175	1.000	0.523	1.045
Sodium	5.400	141.000	22.767	45.533
Thallium	0.060	0.495	0.335	0.669
Vanadium	0.340	13.400	4.765	9.530
Zinc	0.320	11.600	2.162	4.323

Concentrations are in milligrams per kilograms (mg/kg).

Qualifiers have been removed per Baker's standards.

Qualifiers R, U, and UJ have been given one-half the detection value.

Qualifiers J, NJ, and B have been removed with no detection value change.

## **GROUNDWATER**

---

**DRAFT**

**EVALUATION OF METALS IN  
GROUNDWATER**

**MARINE CORPS BASE,  
CAMP LEJEUNE, NORTH CAROLINA**

**CONTRACT TASK ORDER 0177**

**JUNE 3, 1994**

*Prepared for:*

**DEPARTMENT OF THE NAVY  
ATLANTIC DIVISION  
NAVAL FACILITIES  
ENGINEERING COMMAND  
*Norfolk, Virginia***

*Under the:*

**LANTDIV CLEAN Program  
Contract N62470-89-D-4814**

*Prepared by:*

**BAKER ENVIRONMENTAL, INC.  
*Coraopolis, Pennsylvania***

## TABLE OF CONTENTS

	<u>Page</u>
<b>1.0 INTRODUCTION .....</b>	<b>1</b>
<b>2.0 STUDY OBJECTIVES .....</b>	<b>1</b>
<b>3.0 SCOPE OF WORK .....</b>	<b>2</b>
<b>4.0 DATA ANALYSIS .....</b>	<b>3</b>
<b>5.0 ANALYSIS OF STUDY OBJECTIVES .....</b>	<b>8</b>
<b>6.0 CONCLUSIONS .....</b>	<b>10</b>
<b>7.0 RECOMMENDATIONS .....</b>	<b>10</b>

### FIGURES

- 1 Site Location Map
- 2 Positive Detections Above Applicable Federal and State Standards for Total and Filtered Inorganic Analytes in Groundwater-Site 2
- 3 Positive Detections of Total Metals Above Federal MCLs and NCWQS in Shallow Wells-Site 78
- 4 Positive Detections of Total Metals Above Federal MCLs and NCWQS in Intermediate Wells-Site 78
- 5 Positive Detections of Total Metals Above Federal MCLs and NCWQS in Deep Wells-Site 78

### TABLES

- 1 Summary of Total Metals in Shallow Wells
- 2 Comparison of Repeat Sampling in Shallow Wells
- 3 Summary of Dissolved Metals in Shallow Wells
- 4 Summary of Total Metals in Upgradient Wells
- 5 Comparison of Inorganic Subsurface Soil Concentrations in "Clean" and "Contaminated" Wells
- 6 Total Metals in Deep Monitoring Wells
- 7 Summary of Field Parameters in Shallow, Deep, and Supply Wells

## **1.0 INTRODUCTION**

Numerous groundwater investigations have been conducted at Marine Corps Base (MCB), Camp Lejeune under the Department of the Navy (DON) Installation Restoration Program (IRP). These studies have identified elevated levels of total metals in shallow groundwater at almost every site. The degree of contamination, based on dissolved metals analysis of groundwater samples, is limited. It is believed that the presence of elevated metals are not always related to past disposal activities for several reasons, which is the basis of this study.

Currently, Records of Decision (ROD) are being prepared for Operable Units No. 1 (Sites 21, 24, and 78) and No. 5 (Site 2). Both RODs are proposing to not remediate shallow groundwater which contains elevated levels of total metals above State groundwater standards (i.e., North Carolina Water Quality Standards) and/or Federal drinking water standards (i.e., Maximum Contaminant Levels). Specifically, remediation of shallow groundwater due to elevated total metals is not cost effective, or practical, due to the following: (1) the shallow aquifer is not used for potable supply; (2) the source of metals in groundwater cannot be correlated with soil data or previous disposal practices; (3) the extent of shallow groundwater contamination (based on total metals analysis) is widespread and in many cases undefinable, since there are no apparent contaminant plumes or patterns associated with the metals; and (4) deep groundwater, which is the source of potable water, is not significantly contaminated with metals above the standards.

## **2.0 STUDY OBJECTIVES**

The DON/Marine Corps initiated a study on inorganics in groundwater throughout MCB Camp Lejeune to assess whether total metals in groundwater are related to disposal practices or to other factors. The overall goal of this study is to provide information that would be used in consideration of not remediating shallow groundwater at Operable Units No. 1 and No. 5, and possibly other operable units where total metals are elevated without cause. The following study objectives were identified:

- (1) Determine whether the elevated total metals detected in the shallow aquifer are related to past disposal practices, well construction factors, sampling techniques, or suspended particulates in the samples;
- (2) Determine whether total metals in shallow groundwater are elevated throughout the region or MCB Camp Lejeune;
- (3) Determine whether there is a correlation between elevated total metals in groundwater and metals in soil; and

- (4) Determine whether the concentrations of total metals (i.e., low versus high) is related to shallow and deep aquifer characteristics.

### 3.0 SCOPE OF WORK

Groundwater and soil data from a total of 21 sites were compiled as part of the overall study. Three of the 21 sites are located outside the boundary of the base. These sites include the ABC Cleaners Superfund Site, located along Route 24 in Jacksonville, and two sites located along Highway 17 (Off-site Properties No. 1 and No. 2). The two sites along Route 17 were investigated by the DON/Marine Corps as part of a real estate survey. The other 18 sites are located throughout various portions of MCB Camp Lejeune (see Figure 1).

Information from studies conducted by Baker and other consultants were obtained to evaluate metal concentrations in groundwater. The study focused on 14 metals of potential concern to human health and the environment. Some of the information was collected under the IR Program whereas other information was obtained during other investigations (e.g., ABC Cleaners RI/FS). The following data tables were then prepared to determine why total metals are generally elevated in shallow groundwater.

Table 1 - Total Metal Concentrations in Shallow Groundwater by Site

Table 2 - Summary of Repeat Sampling of Shallow Wells (Sites 2 and 78)

Table 3 - Dissolved Metal Concentrations in Shallow Groundwater by Site

Table 4 - Summary of Total Metal Concentrations in Upgradient Wells

Table 5 - Comparison of Subsurface Metal Concentrations in Uncontaminated and Contaminated Wells

Table 6 - Total Metal Concentrations in Deep Groundwater by Site

Table 7 - Summary of Field Parameters in Shallow Monitoring Wells, Deep Monitoring Wells, and Supply Wells

The tables are presented at the end of this report.

#### 4.0 DATA ANALYSIS

The following discussion represents an analysis of the information contained in each of the previously mentioned tables.

Table 1 (Total Metal Concentrations in Shallow Groundwater)

All of the sites had at least one (and in most cases several) metal which exceeded either State water quality standards or Federal drinking water standards. The most frequently detected metals included chromium, lead, and manganese, which were detected at almost every site above drinking water standards. Other frequently detected metals which exceeded drinking water standards included arsenic, beryllium, cadmium, and nickel.

An analysis of the data from Table 1 indicates that elevated total metals are present in shallow groundwater at every site, including the three sites which are located off base. The two sites which did not exhibit significant contamination include the ABC Cleaners site (only chromium exceeded the standards) and Site 48 (only manganese exceeded the standards).

Total metals detected in shallow groundwater at Site 2 exceeded State and/or Federal standards in seven of the 11 shallow monitoring wells. Manganese was the most frequently detected metal (7/11). Lead (3/11), chromium (2/11), and cadmium (1/11) were also detected above the standards, but less frequently (see Figure 2).

With the exception of Wells 78GW03 and 78GW19, total metals were detected at Site 78 (Hadnot Point Industrial Area) above Federal MCLs or NCWQS in every shallow well (see Figure 3). The extent of elevated total metals in groundwater is widespread, encompassing approximately one square mile (or approximately 660 acres) in total area. The distribution and concentration of total metals in shallow groundwater makes it virtually impossible to identify or illustrate contaminant plumes (see Figure 3).

An analysis of the total metals results indicates the following pattern. Samples exhibiting elevated levels of lead, chromium, or other contaminants of concern, also exhibited elevated levels of other metals such as aluminum, antimony, iron, and zinc. Samples which did not exhibit elevated levels of lead, chromium, or manganese also did not exhibit elevated levels of other metals. This pattern indicates that the elevated total metals are not limited to one or

two contaminants, which would be the case if a lead or chromium plume in the groundwater truly existed. In other words, if a site is impacted by a particular metal due to disposal activities (say chromium for example), then other metals such as aluminum, lead, or zinc should not be consistently elevated as in the case of samples collected from the shallow aquifer at MCB Camp Lejeune. This point is depicted in the data summary tables provided in Appendix A for Sites 2 and 78. These tables were taken from the Remedial Investigation Reports for Operable Units No. 1 and No. 5. As an example, note that sample numbers 78-MW08, 78-MW10, 78-MW11, and 78-MW12 all had elevated levels of total metals when compared to samples 78-MW09-2 and 78-MW09-3. It is clear that most of the metal concentrations in a particular sample follow a consistent pattern throughout.

Table 2 (Comparison of Repeat Sampling of Shallow Wells)

Five wells from Sites 2 and 78 were randomly chosen to evaluate total metals concentrations between sampling rounds. The comparison was limited to only chromium, lead, and manganese since these contaminants were frequently detected throughout MCB Camp Lejeune. In several cases, metal concentrations were significantly different between the sampling rounds. If the shallow aquifer was impacted due to former disposal activities, a contaminant plume would be present and concentrations would not significantly deviate. The deviation in metal concentrations may indicate that sampling results are biased due to suspended particulates in the samples.

Table 3 (Dissolved Metal Concentration in Shallow Groundwater by Site)

The data base for Table 3 was limited to 12 sites since many of the previous investigations (i.e., prior to Navy CLEAN) did not analyze for dissolved metals. Nevertheless, an analysis of the 12 sites revealed that elevated levels of dissolved metals in groundwater is limited. Manganese was the most frequently detected metal above drinking water standards (10 of 12 sites exhibited elevated levels). Lead was detected at only one site (Site 21) above drinking water standards. Chromium was also detected at only one site (Site 78) above drinking water standards. No other metal was detected above the standards.

Literature searches have indicated that manganese is a naturally occurring metal in North Carolina. Therefore, the presence of manganese may not be attributable to site-related activities (Greenhorne & O'Mara, 1992).

An analysis of the data from Table 3 clearly shows a significant reduction in metal concentrations when compared to Table 1 (total metals in shallow groundwater). One possible reason for this reduction is that suspended solids or particles are not being introduced into the analysis of the sample due to filtering. A second possibility is that the metals are not significantly present in a dissolved state in shallow groundwater due to the species of metals under site conditions. It should be noted that calcium and sodium did not exhibit such a pattern since the salts of these metals are more soluble in water. For example, the concentrations of total calcium and total sodium versus dissolved calcium and dissolved sodium are similar and are not affected by the removal of the particulates during filtering. The fact that these salts do not exhibit the pattern that the other metals show supports the possibility that total metal concentrations are influenced by particulates in the sample.

**Table 4 (Total Metals in Upgradient Shallow Wells)**

The data base for Table 4 consists of groundwater results from 14 upgradient shallow monitoring wells (i.e., one well per site). These wells were installed to determine baseline groundwater quality to which on-site groundwater conditions could be compared. In some cases, the upgradient wells were located in areas where other base activities may have influenced groundwater quality.

The analysis of this data shows that manganese was the most frequently detected metal above Federal or State standards in upgradient shallow wells. Manganese was detected in 7 of the 14 upgradient wells above drinking water standards. Chromium and lead were also frequently detected above drinking water standards in upgradient (background) wells. These contaminants were detected in 6 of the 14 upgradient wells. At Site 2, samples collected from an upgradient well (2GW9) exhibited elevated levels of chromium ( $83\mu\text{l/l}$ ), lead ( $27.2\mu\text{l/l}$ ) and manganese ( $747\mu\text{l/l}$ ). At Site 78, samples collected from upgradient wells 96W4 and 78GW26 did not exhibit elevated levels of total metals. The concentration range for metals detected above NC WQS and/or Federal MCLs in upgradient wells is provided below:

- beryllium (ND-46.5  $\mu\text{l/l}$ )
- cadmium (ND-10  $\mu\text{l/l}$ )
- chromium (ND-198  $\mu\text{l/l}$ )
- lead (ND-78.8  $\mu\text{l/l}$ )
- manganese (ND-747  $\mu\text{l/l}$ )
- mercury (ND-1.6J  $\mu\text{l/l}$ )

Based on the above range representing upgradient wells, none of the on-site wells at Site 2 exhibited total metals above the maximum background concentrations. However, at Site 78, lead and chromium were detected above the maximum background in several on-site wells.

An analysis of the data from Table 4 indicates that shallow groundwater upgradient of some sites contains total metals above drinking water standards. A comparison of Table 4 data against Table 1 data indicates that shallow groundwater samples from upgradient wells are less contaminated than samples collected from on-site monitoring wells. However, it should be noted that the data base for Table 4 consists of only 14 wells whereas the data base for Table 1 consists of over 130 wells. Therefore, to assume that upgradient groundwater quality is better than on-site groundwater quality may not be justified due to the different data bases.

Table 5 (Comparison of Subsurface Metal Concentrations in Uncontaminated and Contaminated Wells)

The purpose of this table is to determine whether metal concentrations in soils correlate with the elevated levels of metals in shallow groundwater.

To evaluate this, metals in subsurface soils, representing an area of groundwater contamination, were compared to metals in subsurface soil in areas which did not exhibit groundwater contamination. If the elevated total metals in shallow groundwater are present due to former disposal activities, subsurface metals in soil representing an area of groundwater contamination would be expected to be elevated or higher than metals in subsurface soil representing a non-contaminated area. This evaluation assumes that the well exhibiting elevated total metals is within a source area and that the soil sample is representative of soil impacted by metal contamination.

As shown on Table 5, there is no clear pattern or correlation which indicates that elevated total metals are due to soil contamination. Note that in many cases, the concentration of metals which represent "non-contaminated" areas are greater than the metals which represent "contaminated" areas. Also note that the metals in subsurface soil are within or close to background subsurface metal concentrations. Therefore, this supports the possibility that in many cases at MCB Camp Lejeune, the elevated total metals in shallow groundwater cannot be attributable to a source or to past disposal practices.

**Table 6 (Total Metals in Deep Monitoring Wells)**

Table 6 presents total metal concentrations in deep groundwater for each site. The data base is limited to only 8 sites. Metal concentrations in supply wells were also included for comparison purposes.

As shown on Table 6, total metals in deep groundwater are below drinking water standards with a few exceptions. Arsenic and cadmium were detected above the standards in one deep monitoring well at Site 78 (see Figure 4). Manganese was detected in deep groundwater at three sites and a few of the supply wells. Lead was detected in one supply well at 16  $\mu\text{l}$ , which is slightly above the drinking water standard of 15  $\mu\text{l}$ .

Elevated total metals are not widespread in deep groundwater for two possible reasons. First, most metals are not very mobile in the environment. Second, deep groundwater samples may not have significant amounts of suspended particulates due to different geologic conditions. Soils in the deeper aquifer are more compacted and consist primarily of calcareous sands, clays, and limestone fragments. Soils in the shallow aquifer are loosely compacted and consist primarily of fine-grained sands, silts, and clays. This classification may support the possibility that suspended solids are collected during sampling, thereby influencing the analysis for total metals.

**Table 7 (Summary of Field Parameters in Shallow, Deep, and Supply Wells)**

Table 7 provides a range of pH and specific conductivity values representative of shallow and deep groundwater. In general, lower pH values were noted more often in shallow wells than in deep wells (including the supply wells). This condition may influence the leachability and speciation of metals in groundwater.

Deep groundwater usually exhibited higher specific conductivity values. High specific conductivity values are representative of high dissolved conditions. The fact that deep groundwater generally exhibited higher specific conductivity values indicates that most of the metals, if present, are in a dissolved state. The high specific conductivity values could also indicate less suspended particulates due to the geologic conditions of the deep aquifer. The lower specific conductivity values observed in shallow wells indicates that the metals in the shallow aquifer are not in a dissolved state. This also supports the possibility that suspended particulates in the shallow aquifer are influencing the analysis of total metals.

## 5.0 ANALYSIS OF THE STUDY OBJECTIVES

Each of the objectives identified for this study are analyzed below based on the information collected.

Objective No. 1 (Determine whether the elevated total metals in the shallow aquifer are related to past disposal practices, well construction factors, sampling techniques, or suspended particulates in the samples)

Based on the analysis of information provided in Tables 1 through 7 and Appendix A, it appears that suspended particulates in groundwater samples could influence the concentration of total metals in groundwater. Well construction factors and sampling techniques are probably not a significant factor since the data base is representative of data obtained by Baker, ESE (Site 28 and 30), Roy F. Weston (ABC Cleaners), and Halliburton NUS (Site 7). No particular pattern was noted between sites which Baker obtained the samples versus sites in which other consultants obtained the data. Sampling methods were also considered. For Sites 63 and 65 for example, samples were collected with a bailer. At Sites 2 and 78, samples were collected with a low flow pump. All four sites exhibited elevated levels of total metals in groundwater samples. In addition, due to the fact that deep groundwater quality is not significantly impacted with metals indicates that well construction or sampling techniques are probably not factors related to elevated total metals in groundwater.

With respect to past disposal practices, Table 5 clearly shows that soil concentrations do not correlate with elevated total metals in groundwater. Based on this analysis, and on many of the sites previously investigated, the source of total metals in groundwater cannot be attributable to soil contamination or disposal practices in many cases. This is based on both the history of the site as well as the analytical soil results. In some cases, total metals were detected at elevated levels even when the site history did not correlate with the contaminants found. For example, Sites 2 and 21 have a history of pesticide storage and handling, and there are no known disposal areas (i.e., buried debris) within the site boundary. Nevertheless, both of these sites exhibited several metals above drinking water standards that would not be expected to be present at high concentrations based on the historical use of the site. These metals included lead, chromium, beryllium, cadmium, and manganese.

Objective No. 2 (Determine whether total metals in shallow groundwater are elevated throughout the region or MCB Camp Lejeune)

Based on groundwater data obtained from both upgradient wells and off base wells, total metals were detected above drinking water standards in shallow groundwater in areas that would not be influenced by former disposal activities at the sites. Given that some of the upgradient wells are contaminated, it is apparent that total metals in shallow groundwater are elevated in certain areas of the base outside of the influence of site-related disposal activities. However, it is unknown whether the shallow aquifer upgradient of the sites is contaminated due to other base-related activities or whether the levels in groundwater samples are also elevated due to the influence of suspended fines in the samples.

Objective No. 3 (Determine whether there is a correlation between elevated total metals in groundwater and metals in soil)

An evaluation of the data presented in Table 5 shows that metals in soil samples collected in areas of groundwater contamination are not elevated when compared to metals in soil samples collected in areas that did not exhibit groundwater contamination. This supports the possibility that in many cases, elevated levels of total metals in shallow groundwater are not related to the disposal history at the site. As previously mentioned, sites which did not exhibit soil contamination (when compared to background soil levels) or did not have a history of disposal indicative of metals contamination still exhibited elevated levels of total metals in groundwater. Since there is no apparent correlation between metals in soil and total metals in groundwater, then the possibility exists that the elevated total metals in groundwater are biased high due to suspended particulates.

Objective No. 4 (Determine whether the concentrations of total metals in groundwater is related to shallow and deep aquifer characteristics)

There is some evidence that the geologic conditions of the shallow and deep aquifers influence the amount of total metals detected in groundwater samples. The fact that the deep aquifer generally exhibited higher specific conductivity values indicates that there is more dissolved constituents in the deep aquifer when compared to the shallow aquifer. This was evident when comparing Table 1 (total metals in shallow groundwater) to Table 6 (total metals in deep groundwater). Table 6 did not indicate significant levels of total metals in deep groundwater throughout MCB Camp Lejeune.

The geologic conditions of the shallow aquifer would tend to result in samples that may contain suspended particulates. The suspended particulates could influence the total metals concentrations in the samples.

## 6.0 CONCLUSIONS

1. Elevated levels of total metals in the shallow aquifer are probably influenced to some degree by the geologic conditions of the site.
2. There is no correlation between metal levels in soil and total metals in groundwater. Therefore, elevated total metals in groundwater cannot be attributable to soil contamination of past disposal practices.
3. Elevated levels of total metals in the shallow aquifer may be biased high due to suspended particulates in the samples.
4. Dissolved metals in groundwater were generally below Federal MCLs and NC WQS and therefore, do not present a significant problem at MCB Camp Lejeune.
5. Total and dissolved metal concentrations in the Castle Hayne aquifer were generally below drinking water standards and therefore, do not present a significant problem at MCB Camp Lejeune.
6. The presence of manganese in shallow and deep groundwater may be due to naturally occurring geologic conditions.

## **7.0 RECOMMENDATIONS**

- 1. Remediation of total metals in the shallow aquifer at Operable Units 1 and 5 is not recommended based on the following:**
  - Elevated metals in groundwater at both operable units does not appear to be related to soil contamination or past disposal practices;
  - The distribution of total metals in groundwater is not characteristic of a plume that would be present due to a source of contamination;
  - Remediation of total metals would not be practical from an engineering or cost standpoint; and
  - Currently, there is no human or environmental exposure to shallow groundwater.
- 2. Additional background wells should be installed at all sites in order to provide a baseline for comparing on-site groundwater quality.**

## **Tables**

---

---

**TABLE 1**  
**TOTAL METALS BY SITE**  
**SHALLOW MONITORING WELLS**  
**MCB, CAMP LEJEUNE, NORTH CAROLINA**

Site Number Units	NCWQS ug/L	FEDERAL MCL ug/L	Site 1 ug/L	Site 2 ug/L	Site 6 ug/L	Site 7 ug/L	Site 9 ug/L	Site 21 ug/L	Site 24 ug/L	Site 28 ug/L	Site 30 ug/L	Site 41 ug/L	Site 43 ug/L	Site 44 ug/L
Arsenic	50	50	7.2 - 57.4	2.2 - 23.6	ND - 23.3	ND - 43.4J	ND	ND - 101	ND - 116J	5.4 - 13J	6.4 - 12J	2.4 - 36.3	ND - 23.4	ND - 570
Barium	2000	2000	335 - 833	46 - 1420	ND - 1020	427 - 641	ND - 1060	ND - 647	ND - 1120	78.8 - 576	60.1 - 396	55.2 - 999	220 - 745	315 - 3180
Beryllium	NE	4	2.7 J - 43.4	1 - 3	ND - 7.5	ND - 10.3J	ND	ND - 8	ND - 19	ND - 1.2J	ND - 2.4	0.80 - 42.8	1.5 - 4.2	1.4 - 36.6
Cadmium	5	5	ND - 12.9	7	ND	ND	ND	ND	ND - 12	3.3J - 17.3J	ND - 10.7J	3.2 - 110	ND - 6.9	ND - 32
Calcium	NA	NA	8850 - 726000	5710 - 450000	5430 - 64900	5050 - 51300	16100 - 90700	6130J - 63000J	ND - 151000	20200 - 160000	1730 - 11900	8750 - 828000	10300 - 91900	2430 - 191000
Chromium	50	100	172 - 627	11 - 117	ND - 201	47.8 - 220	ND - 214	ND - 348J	19 - 316	9.0J - 140	42.8 - 106J	10.5 - 244	161 - 249	126 - 895
Copper	1000	1300	44.6 - 117	3 - 23	ND - 175	17.7 - 36.4	ND - 39.7	ND - 84	ND - 52	18.8J - 75.4	15.8 - 42.5	16.3 - 1030	64.2 - 104	28.6 - 313
Lead	15	15	40.8J - 176J	2.7 - 44.8	ND - 200	23 - 37.3	ND - 127	ND - 2000J	5.1 - 89	20.3J - 234J	7.7J - 115J	4.8 - 9340	16.5 - 28.8	15.8 - 508
Manganese	50	50 (1)	125 - 1720	21 - 190	ND - 362	56.9 - 220	ND - 91.3	59 - 276J	29 - 518	82.2 - 304	78.5 - 578	56.6 - 2110	72.6 - 297	88 - 1730
Mercury	1.1	2	ND - 1.2J	ND	ND - .46	0.2 - 0.36	ND - 1.4	ND - 2.4J	ND - 3.2	ND - 1.4J	0.88J - 0.9J	0.13 - 0.92	ND - 0.24	ND - 1.1
Nickel	100	100	28.5 - 426	ND	ND - 41.9	ND	ND	ND - 123	ND - 140	ND - 59.8	17.1J - 52.6J	28.8 - 137	20.5 - 143	21.9 - 486
Sodium	NA	NA	9090 - 19000	ND - 103000	1110 - 68700	7040 - 156000	1390 - 4170	7950 - 15700	5230 - 19200	9480 - 74700	5320 - 8100	2080 - 40200	9160 - 22100	4060 - 12600
Vanadium	NE	NE	214 - 640	9 - 184	ND - 330	37.8 - 423	ND - 175	ND - 419	ND - 408	6.1 - 164	57 - 101	20.4 - 244	122 - 233	184 - 759
Zinc	2100	5000 (1)	ND - 1110	6 - 146	ND - 1620	83.6 - 133	ND - 118	27J - 487J	20 - 650	ND	79.2 - 104	25.7 - 5180	19 J - 661J	87.3 - 2800J

Site Number Units	Site 48 ug/L	Site 63 ug/L	Site 65 ug/L	Site 69 ug/L	Site 78 ug/L	Site 82 ug/L	ABC Cleaners ug/L	Offsite Property #1 ug/L	Offsite Property #2 ug/L
Arsenic	ND	ND - 23.4	ND - 308	2.9 - 29.0	ND - 405J	ND - 67.8	ND - 12	10.3 - 160	ND
Barium	18 - 51.3	56.1 - 5410	105 - 638	46.5 - 850	ND - 1250	ND - 540	35 - 220	ND - 468	ND
Beryllium	ND	ND - 3.1	ND	1.3 - 10.6	ND - 19	ND	NA	ND - 8.5	ND
Cadmium	2.2 - 3.3	ND	ND	2.4 - 11.4	ND - 21	ND	NA	ND	ND
Calcium	30600 - 115000	2830 - 24300	33300 - 181000	2010 - 38700	ND - 642000	6580 - 60800	790 - 16000	ND - 22800	ND - 5200
Chromium	5.8 - 17.5	4.4 - 134	50.1 - 364	15.1 - 159	ND - 858J	ND - 174	ND - 57	52.8 - 636	ND - 94
Copper	3.1 - 13.5	10.7 - 126	28.2 - 127	16.2 - 70.8	ND - 699	ND - 29.3	ND - 89	ND - 140	ND
Lead	ND	4.3 J - 369	19.1 - 132	7.8 - 188	ND - 360J	ND - 89	ND - 10	12.3 - 345	6.3 - 62.3
Manganese	38.1 - 585	50.3 - 1020	56.2 - 474	13.0 - 912	26 - 714	26.9 - 283	4 - 44	56 - 973	ND - 60.1
Mercury	0.04 - 0.09	ND - 0.20	ND - 0.29	0.10 - 0.94	ND - 1.5	ND - 0.66	NA	ND	ND
Nickel	ND	19.8 - 54.2	19.4 - 84.3	13.6 - 99.8	ND - 234	ND - 34.6	ND - 77	40.2 - 380	ND
Sodium	5750 - 8760	3150 - 7100	3850 - 11700	4790 - 41300	ND - 42500	5670 - 36500	5800 - 33000	ND - 9390	ND - 7630
Vanadium	3.4 - 12.8	7.9 - 163	59.8 - 433	17.3 - 210	ND - 1700	ND - 236	ND - 45	70 - 739	ND - 64.7
Zinc	ND - 30.3	58.5J - 1110J	148J - 406J	36.2 - 12100	6J - 967J	ND - 204	14 - 220	ND - 736	ND - 40.8

NOTES:

J - Value is estimated.

JB - Value is estimated below the CRDL, but greater than the IDL.

NE - Not established.

NA - Not analyzed.

ND - Not detected.

NCWQS - North Carolina Water Quality Standard

MCL - Maximum Contaminant Level

(1) - Secondary MCL

**TABLE 2**  
**COMPARISON OF REPEAT SAMPLING OF SHALLOW WELLS**  
**MCB, CAMP LEJEUNE, NORTH CAROLINA**

Well Date	2GW01		2GW03		2GW06		2GW08		2GW09	
	5/1993	3/1994	5/1993	3/1994	5/1993	3/1994	5/1993	3/1994	5/1993	3/1994
Chromium	18	ND	11	ND	15	ND	ND	ND	25	83
Lead	15.5 J	ND	3.5 J	ND	6.7 J	ND	ND	3.4	27.2 J	23.6
Manganese	55	47	21	ND	79	140	53	415	290	747

Well Date	78GW05		78GW08		78GW15		78GW16		78GW19	
	1/1991	4/1994	1/1991	4/1994	1/1991	4/1994	1/1991	4/1994	1/1991	4/1994
Chromium	ND	17 J	91.8	491 J	21.4	215 J	209	353 J	13.8	ND
Lead	13.6	13.1 J	54.1	131 J	16.6	53	100	224	31.7	8.3
Manganese	162	161 J	46.5	213 J	18.3	115	98.3	150	79	26

**NOTES:**

J - Value is estimated.

ND - Not detected.

**TABLE 3**  
**DISSOLVED METALS BY SITE**  
**SHALLOW MONITORING WELLS**  
**MCB, CAMP LEJEUNE, NORTH CAROLINA**

Site Number Units	NCWQS ug/L	FEDERAL MCL ug/L	Site 1 ug/L	Site 2 ug/L	Site 6 ug/L	Site 7 ug/L	Site 9 ug/L	Site 21 ug/L	Site 24 ug/L	Site 28 ug/L	Site 30 ug/L	Site 41 ug/L	Site 43 ug/L	Site 44 ug/L
Arsenic	50	50	NA	2.2 - 7.1	ND	NA	ND	ND - 10.6	ND - 16.3	NA	NA	2.2 - 4.7	NA	NA
Barium	2000	2000	NA	25 - 149	ND	NA	ND	ND	ND	NA	NA	12.4 - 451	NA	NA
Beryllium	NE	4	NA	1	ND	NA	ND	ND	ND	NA	NA	0.80 - 3.2	NA	NA
Cadmium	5	5	NA	ND	ND	NA	ND	ND - 5	ND	NA	NA	3.2 - 4.2	NA	NA
Calcium	NA	NA	NA	5800 - 441000	6230 - 57400	NA	15800 - 82400	35900	ND - 113000	NA	NA	4710 - 138000	NA	NA
Chromium	50	100	NA	10	ND	NA	ND	ND	ND	NA	NA	8.3 - 9.6	NA	NA
Copper	1000	1300	NA	2 - 9	ND	NA	ND	ND	ND	NA	NA	16.3 - 23.9	NA	NA
Lead	15	15	NA	2.1	ND	NA	ND	ND - 94	ND	NA	NA	1.0	NA	NA
Manganese	50	50 (1)	NA	17 - 129	ND - 92.7	NA	ND	40 - 134	ND - 320	NA	NA	7.1 - 521	NA	NA
Mercury	1.1	2	NA	ND	ND	NA	ND	ND	ND - 0.5	NA	NA	0.13 - 0.20	NA	NA
Nickel	100	100	NA	ND	ND	NA	ND	ND	ND - 57	NA	NA	28.8 - 31.2	NA	NA
Sodium	NA	NA	NA	ND - 103000	1420 - 70500	NA	1280 - 3860	16200	ND - 183000	NA	NA	2500 - 34200	NA	NA
Vanadium	NE	NE	NA	43	ND	NA	ND	ND	ND	NA	NA	20.4	NA	NA
Zinc	2100	5000 (1)	NA	8 - 35	ND - 350	NA	ND	68 - 50	ND - 437	NA	NA	10.6 - 125	NA	NA

Site Number Units	Site 48 ug/L	Site 63 ug/L	Site 65 ug/L	Site 69 ug/L	Site 78 ug/L	Site 82 ug/L	ABC Cleancers ug/L	Offsite Property #1 ug/L	Offsite Property #2 ug/L
Arsenic	ND	NA	NA	2.9	ND - 21.6	ND	NA	ND - 18.8	ND
Barium	16.8 - 27.6	NA	NA	13.7 - 35.8	ND	ND	NA	ND	ND
Beryllium	ND	NA	NA	1.3	ND	ND	NA	ND	ND
Cadmium	ND - 3.1	NA	NA	2.4	ND	ND	NA	ND	ND
Calcium	72600 - 80700	NA	NA	764 - 10600	ND - 296000	15200 - 58500	NA	ND - 7710	ND
Chromium	ND	NA	NA	7.2	ND - 59	ND	NA	ND - 30.0	ND
Copper	2.6 - 7.6	NA	NA	16.2	ND - 121	ND	NA	ND - 10.7	ND
Lead	ND	NA	NA	1	ND - 17.2	ND	NA	ND - 15.8	ND
Manganese	39.7 - 539	NA	NA	8.5 - 139	ND - 152	21 - 127	NA	ND - 63.8	ND - 21.3
Mercury	0.05 - 0.09	NA	NA	0.1	ND - 0.6	ND	NA	ND	ND
Nickel	ND	NA	NA	13.6	ND	ND	NA	ND	ND
Sodium	6430 - 8920	NA	NA	5170 - 41100	ND - 42200	5980 - 36000	NA	ND - 9540	ND - 6750
Vanadium	ND	NA	NA	16.6	ND	ND	NA	ND	ND
Zinc	ND	NA	NA	7.0 - 7670	ND - 58	ND - 119	NA	ND - 468	ND - 222

**NOTES:**

J - Value is estimated.

JB - Value is estimated below the CRDL, but greater than the IDL.

NE - Not established.

NA - Not analyzed.

ND - Not detected.

NCWQS - North Carolina Water Quality Standard

MCL - Maximum Contaminant Level

(1) - Secondary MCL

**TABLE 4**  
**SUMMARY OF TOTAL METALS IN UPGRAIDENT WELLS**  
**SHALLOW MONITORING WELLS**  
**MCB, CAMP LEJEUNE, NORTH CAROLINA**

Well Number	NCWQS	FEDERAL MCL	Upgradient of Site	Upgradient of Sites	Upgradient of Site									
			1	2	6	7	9	21 and 78	24	28	30	41	43	44
			Units	ug/L	ug/L	ug/L	ug/L	ug/L	ug/L	ug/L	ug/L	ug/L	ug/L	ug/L
Arsenic	50	50	17.8 J	12.9	ND	ND	ND	ND	3.7 J	7.4 J	13.1			
Barium	2000	2000	548	328	257	428	71.3	ND	ND	576	55.7			
Beryllium	NE	4	3.2 J	3	ND	ND	ND	ND	ND	9.3 J	1.6			
Cadmium	5	5	ND	ND	ND	ND	ND	not reported	ND	3.3 J	10			
Chromium	50	100	193	75	198	124	ND	13	37	122	54.4			
Copper	1000	1300	64.8	25	35.6	36.4	ND	ND	ND	20.7 J	27			
Lead	15	15	78.8 J	27.2	64.4	30.3 J	ND	9	11.4	22.4 J	23.7			
Manganese	50	50 (1)	202	747	84.5	56.9 J	ND	ND	39	206	203			
Mercury	1.1	2	1.6 J	ND	ND	0.36	ND	ND	ND	ND	0.16			
Nickel	100	100	51.6	ND	ND	ND	ND	ND	ND	59.8	38			
Vanadium	NE	NE	214	86	209	152	ND	149	64	85.3	38.1			
Zinc	2100	5000 (1)	ND	103	56.6	86.4 J	ND	68.1	41	ND	173			

Well Number	Upgradient of Site	Upgradient of ABC Cleaners	Upgradient of Offsite Property #1	Upgradient of Offsite Property #2				
	48	63	65	69	78	82	MW-S01	
	48GW1			69GW07	9GW04	6MW3S	MW-S01	
Units	ug/L	ug/L	ug/L	ug/L	ug/L	ug/L	ug/L	
Arsenic	ND			2.9	ND	ND	ND	
Barium	29.4 J			46.5	ND	ND	35	
Beryllium	ND			1.3	ND	ND	NA	
Cadmium	2.5 J			2.4	ND	ND	NA	
Chromium	ND			15.8	ND	ND	ND	
Copper	ND			16.2	ND	ND	ND	
Lead	ND			7.8	ND	ND	3	
Manganese	70.6			13	ND	ND	10	
Mercury	ND			0.1	ND	ND	NA	
Nickel	ND			13.6	ND	ND	ND	
Vanadium	3.4 J			17.3	ND	ND	9	
Zinc	ND			36.2	ND	ND	23	

NOTES:

J - Value is estimated.

JB - Value is estimated below the CRDL, but greater than the IDL.

NE - Not established.

NA - Not analyzed.

ND - Not detected.

NCWQS - North Carolina Water Quality Standard

MCL - Maximum Contaminant Level

(1) - Secondary MCL

**TABLE 5**  
**COMPARISON OF INORGANIC SUBSURFACE SOIL CONCENTRATIONS IN "CLEAN" AND "CONTAMINATED" WELLS**  
**MCB, CAMP LEJEUNE, NORTH CAROLINA**

Units Well Number Soil Sample Number	Camp Lejeune Background Subsurface Soil Data mg/kg	Site 1		Site 2		Site 6		Site 7		Site 9		Site 21	
		"Clean" mg/kg	"Contaminated" mg/kg										
		—	—	2-GW07	2-GW09	6-GW18	6-GW15	7-GW03	7-GW02	9-GWS	9-GW1	21-GW03	21-GW02
Arsenic	0.03 - 0.47	NA	NA	1.7 J	ND	ND	ND	1.5	ND	ND	ND	ND	0.55 J
Barium	2 - 11	NA	NA	12.5 J	ND	ND	ND	6.6	71	ND	ND	ND	4.4 J
Beryllium	0.03 - 0.23	NA	NA	ND	ND								
Cadmium	0.17 - 1.2	NA	NA	ND	ND	ND	ND	1.3	4.5	ND	ND	ND	ND
Chromium	2 - 9	NA	NA	10.9 J	4.6	ND	ND	5.2	16	ND	2.6 J	15.2	3.2 J
Copper	0.47 - 2	NA	NA	0.97 J	ND	ND	ND	ND	ND	ND	ND	ND	ND
Lead	1 - 12	NA	NA	8 J	4.3	3.3 J	ND	2.5	34.4	1.6	8.3	7.1	6.9 J
Manganese	0.40 - 8	NA	NA	4.3 J	4.1	ND	1.8 B	3	13	ND	3.7 J	0.9 J	3.4 J
Mercury	0.01 - 0.11	NA	NA	0.3 J	ND	ND	ND	10.13	0.48	ND	ND	ND	ND
Nickel	0.70 - 5.0	NA	NA	ND	ND	ND	ND	3.4	11.8	ND	ND	ND	ND
Vanadium	0.75 - 13	NA	NA	13.8 J	ND	ND	2.9 B	5.5	4.5	ND	ND	15.5	4.4 J
Zinc	0.40 - 12	NA	NA	ND	ND	ND	ND	1.3	ND	ND	6.1 J	5.7	3 J

**NOTES:**

Shaded area indicates inorganic which exceeded a MCL and/or NCWQS in groundwater sample.

J - Value is estimated.

JB - Value is estimated below the CRDL, but greater than the IDL.

NA - No available wells to compare OR compound was not analyzed.

ND - Not detected.

NCWQS - North Carolina Water Quality Standard

MCL - Maximum Contaminant Level

(1) - Secondary MCL.

**TABLE 5**  
**COMPARISON OF INORGANIC SUBSURFACE SOIL CONCENTRATIONS IN "CLEAN" AND "CONTAMINATED" WELLS**  
**MCB, CAMP LEJEUNE, NORTH CAROLINA**

Well Number Soil Sample Number	Site 24		Site 28		Site 30		Site 41		Site 43		Site 44		
	Units	"Clean" mg/kg	"Contaminated" mg/kg										
	24GW10	24GW02	--	--	--	--	41GW04	41-GW11	43GW01	43GW02	44GW02	44GW01	
24-GW10	24-BDA-SB09	--	--	--	--	41-GW04-DW	41-GW11-01	43-GW01-00	43-GW02-00	44-GW02-035	--		
Arsenic	ND	ND	NA	NA	NA	NA	0.51	1.6	ND	ND	ND	1.7	
Barium	ND	ND	NA	NA	NA	NA	9.4	22.6	ND	ND	ND	17.9	
Beryllium	ND	ND	NA	NA	NA	NA	0.18	0.18	ND	ND	ND	ND	
Cadmium	ND	ND	NA	NA	NA	NA	0.73	0.73	8.3	ND	ND	ND	
Chromium	11.2	9.7	NA	NA	NA	NA	3.6	11.2	4.3	6.7	5.6	10.1	
Copper	ND	ND	NA	NA	NA	NA	3.7	22.3	3.4	ND	6.2 J	25.4 J	
Lead	4.6 J	4.4 J	NA	NA	NA	NA	4.8	11.0	9.8	6.1	5.4	10.7	
Manganese	4.7	4.4 J	NA	NA	NA	NA	3.7	75.3	31.2	5.4	5.4	23.4	
Mercury	ND	ND	NA	NA	NA	NA	0.06	0.31	ND	ND	ND	ND	
Nickel	ND	ND	NA	NA	NA	NA	6.6	6.4	7.6	7.1	3.1	3.4	
Vanadium	18.4	10	NA	NA	NA	NA	6.8	9.3	7.2	5.8	5	14.7	
Zinc	ND	7.8	NA	NA	NA	NA	7.7	130	20.1	3	3.2	34.9	

**NOTES:**

Shaded area indicates inorganic which exceeded a MCL and/or NCWQS in groundwater sample.

J - Value is estimated.

JB - Value is estimated below the CRDL, but greater than the IDL.

NA - No available wells to compare OR compound was not analyzed.

ND - Not detected.

NCWQS - North Carolina Water Quality Standard

MCL - Maximum Contaminant Level

(1) - Secondary MCL

**TABLE 5**  
**COMPARISON OF INORGANIC SUBSURFACE SOIL CONCENTRATIONS IN "CLEAN" AND "CONTAMINATED" WELLS**  
**MCB, CAMP LEJEUNE, NORTH CAROLINA**

Units Well Number Soil Sample Number	Site 48		Site 63		Site 65		Site 69		Site 78		Site 82	
	"Clean" mg/kg	"Contaminated" mg/kg										
	48-GW01	48-GW03	63-MW03	63-MW02	65-MW03	65-MW02	69-CW11	69-GW03	78-GW34	78-GW24-1	6-GW28	82-MW3
48-GW1A-01	48-C3-03	63-MW03-04	63-MW02-06	65-MW03-11	65-MW02-06	69-GW11-04	69-CSA-SB23-00	78-GW34	78-B903-SB03	6-GW28-09	6-GW27D-06	
Arsenic	1.3	0.77 J	ND	ND	ND	13	0.68	0.63	ND	ND	0.31	15.9
Barium	21.1	15	ND	ND	3.4	6.8	5.6	3	ND	ND	ND	ND
Beryllium	0.2	0.19	ND	ND	ND	ND	0.3	0.28	ND	ND	ND	ND
Cadmium	1.4	1.8 J	ND	ND	NA	NA	0.56	0.52	ND	ND	ND	ND
Chromium	18.2	18.6	7.7	ND	3.9	3.7	6.8	1.2	18.5	9.3	2.6	5
Copper	3.5	3.8	ND	ND	1.5	3.1	3.8	3.5	3.4 B	ND	ND	ND
Lead	32.3	14.3	4.2	2.5	1.7	1.7	4.3	1.1	4.5 J	2.6 J	2.7	4.1
Manganese	4.1	7	4.9	13.4	3.5	6.9	4	1.2	9.2	ND	ND	ND
Mercury	ND	ND	ND	ND	NA	NA	0.06	0.05	ND	ND	ND	ND
Nickel	2.2	1.9 J	ND	ND	ND	ND	3.2	3	ND	ND	ND	ND
Vanadium	28.3	20.8 J	ND	ND	4.4	3	4.4	3.6	18.7	19.2	ND	ND
Zinc	ND	ND	ND	ND	2.7	5	3.2	3.1	7.9	ND	ND	ND

**NOTES:**

Shaded area indicates inorganic which exceeded a MCL and/or NCWQS in groundwater sample.

J - Value is estimated.

JB - Value is estimated below the CRDL, but greater than the IDL.

NA - No available wells to compare OR compound was not analyzed.

ND - Not detected.

NCWQS - North Carolina Water Quality Standard

MCL - Maximum Contaminant Level

(I) - Secondary MCL

TABLE 5  
COMPARISON OF INORGANIC SUBSURFACE SOIL CONCENTRATIONS IN "CLEAN" AND "CONTAMINATED" WELLS  
MCB, CAMP LEJEUNE, NORTH CAROLINA

Well Number Soil Sample Number	Units	ABC Cleaners		Offsite Property #1		Offsite Property #2	
		"Clean" mg/kg	"Contaminated" mg/kg	"Clean" mg/kg	"Contaminated" mg/kg	"Clean" mg/kg	"Contaminated" mg/kg
		-	-	-	-	-	-
Arsenic	NA	NA	NA	NA	NA	NA	NA
Barium	NA	NA	NA	NA	NA	NA	NA
Beryllium	NA	NA	NA	NA	NA	NA	NA
Cadmium	NA	NA	NA	NA	NA	NA	NA
Chromium	NA	NA	NA	NA	NA	NA	NA
Copper	NA	NA	NA	NA	NA	NA	NA
Lead	NA	NA	NA	NA	NA	NA	NA
Manganese	NA	NA	NA	NA	NA	NA	NA
Mercury	NA	NA	NA	NA	NA	NA	NA
Nickel	NA	NA	NA	NA	NA	NA	NA
Vanadium	NA	NA	NA	NA	NA	NA	NA
Zinc	NA	NA	NA	NA	NA	NA	NA

NOTES:

Shaded area indicates inorganic which exceeded a MCL and/or NCWQS in groundwater sample.

J - Value is estimated.

JB - Value is estimated below the CRDL, but greater than the IDL.

NA - No available wells to compare OR compound was not analyzed.

ND - Not detected.

NCWQS - North Carolina Water Quality Standard

MCL - Maximum Contaminant Level

(1) - Secondary MCL

TABLE 6  
TOTAL METALS BY SITE  
DEEP MONITORING WELLS  
MCB, CAMP LEJEUNE, NORTH CAROLINA

	Site 1	Site 2	Site 6	Site 7	Site 9	Site 21	Site 24	Site 28	Site 30	Site 41	Site 43	Site 44	Site 48	Site 63	Site 65	Site 69	Site 78	Site 82	ABC Cleaners	Base Supply Wells (1)
Arsenic			ND			ND				2.2 - 9.6						2.2 - 3.5	2 - 118 J	ND	ND - 14	ND
Barium		1420	ND			ND				22.6 - 186						42.3 - 58.0	ND - 547	ND	4 - 36	ND
Beryllium		ND	ND			ND				3.2						0.80 - 0.89	ND	ND	NA	NA
Cadmium		ND	ND			ND				4.2 - 4.7						3.2	ND - 21	ND	NA	ND
Chromium		16	ND			ND				9.6 - 40.5						8.3 - 20.7	ND - 10	ND	ND - 32	ND
Copper		ND	ND			ND				23.9						16.3	ND	ND	ND - 41	ND - 130
Lead		ND	ND			ND				1.0 - 11.1						3.1 - 6.8	ND	ND	ND - 10	ND - 16
Manganese			ND - 33.5			ND				16.9 - 101						53.7 - 114	ND - 591	ND - 21.6	ND - 45	10 - 120
Mercury		ND	ND			ND				0.15 - 0.17						0.16 - 0.17	ND - 0.3	ND	NA	ND
Nickel		ND	ND			ND				31.2						28.8	ND	ND	ND - 14	NA
Vanadium		ND	ND			ND				20.4 - 49.8						20.4	ND - 24 J	ND	ND - 15	NA
Zinc		ND	ND			ND				17.8 - 83.8						31.1 - 48.7	ND - 181 J	ND	58 - 390	ND - 120

NOTES:

J - Value is estimated.

NA - Not analyzed.

ND - Not detected.

(1) - Range is based on 67 supply wells located throughout MCB, Camp Lejeune, NC.

**TABLE 7**  
**SUMMARY OF FIELD PARAMETERS IN**  
**SHALLOW, DEEP, AND SUPPLY WELLS**  
**MCB; CAMP LEJEUNE, NORTH CAROLINA**

	Shallow Wells		Deep Wells		Supply Wells	
	Range (1)	Average Maximum	Range (2)	Average Maximum	Range (3)	Average Maximum
pH (standard units)	4.5 - 7.28	6.08	7.52 - 11.34	8.88	6.91 - 7.45	7.32
Specific Conductivity (micromhos/cm)	40 - 580	267	149 - 525	350	212 - 511	353

(1) - Based on data from 11 sites.

(2) - Based on data from 6 sites.

(3) - Based on data from 9 supply wells.

## **Figures**

---






FIGURE 1  
SITE LOCATION MAP  
INORGANIC GROUNDWATER STUDY


MARINE CORPS BASE, CAMP LEJEUNE  
NORTH CAROLINA

01508TB1Z





**FIGURE 3**  
POSITIVE DETECTIONS OF TAL METALS ABOVE  
FEDERAL MCLs AND/OR NCWQS IN SHALLOW WELLS  
SITE 78  
REMEDIAL INVESTIGATION CTO-0177  
MARINE CORPS BASE, CAMP LEJEUNE  
NORTH CAROLINA



**LEGEND**

78GW04-3 EXISTING DEEP MONITORING WELL INSTALLED BY ESE, 1991  
 → APPROXIMATE DIRECTION OF GROUNDWATER FLOW  
 (NS) NOT SAMPLED FOR TAL METALS  
 HP-603 WATER SUPPLY WELL (ACTIVE)-NOT SAMPLED  
 HP-601 WATER SUPPLY WELL (INACTIVE)-NOT SAMPLED  
 SOURCE: LANTDIV, FEBRUARY 1992

**FIGURE 4**  
**POSITIVE DETECTIONS OF TAL METALS ABOVE FEDERAL MCLs AND/OR NCWQS IN DEEP WELLS SITE 78**  
**REMEDIAL INVESTIGATION CTO-0177**  
**MARINE CORPS BASE, CAMP LEJEUNE**  
**NORTH CAROLINA**

**Appendix A  
Data Summary Tables  
for Sites 2 and 78**

---

---

**OPERABLE UNIT NO. 1 - SITES 21, 24, 78**  
**SHALLOW, INTERMEDIATE AND DEEP MONITORING WELLS**  
**GROUNDWATER DATA AND FREQUENCY SUMMARY**  
**REMEDIAL INVESTIGATION CTO - 19177**  
**MCB CAMP LEJEUNE, NORTH CAROLINA**  
**TAL METALS AND CYANIDE**

	MINIMUM NONDETECTED UG/L	MAXIMUM NONDETECTED UG/L	MINIMUM DETECTED UG/L	MAXIMUM DETECTED UG/L	LOCATION OF MAXIMUM DETECTED	FREQUENCY OF DETECTION
ALUMINUM	NA	NA	68 J	542000 J	78-GW06-01	59 / 59
ANTIMONY	3 U	20 U	3.3 B	169 J	78-GW02-01	7 / 33
ARSENIC	2 U	10 U	2.3 J	405 J	78-GW02-01	44 / 48
BARIUM	NA	NA	17 B	1250	78-GW07-01	59 / 59
BERYLLIUM	1 U	4 U	1 B	19	24-GW02-01	52 / 59
CADMIUM	5 U	25 U	5	21	78-GW04-3-01	9 / 59
CALCIUM	NA	NA	2420 B	642000	78-GW04-1-01	59 / 59
CHROMIUM	10 U	50 U	10	858 J	78-GW06-01	46 / 59
COBALT	8 U	8 U	8 B	170	78-GW22-2-01	25 / 59
COPPER	2 U	2 U	3 B	699	78-GW39-01	58 / 59
IRON	NA	NA	32 B	523000	78-GW04-3-01	59 / 59
LEAD	1.8 U	4.9 U	2.9 B	2000 J	21-GW0B-01	50 / 59
MAGNESIUM	NA	NA	88 B	37100	24-GW03-01	59 / 59
MANGANESE	2 U	2 U	2 B	714	78-GW24-1-01	57 / 59
MERCURY	0.2 U	0.2 U	0.23 J	3.2	24-GW06-01	24 / 52
NICKEL	20 U	20 U	20 B	234	78-GW22-2-01	31 / 59
POTASSIUM	NA	NA	982 B	67300	78-GW32-3-01	59 / 59
SELENIUM	1 U	5 U	1.1 J	99.5 J	78-GW32-2-01	41 / 54
SILVER	3 U	15 U	5 J	5 J	78-GW09-3-01	1 / 59
SODIUM	NA	NA	2450 B	42500	78-GW32-3-01	59 / 59
THALLIUM	1 U	1 U	1 B	7.3 J	78-GW32-2-01	16 / 59
VANADIUM	4 U	4 U	4 J	1700	78-GW08-01	55 / 59
ZINC	6 U	6 U	6 J	967 J	78-GW22-2-01	57 / 59
CYANIDE	10 U	10 U	ND	ND	ND	0 / 54

OPERABLE UNIT NO. 1 - SITES 21, 24, 78  
 SHALLOW, INTERMEDIATE AND DEEP MONITORING WELLS  
 GROUNDWATER DATA AND FREQUENCY SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19177  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 TAL METALS AND CYANIDE

SAMPLE NO. UNITS	21-GW01-01 UG/L	21-GW02-01 UG/L	21-GW03-01 UG/L	21-GW04-01 UG/L	21-GW0A-01 UG/L	21-GW0B-01 UG/L
ALUMINUM	4910 J	319000 J	4820 J	20100 J	16900 J	118000 J
ANTIMONY	7 UJ	7 U	7 U	7 U	7 R	7 U
ARSENIC	15	10	2 U	11.8	45.2 J	30.4
BARIUM	32 B	647	51 B	119 B	100 B	386
BERYLLIUM	1 B	5	1 B	1 B	1 B	6
CADMUM	5 U	10 U	5 U	5 U	5 U	10 U
CALCIUM	63000 J	24100 J	6130 J	21700 J	23800	6250 J
CHROMIUM	10 UJ	348 J	10 UJ	33 J	21 J	192 J
COBALT	8 U	18 B	8 U	10 B	8 U	36 B
COPPER	4 B	79	7 B	28	24 B	38
IRON	9920 J	122000 J	13400 J	24900 J	38900 J	72900 J
LEAD	1.8 UJ	214 J	4.9 UJ	33 J	29	2000 J
MAGNESIUM	5070	15400	4550 B	5490	4850 B	11600
MANGANESE	64 J	179 J	134 J	193 J	59	276 J
MERCURY	0.2 R	2.4 J	0.2 R	0.2 R	0.2 U	0.2 R
NICKEL	20 U	86	20 U	20 U	20 U	60
POTASSIUM	2390 B	10500	2240 B	3800 B	2360 B	9520
SELENIUM	1 U	11 J	1 U	1 U	1 UJ	3.7 J
SILVER	3 U	3 U	3 U	3 U	3 UJ	3 U
SODIUM	15700	12600	7950	14400	12600	14400
THALLIUM	1 U	1 UJ	1 U	1 UJ	1 UJ	1 U
VANADIUM	30 B	281	11 B	42 B	48 B	243
ZINC	65 J	136 J	27 J	57 J	41 J	175 J
<u>CYANIDE</u>	10 U					

OPERABLE UNIT NO. 1 - SITES 21, 24, 78  
 SHALLOW, INTERMEDIATE AND DEEP MONITORING WELLS  
 GROUNDWATER DATA AND FREQUENCY SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19177  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 TAL METALS AND CYANIDE

SAMPLE NO. UNITS	21-GW0C-01 UG/L	24-GW01-01 UG/L	24-GW02-01 UG/L	24-GW03-01 UG/L	24-GW04-01 UG/L	24-GW06-01 UG/L
ALUMINUM	209000 J	262000	93700	50200	58900	19800
ANTIMONY	.7 U	3 U	3 UJ	3 U	4.6 B	3.5 B
ARSENIC	101	10 UJ	2.3 J	4.7 J	116 J	10.1 J
BARIUM	467	380	1120	480	290	159 B
BERYLLIUM	8	3 B	19	5	2 B	9
CADMUM	10 U	5 U	12	5 U	5 U	5
CALCIUM	35200 J	4120 B	2420 B	124000	65600	151000
CHROMIUM	291 J	296	316	110	153	78
COBALT	60	8 U	41 B	66	8 U	35 B
COPPER	84	49	52	22 B	31	15 B
IRON	106000 J	58600	395000	16300	70500	69500
LEAD	92.5 J	89	17.9	21.6	23.6	7.4
MAGNESIUM	16300	12200	7240	37100	7690	4320 B
MANGANESE	273 J	117	518	393	66	431
MERCURY	0.23 J	0.23	2.6	0.2 U	0.2 U	3.2
NICKEL	123	38 B	140	85	20 U	93
POTASSIUM	11800	12000	7550	15400	6130	3370 B
SELENIUM	4.3 B	1.3 J	1.1 J	16.2 J	4.3 J	1 UJ
SILVER	3 U	3 UJ	15 UJ	3 UJ	3 UJ	3 UJ
SODIUM	15200	6030	11600	19200	5230	7280
THALLIUM	1 U	1 U	1 U	2.4 B	1 U	1 B
VANADIUM	419	304	408	92	202	83
ZINC	487 J	118	461	650	80	489
CYANIDE	10 U					

OPERABLE UNIT NO. 1 - SITES 21, 24, 78  
 SHALLOW, INTERMEDIATE AND DEEP MONITORING WELLS  
 GROUNDWATER DATA AND FREQUENCY SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19177  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 TAL METALS AND CYANIDE

SAMPLE NO. UNITS	24-GW07-01 UG/L	24-GW08-01 UG/L	24-GW09-01 UG/L	24-GW10-01 UG/L	78-GW02-01 UG/L	78-GW03-01 UG/L
ALUMINUM	36000	61100	12800	23300	29200 J	23900 J
ANTIMONY	3 U	3 U	3.3 B	5.7 B	169 J	38.5 J
ARSENIC	3.7 J	8 J	4.3 J	2.5 J	405 J	5.7 J
BARIUM	85 B	112 B	164 B	59 B	109 B	36 B
BERYLLIUM	1 B	2 B	1 B	1 U	12	2 B
CADMIUM	5 U	5 U	5 U	5 U	8	5 U
CALCIUM	4960 B	27000	9530	3820 B	37000	32900
CHROMIUM	37	85	19	21	18 J	10 UJ
COBALT	8 U	8 U	11 B	8 U	8 U	8 U
COPPER	19 B	24 B	11 B	13 B	20 B	8 B
IRON	13700	27500	13100	7010	427000 J	5020 J
LEAD	11.4	23.8	5.1	7.3	19.6	3.4
MAGNESIUM	2670 B	5050	7630	1760 B	3650 B	2210 B
MANGANESE	39	47	180	29	141	27
MERCURY	0.2 U					
NICKEL	20 U					
POTASSIUM	3870 B	5580	4280 B	2620 B	2770 B	1320 B
SELENIUM	2.1 J	1.9 J	2.6 J	1 UJ	19.8 J	2.4 J
SILVER	3 UJ	3 UJ	3 UJ	3 UJ	15 UJ	3 UJ
SODIUM	6520	6550	6010	6650	5120	4270 B
THALLIUM	1 U	1 U	1 U	1 U	1 UJ	1 UJ
VANADIUM	64	129	26 B	34 B	1660	50
ZINC	41	47	50	20	58 J	12 J
CYANIDE	10 U					

OPERABLE UNIT NO. 1 - SITES 21, 24, 78  
 SHALLOW, INTERMEDIATE AND DEEP MONITORING WELLS  
 GROUNDWATER DATA AND FREQUENCY SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19177  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 TAL METALS AND CYANIDE

SAMPLE NO.	78-GW04-1-01	78-GW04-2-01	78-GW04-3-01	78-GW05-01	78-GW06-01	78-GW07-01
UNITS	UG/L	UG/L	UG/L	UG/L	UG/L	UG/L
ALUMINUM	297000 J	286	115 B	23000 J	542000 J	207000 J
ANTIMONY	7 R	7 R	7 R	7 U	7 U	7 U
ARSENIC	18.6 J	2 R	118 J	5.2 J	26 B	16.2
BARIUM	728	519	547	54 B	1200	1250
BERYLLIUM	19	1 B	1 B	2 B	9	5
CADMUM	12	5 U	21	5 U	5 U	5 U
CALCIUM	642000	170000	105000	90200 J	7180 J	18700 J
CHROMIUM	496 J	10 U	50 U	17 J	858 J	400 J
COBALT	28 B	8 U	8 U	8 U	11 B	20 B
COPPER	87	4 B	7 B	8 B	127	53
IRON	267000 J	32 B	523000	14900 J	142000 J	96700 J
LEAD	126	2 U	2 U	13.1 J	155 J	61.5 J
MAGNESIUM	25500	88 B	3210 B	12700	24000	20000
MANGANESE	703	51	591	161 J	184 J	135 J
MERCURY	0.75	0.2 U	0.3	0.2 R	1.1 J	0.44 J
NICKEL	136	20 B	20 U	20 U	86	54
POTASSIUM	18800	21800	11300	4770 B	25600	13200
SELENIUM	9 J	1 R	1 R	6.4	5.5 B	9.1
SILVER	6 UJ	3 U	15 U	3 U	3 U	3 U
SODIUM	8870	11500	9290	23900	5090	9260
THALLIUM	1.2 J	1 U	1 U	1 UJ	1.1 B	1 UJ
VANADIUM	591	4 UJ	24 J	28 B	811	406
ZINC	373 J	7 J	79 J	32 J	223 J	158 J
CYANIDE	10 U	10 U	10 U	10 U	10 U	10 U

OPERABLE UNIT NO. 1 - SITES 21, 24, 78  
 SHALLOW, INTERMEDIATE AND DEEP MONITORING WELLS  
 GROUNDWATER DATA AND FREQUENCY SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19177  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 TAL METALS AND CYANIDE

SAMPLE NO. UNITS	78-GW08-01 UG/L	78-GW09-2-01 UG/L	78-GW09-3-01 UG/L	78-GW10-01 UG/L	78-GW11-01 UG/L	78-GW12-01 UG/L
ALUMINUM	483000 J	68 J	2710 J	404000 J	332000	108000 J
ANTIMONY	7 U	7 R	7 R	7 R	7 R	7 R
ARSENIC	60.3	2 R	2 R	43 J	10 R	9.6 J
BARIUM	740	27 B	41 B	582	631	155 B
BERYLLIUM	9	1 U	1 B	8	5	2 B
CADMUM	25 U	5 U	5 U	10 U	25 U	10 U
CALCIUM	28200 J	114000	99100	54400	9130	31200
CHROMIUM	491 J	10 UJ	10 UJ	362 J	412	114 J
COBALT	29 B	8 U	8 U	31 B	8 U	8 U
COPPER	86	4 B	4 B	91	84	30
IRON	138000 J	955 J	99 J	157000 J	120000	26400 J
LEAD	131 J	2 U	2 U	257	195	35.5
MAGNESIUM	18500	2550 B	249 B	17400	15400	7220
MANGANESE	213 J	19	2 U	326	174	47
MERCURY	1.3 J	0.2 U	0.2 U	1.5	0.75	0.2 U
NICKEL	89	20 U	20 U	108	79	20 U
POTASSIUM	14700	1220 B	7820	15800	13000	6090
SELENIUM	25.3	1 UJ	1 UJ	18 J	12 J	3.6 J
SILVER	3 U	3 UJ	5 J	3 UJ	3 U	3 UJ
SODIUM	4710 B	5820	7280	3340 B	3490 B	5420
THALLIUM	1.3 J	1 UJ	1 UJ	1 UJ	1 U	1 UJ
VANADIUM	1700	4 U	9 B	499	526	145
ZINC	200 J	11 J	181 J	217 J	120 J	64 J
CYANIDE	10 U	10 U	10 U	10 U	10 U	10 U

OPERABLE UNIT NO. 1 - SITES 21, 24, 78  
 SHALLOW, INTERMEDIATE AND DEEP MONITORING WELLS  
 GROUNDWATER DATA AND FREQUENCY SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19177  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 TAL METALS AND CYANIDE

SAMPLE NO.	78-GW13-01 UG/L	78-GW14-01 UG/L	78-GW15-01 UG/L	78-GW16-01 UG/L	78-GW17-1-01 UG/L	78-GW17-2-01 UG/L
UNITS						
ALUMINUM	61800 J	103000 J	205000 J	341000 J	168000 J	541 J
ANTIMONY	7 U	7 R	7 R	7 R	7 R	7 R
ARSENIC	38.3	18.4 J	4 R	19 J	11.6 J	2 R
BARIUM	236	321	469	511	261	57 B
BERYLLIUM	3 B	1 B	4 B	6	4 B	1 B
CADMUM	5 U	10 U	5 U	5 U	10 U	5 U
CALCIUM	4040 J	5300	29100	62700	86900	144000
CHROMIUM	222 J	113 J	215 J	353 J	200 J	10 UJ
COBALT	20 B	8 U	9 B	13 B	9 B	8 U
COPPER	18 B	33	49	80	40	5 B
IRON	61800 J	49600 J	43300 J	80900 J	48700 J	2120 J
LEAD	26.4 J	63	53	224	81	5.9
MAGNESIUM	11800	10600	13400	10800	9940	2570 B
MANGANESE	57 J	68	115	150	96	33
MERCURY	0.3 J	0.38	0.2 U	0.38	0.2 U	0.2 U
NICKEL	40	34 B	29 B	61	30 B	20 U
POTASSIUM	8210	6460	12000	14000	11600	1630 B
SELENIUM	4.7 B	12.4 J	2.1 J	14.5 J	5 UJ	1 UJ
SILVER	3 U	3 UJ	3 UJ	3 UJ	3 UJ	3 UJ
SODIUM	15000	15400	6410	4120 B	3180 B	9480
THALLIUM	1 U	1 UJ	1 J	1.4 J	1 J	1 UJ
VANADIUM	158	122	248	371	289	4 U
ZINC	96 J	51 J	116 J	157 J	98 J	6 UJ
CYANIDE	10 U	10 U				

OPERABLE UNIT NO. 1 - SITES 21, 24, 78  
 SHALLOW, INTERMEDIATE AND DEEP MONITORING WELLS  
 GROUNDWATER DATA AND FREQUENCY SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19177  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 TAL METALS AND CYANIDE

	SAMPLE NO. UNITS	78-GW19-01 UG/L	78-GW20-01 UG/L	78-GW21-01 UG/L	78-GW22-01 UG/L	78-GW22-1-01 UG/L	78-GW22-2-01 UG/L
ALUMINUM		4110 J	149000 J	23800 J	78900 J	257000	190000 J
ANTIMONY		7 R	7 U	7 U	14 J	7 R	7 UJ
ARSENIC		3.1 J	30.3	6.3 J	10 J	59.5 J	75.6
BARIUM		101 B	430	382	107 B	411	471
BERYLLIUM		1 B	4 B	2 B	1 B	4 B	12
CADMUM		5 U	5 U	5 U	10 U	25 U	6
CALCIUM		3700 B	5450 J	32900 J	90100	44500	118000 J
CHROMIUM		10 UJ	231 J	22 J	83 J	238	389 J
COBALT		8 U	35 B	10 B	8 U	8 U	170
COPPER		3 B	61	11 B	34	54	92
IRON		8500 J	101000 J	26400 J	27600 J	62300	140000 J
LEAD		8.3	119 J	19.1 J	37.2	272	360 J
MAGNESIUM		5740	13100	9110	5500	12000	13000
MANGANESE		26	93 J	85 J	70	158	348 J
MERCURY		0.2 U	0.37 J	0.2 R	0.3	0.45	0.2 R
NICKEL		20 U	75	20 U	21 B	99	234
POTASSIUM		2130 B	9100	4100 B	6180	12000	10200
SELENIUM		1 UJ	4.2 B	1.1 B	4.2 J	7.5 J	45
SILVER		3 UJ	3 U	3 U	3 UJ	3 U	3 U
SODIUM		24000	11900	9480	12100	9910	8230
THALLIUM		1 UJ	1.8 B	1 U	1.7 J	1 U	3 B
VANADIUM		9 B	236	86	114	269	547
ZINC		6 J	250 J	108 J	50 J	150 J	967 J
CYANIDE		10 U	10 U				

OPERABLE UNIT NO. 1 - SITES 21, 24, 78  
 SHALLOW, INTERMEDIATE AND DEEP MONITORING WELLS  
 GROUNDWATER DATA AND FREQUENCY SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19177  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 TAL METALS AND CYANIDE

SAMPLE NO. UNITS	78-GW23-01 UG/L	78-GW24-1-01 UG/L	78-GW24-2-01 UG/L	78-GW24-3-01 UG/L	78-GW25-01 UG/L	78-GW29-01 UG/L
ALUMINUM	111000 J	160000	1340	304	101000 J	78800 J
ANTIMONY	7 R	7 R	7 R	7 R	7 R	7 R
ARSENIC	7.6 J	100 J	2 R	2 R	11.4 J	19 J
BARIUM	230	396	34 B	17 B	119 B	1070
BERYLLIUM	2 B	7	1 B	1 U	2 B	12
CADMIUM	5 U	5 U	5	5	5 U	5 U
CALCIUM	10800	34400	107000	73400	37800	41600
CHROMIUM	101 J	264	10	10 U	82 J	252 J
COBALT	8 B	39 B	8 U	8 U	8 U	17 B
COPPER	25	71	6 B	5 B	26	34
IRON	30800 J	159000	2320	2370	26300 J	125000 J
LEAD	50	152	3.3	2.9 B	30.5	25.5
MAGNESIUM	7110	11600	1740 B	1500 B	4500 B	21900
MANGANESE	87	714	21	41	33	341
MERCURY	0.3	0.75	0.2 U	0.2 U	0.2 U	0.2 U
NICKEL	42	91	20 U	20 U	20 U	125
POTASSIUM	5450	9090	1050 B	982 B	4950 B	11600
SELENIUM	4.4 J	17.6 J	1 R	1 R	1.6 J	2.5 J
SILVER	3 UJ	3 U	3 U	3 U	3 UJ	3 UJ
SODIUM	7450	10800	8350	7050	16400	21200
THALLIUM	1.7 J	1.5 B	1 U	1 U	1.3 J	1 UJ
VANADIUM	108	436	4 J	4 UJ	144	183
ZINC	67 J	291 J	11 J	16 J	34 J	330 J
CYANIDE	10 U	10 U	10 U	10 U	10 U	10 U

OPERABLE UNIT NO. 1 - SITES 21, 24, 78  
 SHALLOW, INTERMEDIATE AND DEEP MONITORING WELLS  
 GROUNDWATER DATA AND FREQUENCY SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19177  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 TAL METALS AND CYANIDE

SAMPLE NO.	78-GW31-2-01 UNITS	78-GW31-3-01 UG/L	78-GW32-2-01 UG/L	78-GW32-3-01 UG/L	78-GW33-01 UG/L	78-GW34-01 UG/L
ALUMINUM		110 B	1200	112000 J	539 J	78200
ANTIMONY		7 R	7 R	7 R	7 R	3 U
ARSENIC		2 R	2 R	21.6 J	2 R	5.6 J
BARIUM		17 B	415	476	42 B	162 B
BERYLLIUM		1 B	1 B	10	1 B	1 B
CADMIUM		5 U	5 U	10	5 U	5 U
CALCIUM	77600		308000	94600	5440	64800
CHROMIUM	10 U		21	215 J	10 UJ	65
COBALT	8 U		8 U	84	8 U	8 U
COPPER	3 B		5 B	87	2 U	20 B
IRON	280		72 B	98500 J	112 J	14900
LEAD	2 U		2 U	146	2 U	18.1
MAGNESIUM	2200 B		151 B	13700	319 B	7290
MANGANESE	8 B		2 B	328	2 U	86
MERCURY	0.3		0.2 U	0.3	0.2 U	0.2 U
NICKEL	20 U		20 U	166	20 U	20 B
POTASSIUM	1640 B		61600	8460	67300	6900
SELENIUM	1 R		1.7 J	99.5 J	1 UJ	12.8 J
SILVER	3 U		3 U	3 UJ	3 UJ	3 UJ
SODIUM	10400		26100	7510	42500	7030
THALLIUM	1 U		1 UJ	7.3 J	1.3 J	1 U
VANADIUM	4 J		10 J	462	5 B	74
ZINC	23 J		10 J	826 J	6 UJ	37
CYANIDE	10 U		10 U	10 U	10 U	10 U

OPERABLE UNIT NO. 1 - SITES 21, 24, 78  
 SHALLOW, INTERMEDIATE AND DEEP MONITORING WELLS  
 GROUNDWATER DATA AND FREQUENCY SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19177  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 TAL METALS AND CYANIDE

SAMPLE NO.	78-GW35-01	78-GW36-01	78-GW37-01	78-GW38-01	78-GW39-01
UNITS	UG/L	UG/L	UG/L	UG/L	UG/L
ALUMINUM	47100	120000	73500	102000	60000
ANTIMONY	3 U	20 U	3 U	20 U	20 U
ARSENIC	2 UJ	3.1 J	4 J	33.6 J	4 UJ
BARIUM	261	152 B	123 B	420	256
BERYLLIUM	1 B	2 U	2 B	4 U	1 U
CADMUM	5 U	5 U	5 U	25 U	5 U
CALCIUM	7480	35400	10100	62200	16800
CHROMIUM	55	111	65	201	60
COBALT	8 U	8 U	8 U	8 U	10 B
COPPER	15 B	29	22 B	110	699
IRON	11800	21200	18800	67500	28800
LEAD	13.2	30.2	21.8	41.2	186
MAGNESIUM	5680	5740	4600 B	17500	14300
MANGANESE	57	62	62	106	84
MERCURY	0.2 U	0.3	0.2 U	0.2 U	0.52
NICKEL	20 U	24 B	20 U	32 B	32 B
POTASSIUM	6150	5820	5990	8180	3840 B
SELENIUM	3.5 J	1.7 J	1.1 J	1.3 J	4.3 J
SILVER	3 UJ				
SODIUM	10300	2450 B	7270	10300	19500
THALLIUM	1 U	1 U	1 U	1 U	1 U
VANADIUM	59	98	106	235	67
ZINC	30	57	58	134	138
CYANIDE	10 U				

**OPERABLE UNIT NO. 5 - SITE 2**  
**SHALLOW AND DEEP MONITORING WELLS**  
**GROUNDWATER STATISTICAL SUMMARY**  
**REMEDIAL INVESTIGATION CTO - 19174**  
**MCB CAMP LEJEUNE, NORTH CAROLINA**  
**TAL METALS AND CYANIDE**

SAMPLE NO.	2-GW01-01	2-GW02-01	2-GW03-01	2-GW03DW-01	2-GW04-01	2-GW05-01
UNITS	UG/L		UG/L	UG/L	UG/L	UG/L
ALUMINUM	36000		5200	269	16800	4050
ANTIMONY	10 U		10 U	3.5 U	10 U	10 U
ARSENIC	21.2		2.5 B	1 UJ	23.6	2.2 B
BARIUM	52 B		46 B	1420	95 B	100 B
BERYLLIUM	1 B		0.5 U	0.5 U	2 B	0.5 U
CADMIUM	7		2.5 U	2.5 U	2.5 U	2.5 U
CALCIUM	23700		8460	450000	11100	21000
CHROMIUM	18		11	16	5 U	5 U
COBALT	10 B		4 U	4 U	4 U	4 U
COPPER	10 B		4 B	8 B	5 B	3 B
IRON	10300		7190	127	28100	12700
LEAD	15.5 L		3.5 J	1.1 UJ	2.7 J	0.5 UJ
MAGNESIUM	5660		1600 B	75 B	1920 B	4800 B
MANGANESE	35		21	2 U	21	46
MERCURY	0.1 U		0.1 U	0.1 U	0.1 U	0.1 U
NICKEL	10 U		10 U	10 U	10 U	10 U
POTASSIUM	2560 B		1030 B	187000	1210 B	2130 B
SELENIUM	4.2 B		0.5 U	0.5 U	0.5 U	0.5 U
SILVER	1.5 U		1.5 U	1.5 U	1.5 U	1.5 U
SODIUM	4040 B		5490	103000	5560	10100
THALLIUM	0.5 U		0.5 U	0.5 UJ	0.5 U	0.5 U
VANADIUM	72		10 B	2 U	89	9 B
ZINC	146		13 B	9 B	16 B	6 B
CYANIDE	5 U		5 U	5 U	5 U	5 U

OPERABLE UNIT NO. 5 - SITE 2  
 SHALLOW AND DEEP MONITORING WELLS  
 GROUNDWATER STATISTICAL SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19174  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 TAL METALS AND CYANIDE

SAMPLE NO. UNITS	2-GW06-01 UG/L	2-GW07-01 UG/L	2-GW08-01 UG/L	2-GW09-01 UG/L
ALUMINUM	13600	8550	6380	56300
ANTIMONY	10 U	10 U	3.5 UJ	10 U
ARSENIC	5.4 B	5.7 B	9.2 B	12.9
BARIUM	173 B	98 B	98 B	328
BERYLLIUM	0.5 U	0.5 U	0.5 U	3 B
CADMIUM	2.5 U	2.5 U	2.5 U	2.5 U
CALCIUM	7940	9350	5710	22100
CHROMIUM	15	15	5 U	75
COBALT	12 B	4 U	4 U	10 B
COPPER	5 B	7 B	6 B	25
IRON	11700	12500	9150	42000
LEAD	6.7 J	8.3 J	1.8 UJ	27.2 L
MAGNESIUM	4120 B	3620 B	2020 B	9980
MANGANESE	79	72	53	290
MERCURY	0.1 U	0.1 U	0.1 U	0.1 U
NICKEL	10 U	10 U	10 U	25 B
POTASSIUM	2570 B	1940 B	1550 B	6610
SELENIUM	0.5 U	0.5 U	0.5 U	0.5 U
SILVER	1.5 U	1.5 U	1.5 U	1.5 U
SODIUM	21900	8180	11800	18300
THALLIUM	0.5 U	0.5 U	0.5 U	0.5 U
VANADIUM	15 B	18 B	12 B	86
ZINC	26	22	27	103
CYANIDE	5 U	5 U	5 U	5 U

OPERABLE UNIT NO. 5 - SITE 2  
 SHALLOW AND DEEP MONITORING WELLS  
 GROUNDWATER STATISTICAL SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19174  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 DISSOLVED METALS

SAMPLE NO. UNITS	2-GW01D-01 UG/L	2-GW02D-01	2-GW03D-01 UG/L	2-GW03DWD-01 UG/L	2-GW04D-01 UG/L	2-GW05D-01 UG/L
ALUMINUM	1930		66 B	89 B	60 B	1990
ANTIMONY	10 U		10 U	3.5 U	10 U	10 U
ARSENIC	2.2 B		1 U	1 U	6.1 B	1 U
BARIUM	42 B		25 B	1400	64 B	98 B
BERYLLIUM	1 B		0.5 U	0.5 U	0.5 U	1 B
CADMUM	2.5 U		2.5 U	2.5 U	2.5 U	2.5 U
CALCIUM	24400		7100	441000	11300	21800
CHROMIUM	5 U		5 U	11	5 U	5 U
COBALT	4 U		4 U	4 U	4 U	4 U
COPPER	4 B		2 B	6 B	9 B	4 B
IRON	2560		2170	10 U	2720	7400
LEAD	2.1 J		0.5 UJ	0.5 UJ	0.5 UJ	0.5 UJ
MAGNESIUM	5220		1030 B	26 B	1840 B	4900 B
MANGANESE	51		4.5 U	1 U	17	46
MERCURY	0.1 U		0.1 U	0.1 U	0.1 U	0.1 U
NICKEL	10 U		10 U	10 U	10 U	10 U
POTASSIUM	2140 B		589 B	188000	1130 B	2170 B
SELENIUM	0.5 U		0.5 U	0.5 U	0.5 U	0.5 U
SILVER	1.5 U		1.5 U	1.5 U	1.5 U	1.5 U
SODIUM	3590 B		5400	103000	5710	9970
THALLIUM	0.5 U		0.5 U	0.5 U	0.5 U	0.5 U
VANADIUM	2 U		2 U	2 U	2 U	2 U
ZINC	28		3 U	3 U	8 B	9 B
<u>CYANIDE</u>						

OPERABLE UNIT NO. 5 - SITE 2  
 SHALLOW AND DEEP MONITORING WELLS  
 GROUNDWATER STATISTICAL SUMMARY  
 REMEDIAL INVESTIGATION CTO - 19174  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 DISSOLVED METALS

SAMPLE NO.	2-GW06D-01 UNITS	2-GW07D-01 UG/L	2-GW08D-01 UG/L	2-GW09D-01 UG/L
ALUMINUM		149 B	43 B	95 B
ANTIMONY		10 U	10 U	3.5 U
ARSENIC		2.9 B	1 U	7.1 B
BARIUM		126 B	49 B	62 B
BERYLLIUM		0.5 U	0.5 U	0.5 U
CADMIUM		2.5 U	2.5 U	2.5 U
CALCIUM		8080	9590	5800
CHROMIUM		5 U	5 U	5 U
COBALT		10 B	8 B	4 U
COPPER		2 B	5 B	4 B
IRON		7070	4660	6180
LEAD		0.5 UJ	0.5 UJ	0.5 UJ
MAGNESIUM		3610 B	3060 B	1730 B
MANGANESE		65	48	40
MERCURY		0.1 U	0.1 U	0.1 U
NICKEL		10 U	10 U	10 U
POTASSIUM		1970 B	1490 B	1150 B
SELENIUM		0.5 U	0.5 U	0.5 U
SILVER		1.5 U	1.5 U	1.5 U
SODIUM		22600	8720	12100
THALLIUM		0.5 U	0.5 U	0.5 U
VANADIUM		2 U	2 U	2 U
ZINC		12 B	13 B	19 B
CYANIDE				35

## **SURFACE WATER**

**BASE BACKGROUND  
SURFACE WATER  
MCB, CAMP LEJEUNE, NORTH CAROLINA  
TAL INORGANICS**

Sample ID:	6-BH01-SW-06B	6-BH01-SW-06M	6-BH02-SW-06M	6-WC01-SW-06B	6-WC01-SW-06M	6-WC02-SW-06B	6-WC03-SW-06B
ALUMINUM	1210	1230	868	1350	1220	633	747
ANTIMONY	17.2 UJ	14 U	14 U	14 U	14 U	16.2 UJ	49 U
ARSENIC	3 U	3 U	3 UJ	3 UJ	3 UJ	2 U	2 U
BARIUM	13.4 JB	14 JB	25.1 JB	16 JB	16.2 JB	19.3 B	21 U
BERYLLIUM	0.3 U	1 U					
CADMIUM	1.9 UJ	2.6 UJ	1.9 U	1.9 U	1.9 U	1.9 U	3 U
CALCIUM	612 B	600 B	16100	3640 B	3670 B	9990	9360
CHROMIUM	3.6 U	3.6 U	7 U	3.6 UJ	3.6 UJ	3.6 U	5 U
COBALT	2 U	2 U	3 UJ	2 U	2 U	2 U	6 U
COPPER	3.2 UJ	3 UJ	7 UJ	1.9 U	1.9 U	1.9 U	4 U
CYANIDE	10 U	10 UJ	10 U				
IRON	958	818	921	1050	941	844	849
LEAD	1 U	1 U	3 U	2.3 JB	1.9 JB	1.2 B	5
MAGNESIUM	588 B	612 B	1010 B	632 B	639 B	1110 B	916 B
MANGANESE	6.5 B	6.2 B	14 JB	9 UJ	8.9 UJ	8.8 B	9.8 JB
MERCURY	0.04 U	0.05 U	0.04 U	0.04 U	0.04 U	0.07 U	0.2 U
NICKEL	7.9 UJ	7.9 UJ	7.9 U	7.9 UJ	7.9 UJ	7.9 U	17 U
POTASSIUM	117 UJ	146 UJ	685 B	376 B	341 B	604 B	610 B
SELENIUM	5 U	5 U	5 U	5 UJ	5 UJ	5 U	5 U
SILVER	2 UJ	2 UJ	4 UJ	2 UJ	2 UJ	3.8 UJ	10 U
SODIUM	4680 B	4850 B	5250	3930 B	3980 B	7790	6240
THALLIUM	2 UJ	2 UJ	2 UJ	2 U	2 UJ	2 UJ	2 UJ
VANADIUM	1.8 UJ	1.8 UJ	2 JB	3.3 JB	1.9 JB	2.1 JB	5 U
ZINC	4.5 U	4.9 U	13.1 U	8.7 U	7.6 U	7.5 U	7.4 U

Concentrations presented in micrograms per liter (UG/L)

**BASE BACKGROUND  
SURFACE WATER  
MCB, CAMP LEJEUNE, NORTH CAROLINA  
TAL INORGANICS**

	Sample ID: 6-WC03-SW-06M	6-WC03-SW-312M	41-TC-SW06	41-UN-SW01	41-NE-SW05	2-OC-SW01	69-UT1-SW-06
ALUMINUM	633	676	390.0	447.0 J	178.0	556	1110
ANTIMONY	49 U	49 U	7.60 U	7.60 U	7.60 U	7 U	49 U
ARSENIC	2 U	2 U	2.90 U	2.20 U	2.90 U	2 U	3 U
BARIUM	21 U	21 U	23.6	23.3	27.2	18 B	23 B
BERYLLIUM	1 U	1 U	0.760 U	0.760 U	0.760 U	1 U	1 U
CADMIUM	3 U	3 U	3.19 U	3.19 U	3.19 U	5 U	3 JB
CALCIUM	8890	9430	18900.0	41600.0	40300.0	22900	1380 B
CHROMIUM	5 U	5 U	8.31 U	8.31 U	8.31 U	10 U	5 U
COBALT	6 U	6 U	16.0 U	16.0 U	16.0 U	8 U	8 JB
COPPER	4 U	129	16.3 U	16.3 U	16.3 U	4 B	7 JB
CYANIDE	10 U	10 U	NZ	NZ	NZ	NZ	10 U
IRON	756	830	1460.0	1300.0 J	469.0	413	1000
LEAD	5	10.4	1.40	1.85	1.17	2 U	2 B
MAGNESIUM	883 B	936 B	1620.0	1770.0	2410.0	1960 B	846 B
MANGANESE	8.2 JB	9.2 JB	25.7	17.5	40.0	24	9 JB
MERCURY	0.2 U	0.52	0.171 U	0.182 UJ	0.160 U	0.2 U	0.2 U
NICKEL	17 U	1380	17.4 U	28.8 U	17.4 U	20 U	17 U
POTASSIUM	603 B	640 B	2210	1860	1620	809 B	385 B
SELENIUM	5 U	5 U	1.60 UJ	1.60 UJ	1.60 UJ	1 U	5 U
SILVER	10 U	10 U	1.60 U	1.60 U	1.60 U	3 UJ	10 U
SODIUM	6100	6500	15000	22100	12300	6190	4790 JB
THALLIUM	2 UJ	2 UJ	3.00 U	3.00 U	3.00 U	1 U	2 UJ
VANADIUM	5 U	5 U	20.4 U	20.4 U	20.4 U	4 U	10 JB
ZINC	10.4 U	111	21.4	24.9	33.2	23 UJ	18 B

Concentrations presented in micrograms per liter (UG/L)

**BASE BACKGROUND**  
**SURFACE WATER**  
**MCB, CAMP LEJEUNE, NORTH CAROLINA**  
**TAL INORGANICS**

	Minumium (ug/L)	Maximum (ug/L)	Average (ug/L)
ALUMINUM	178	1350	803.4
ANTIMONY	ND	ND	NA
ARSENIC	ND	ND	NA
BARIUM	13.4	27.2	17.9
BERYLLIUM	ND	ND	NA
CADMUM	3	3	1.5
CALCIUM	600	41600	13383.7
CHROMIUM	ND	ND	NA
COBALT	8	8	3.7
COPPER	4	129	12.7
CYANIDE	ND	ND	NA
IRON	413	1460	900.6
LEAD	1.17	10.4	2.6
MAGNESIUM	588	2410	1138.0
MANGANESE	6.2	40	13.4
MERCURY	0.52	0.52	0.1
NICKEL	1380	1380	105.1
POTASSIUM	341	2210	776.8
SELENIUM	ND	ND	NA
SILVER	ND	ND	NA
SODIUM	3930	22100	7835.7
THALLIUM	ND	ND	NA
VANADIUM	1.9	10	4.4
ZINC	18	111	18.0

Qualifiers have been removed per Baker's standards.

Qualifiers U and UJ have been given one-half the detection value.

Qualifiers J, NJ, and B have been removed with no detection value change.

**SEDIMENT**

---

**BASE BACKGROUND**  
**SEDIMENT**  
**MCB, CAMP LEJEUNE, NORTH CAROLINA**  
**TAL INORGANICS**

	Sample ID: 2-OCSD01-06	2-OCSD01-612	6-BH01-SD-612B	6-BH01-SD-612M	6-BH01-SD-6B	6-BH01-SD-6M	6-BH02-SD-06M
ALUMINUM	8680	9090	6760	7790	5610	6360	3010
ANTIMONY	R	R	4.7 UJ	5.9 U	4.9 UJ	4.8 U	3.8 U
ARSENIC	0.56 UJ	0.57 UJ	1 U	1.1 U	1.1 U	0.93 U	0.77 U
BARIUM	30.5 B	30 B	9.7 JB	14.4 B	8.5 UJ	9.9 JB	12.5 B
BERYLLIUM	0.85 B	0.86 B	0.13 B	0.17 B	0.14 B	0.1 U	0.08 U
CADMUM	1.4 U	1.4 U	0.51 UJ	0.8 UJ	0.86 UJ	0.65 UJ	0.54 JB
CALCIUM	6320	6180	59.3 U	82.8 U	61.9 U	70.2 U	1410
CHROMIUM	9.9	10	5.1	4.7	4.9	3.6	3.3 U
COBALT	2.3 U	2.3 U	0.53 U	0.84 U	0.55 U	0.69 U	1.1 UJ
COPPER	1.1 B	0.86 B	3.2 JB	10.1 JB	4.2 JB	6.2 JB	2.5 UJ
IRON	842	845	765	1590	638	956	1240
LEAD	8.8	8	8.9	12.3	11.3	10.2	6.9
MAGNESIUM	322 B	307 B	128 B	160 B	103 B	130 B	77.9 B
MANGANESE	4.8	5.7	4.9	6 B	4.7	4.9 B	4.4 J
MERCURY	0.14 U	0.14 U	0.05 U	0.05 U	0.05 U	0.04 UJ	0.03 U
NICKEL	5.6 U	5.7 U	2.1 UJ	3.3 UJ	2.2 UJ	2.7 UJ	2.7 UJ
POTASSIUM	229 B	237 B	125 B	163 B	122 B	140 B	76.8 UJ
SELENIUM	1.7 J	2.1 J	1.7 UJ	1.9 U	1.8 UJ	1.6 UJ	1.3 U
SILVER	0.85 UJ	0.86 UJ	0.53 UJ	0.84 UJ	0.55 UJ	0.69 UJ	0.82 UJ
SODIUM	86.2 B	78.9 B	35.5 UJ	42.8 UJ	41.5 UJ	39.4 UJ	25.4 UJ
THALLIUM	0.31 J	0.29 J	0.69 U	0.76 U	0.73 U	0.62 U	0.51 U
VANADIUM	6.8 B	6.6 B	5.7 B	6.5 B	4.8 B	4.9 B	3.3 JB
ZINC	18.9	18.9	2.1 U	1.4 U	1.6 U	1.8 U	12

Concentrations presented in milligrams per kilogram (mg/kg).

**BASE BACKGROUND**  
**SEDIMENT**  
**MCB, CAMP LEJEUNE, NORTH CAROLINA**  
**TAL INORGANICS**

	Sample ID:	6-BH02-SD-612M	6-WC01-SD-06B	6-WC01-SD-612B	6-WC02-SD-06B	6-WC02-SD-612B	6-WC03-SD-06B	6-WC03-SD-06M
ALUMINUM		7780	2090 J	2510	6540 J	5390 J	6480 J	4780 J
ANTIMONY		4.6 U	3.3 U	3.1 U	3.1 U	4.1 U	6.8 UJ	3.4 U
ARSENIC		1.6 JB	1.2 JB	0.73 UJ	0.81 U	0.64 U	1.4 UJ	0.82 UJ
BARIUM		30 B	5.2 JB	15.3 B	19.6 JB	23.7 JB	15.8 JB	37.1 JB
BERYLLIUM		0.33 B	0.07 U	0.07 U	0.26 U	0.33 U	0.27 U	0.32 U
CADMIUM		1.3 JB	0.45 U	0.42 U	0.42 U	0.74 UJ	1.2 UJ	0.46 U
CALCIUM		3890	329 B	1060 B	1090 JB	1790 J	2850 J	22200 J
CHROMIUM		9.9	3 UJ	2.5 UJ	4.2	3.4	6.2	6.4
COBALT		2.6 UJ	0.48 U	0.44 U	0.6 JB	0.87 JB	0.94 U	1.3 JB
COPPER		2.3 UJ	0.86 UJ	0.64 UJ	0.43 JB	0.62 JB	5.8 JB	53200
IRON		3150	724 J	1430 J	1200 J	1570 J	6870 J	6940 J
LEAD		8.9	9.7 J	2.3 J	4.8 J	4.8 J	9 J	314 J
MAGNESIUM		187 B	50.5 B	57 B	372 JB	356 JB	440 JB	852 JB
MANGANESE		8.6 J	2.4 UJ	4.7 J	8.8	6.5	9.7	23
MERCURY		0.07 U	0.03 U	0.04 U	0.08 U	0.06 U	0.11 U	0.06 U
NICKEL		7.2 UJ	1.9 UJ	1.8 UJ	1.7 UJ	2.8 B	3.7 UJ	1.9 UJ
POTASSIUM		151 U	92.1 B	98.1 B	145 B	97 U	220 B	360 B
SELENIUM		2.9	1.4 UJ	1.2 UJ	1 U	1.3 U	2.7 U	1 UJ
SILVER		1.3 UJ	0.48 UJ	0.44 UJ	0.52 UJ	1.2 UJ	1.5 UJ	7.3
SODIUM		39.9 UJ	38.3 UJ	27 UJ	491 JB	469 JB	277 UJ	489 JB
THALLIUM		0.65 UJ	0.55 U	0.49 U	0.4 UJ	0.5 UJ	1.1 UJ	0.4 UJ
VANADIUM		14.1 B	5.7 B	4.4 B	5.8 B	7 B	11.6 B	9.1 B
ZINC		12.6	3.1 U	3.1 U	1.6 U	2.4 U	16.3 U	926

Concentrations presented in milligrams per kilogram (mg/kg).

**BASE BACKGROUND**  
**SEDIMENT**  
**MCB, CAMP LEJEUNE, NORTH CAROLINA**  
**TAL INORGANICS**

	Sample ID:	6-WC03-SD-612B	41-UN-SD01-06	41-UN-SD01-612	41-NE-SD05-06	41-NE-SD05-612	41-TC-SD06-06
ALUMINUM		7040 J	1720.0	2780.0	437 J	351 J	2580.0 J
ANTIMONY		6.8 U	2.15 U	2.09 U	1.91 U	1.88 U	2.28 U
ARSENIC		1.3 JB	0.789 U	0.768 U	0.542 U	0.532 U	0.702
BARIUM		25.2 JB	5.24	7.66	3.2 U	3.14 U	13.5
BERYLLIUM		0.26 U	0.351 U	0.342 U	0.196 U	0.193 U	0.234 U
CADMUM		0.92 U	0.639 U	0.622 U	0.823 U	0.809 U	0.982 U
CALCIUM		4500 J	1250.0	1660.0	314 J	216 J	1090.0 J
CHROMIUM		8.3	4.81 U	3.18 U	2.42 J	2.11 UJ	3.42 J
COBALT		0.97 U	2.65 U	2.58 U	4.13 U	4.06 U	4.92 U
COPPER		79.6	4.41 U	4.29 U	4.21 U	4.13 U	5.02 U
IRON		6050 J	924.0 J	1160.0 J	354 J	262 J	2840.0 J
LEAD		10.3 J	13.8 J	12.6 J	1.94	2.19	18.7
MAGNESIUM		333 JB	62.5	59.4	21.5	18.2 U	99.8
MANGANESE		8.3	2.94	2.67	1.96 J	1.79 UJ	8.72 J
MERCURY		0.11 U	0.068 U	0.066 U	0.064 U	0.063 U	0.077 U
NICKEL		3.8 UJ	5.97	3.79	7.4 U	7.3 U	8.90 U
POTASSIUM		457 B	136.0 U	132.0 U	197 U	193 U	235.0 U
SELENIUM		2.3 U	0.688 U	0.670 U	0.387 UJ	0.38 UJ	0.462 UJ
SILVER		1.3 UJ	0.435 U	0.424 U	0.413 UJ	0.406 UJ	0.492 UJ
SODIUM		382 UJ	73.6 J	49.3 UJ	95 U	117	347.0
THALLIUM		0.93 UJ	1.25 U	1.22 U	0.748 UJ	0.735 UJ	0.892 UJ
VANADIUM		15.7 B	4.52 U	4.40 U	5.26 U	5.17 U	6.28 U
ZINC		12.3 U	10.5 U	15.2 U	7.41 U	13.6	18.0

Concentrations presented in milligrams per kilogram (mg/kg).

BASE BACKGROUND  
SEDIMENT  
MCB, CAMP LEJEUNE, NORTH CAROLINA  
TAL INORGANICS

Sample ID: 410TC-SD06-612      69-UT1-SD-06

ALUMINUM	6600.0 J	1240
ANTIMONY	2.11 U	9.4 U
ARSENIC	0.864	0.62 U
BARIUM	25.3	4 U
BERYLLIUM	0.377	0.19 U
CADMIUM	0.909 U	0.58 U
CALCIUM	1230.0 J	264 B
CHROMIUM	8.72 J	3.3
COBALT	4.56 U	1.2 UJ
COPPER	4.64 U	1.5 UJ
IRON	6030.0 J	3530
LEAD	13.6	1
MAGNESIUM	235.0	48.9 B
MANGANESE	13.7 J	2.9 J
MERCURY	0.071 U	0.11 U
NICKEL	8.20 U	3.3 U
POTASSIUM	381.0	81.1 B
SELENIUM	0.862 J	1 U
SILVER	0.456 UJ	1.9 U
SODIUM	105.0 U	122 JB
THALLIUM	0.826 UJ	0.42 UJ
VANADIUM	12.7	4 UJ
ZINC	19.9	4.4 U

Concentrations presented in milligrams per kilogram (mg/kg).

**APPENDIX N**  
**WHITE OAK RIVER BASIN REFERENCE STATION**

---

## WHITE OAK RIVER BASIN REFERENCE STATIONS

### *Water Body Description*

Hadnot Creek, Holland Mill Creek (including Cartwheel Branch) and the section of the White Oak River that encompasses Hadnot Creek, Holland Mill Creek, and Webb Creek are classified as SA from their source to the White Oak River. The SA classifies the water body as a tidal saltwater with shellfishing for market purposes and the following uses: primary recreation, aquatic life propagation and survival, fishing, wildlife, and secondary recreation. Webb Creek is classified as C from its source to the White Oak River. The C classifies the water body as a fresh water with the following uses: aquatic life propagation and survival, fishing, wildlife, and secondary recreation. The section of the White Oak River that encompasses these three creeks is designated by the North Carolina Fisheries Rule as Class C - coastal fishing waters (NCMFC, 1993).

### *Biological Sampling*

Biological samples collected at the background stations consisted of fish and benthic macroinvertebrate. The biological samples were collected to obtain population statistics for fish and benthic macroinvertebrates and to obtain fish tissue samples for chemical analysis (Hadnot Creek only). Prior to initiating the sampling event at each station, the following information describing the site was recorded in the field log book:

Average width, depth and velocity of the water body

Description of substrate

Description of "abiotic" characteristics of the reach such as pools, riffles, runs, channel shape, degree of bank erosion, and shade/sun exposure

Description of "biotic" characteristics of the reach including aquatic and riparian vegetation and wetlands

Water quality measurements were collected during the benthic macroinvertebrate sampling, at a minimum, and during collection of some of the fish samples. On-site water quality measurements at these stations consisted of temperature, pH, specific conductance, salinity and dissolved oxygen. These measurements were conducted prior to sample collection. The station locations and sampling procedures for the collection of the fish and benthic macroinvertebrates is discussed later in this appendix.

### Fish and Shellfish

This section discusses collection of the fish and shellfish samples in the reference stations at Webb Creek, Hadnot Creek, and Holland Mill Creek.

A literature review was conducted to determine the fish species that may potentially be exposed to contaminants in the surface water/sediment exposure pathway. This review included compiling information from State and Federal natural resources agencies. In addition, Bakers experience in sampling similar areas formed a basis for a database of expected species for the area.

Sampling variability can prevent the same species of fish from being sampled at each station because either the preferred species was not captured, or adequate numbers of uniform-size individuals were not captured. Therefore, if the preferred species was not successfully collected to satisfy the above requirements, a substitute species was collected that, if possible, exhibiting a similar trophic position in the estuarine ecosystem.

The collected fish species were identified, measured, and counted. The small fish (less than 20 mm) were weighed in groups of 10 or 20 because of their low individual weight; the larger fish were weighed individually. The

proportion of individuals as hybrids and the proportion of individuals with disease, tumors, fin damage, and skeletal anomalies was recorded at each station.

Fish that exhibited signs of being dead for an extended period of time (i.e., brown gills, bloating) were not retained for tissue analysis because of the potential for decomposition and leaching of contaminants from the organs into the edible portions of the fish.

#### Webb Creek

This section discusses collection of the fish samples in Webb Creek including the station locations and sampling procedures.

##### *Station Location*

The fish station WC02 was located on Webb Creek approximately 300 feet upstream from the Camp Lejeune railroad crossing. Station WC03 was located in the White Oak River approximately 25 feet downstream from its confluence with Webb Creek. See fish and benthic macroinvertebrate sampling station figure found later in this appendix for approximate sample locations.

##### *Sampling Procedures*

Fish were collected in Webb Creek using gill nets and hoop nets. All fish that were collected were processed for population statistics; no fish at these stations were collected for tissue analysis.

The gill nets were six feet deep by 50 to 100 feet long with a stretch mesh size ranging from two to four inches, and an approximate twine break strength of 29 pounds. The nets were deployed approximately at the locations shown on the figure found later in this appendix. Weights were attached to the nets to secure them on the bottom of the stream and yellow buoys marked with "Baker Environmental" were attached to the tops of the nets. The nets were deployed in the morning or evening, and they were checked for fish within twelve hours after deployment.

The hoop nets were three to four feet in diameter and fourteen to sixteen feet in length. Twenty-five foot wings were attached to the nets to help direct fish into the net. The nets were deployed in the middle of the channel with the wings stretched across the creek in a forty-five degree angle. The end of the net and the wings were secured using 6.5 foot wooden posts. The nets were checked at least once daily, as the fish usually survive when captured in these nets.

#### Hadnot Creek

This section discusses collection of the fish samples in Hadnot Creek including the station locations and sampling procedures.

##### *Station Location*

Fish were collected from four stations in Hadnot Creek (HC01, HC02, HC03 and HC04). HC01 was located approximately 100 feet upstream of Rt. 1104. Station HC02 was located approximately 2,500 feet upstream of Rt. 58. Station HC03 was located in the White Oak River approximately 100 feet upstream from its confluence with Hadnot Creek. Finally, station HC04 was located in Hadnot Creek by the road off of the Rt. 1105 crossing. In October, 1993, fish were collected by Baker in Hadnot Creek as part of another investigation (Baker, 1993). Fillet samples of these fish were chemically analyzed and the results are included in this ERA.

##### *Sampling Procedures*

Fish were collected at these stations for population statistics; fish were not collected at these stations for tissue analysis. Fish were collected in Hadnot Creek using hoop nets, gill nets, a haul seine, pole fishing, and the

backpack electroshocker. The same sample collection and sample processing procedures used in Webb Creek were conducted at the Hadnot Creek stations for the gill nets and hoop nets. Pole fishing only was conducted during the October 1993 sampling.

Fish were collected in the furthest upstream stations using electrofishing, conducted with a Smith-Root, Inc., backpack electrofisher powered by a 300-watt portable generator. A DC current was applied utilizing a "rattail" as the cathode and a hand-held electrode as the anode. Blocking seines were placed downstream and upstream of the shocking areas to aid in the collection of the fish. The length of the shocking time per subsection was recorded as seconds of applied current. Stunned fish were collected with one-inch mesh or smaller dip nets handled by members of the field sampling team.

#### Holland Mill Creek

This section discusses collection of the fish samples in Holland Mill Creek including the station locations and sampling procedures.

##### *Station Location*

Fish were collected from three stations in Holland Mill Creek (HM01, HM02, and HM03). HM01 was located on Cartwheel Branch just upstream of Rt. 1444. Station HM02 was located at the confluence of Holland Mill Creek and Cartwheel Branch. Station HM03 was located in the White Oak River approximately 50 feet downstream from Holland Mill Creek.

##### *Sampling Procedures*

Fish were collected at these stations for population statistics. Fish were not collected at these stations for tissue analysis. Fish were collected in Holland Mill Creek using hoop nets, gill nets, a haul seine, and the backpack electroshocker. The same sample collection and sample processing procedures used in the Webb Creek and Hadnot Creek stations were conducted at the Holland Mill Creek stations.

#### Benthic Macroinvertebrates

This section discusses collection of benthic macroinvertebrate samples in the reference stations at Webb Creek, Hadnot Creek, and Holland Mill Creek.

#### Webb Creek

Benthic macroinvertebrates were collected in Webb Creek using the ponar grab deployed from the boat.

Benthic macroinvertebrates were collected from a boat using a standard ponar grab. The dimensions of the ponar are 23 x 23 cm (9 x 9 in.) for a sampling area of 529 cm<sup>2</sup> or 0.0523 m<sup>2</sup> (81 in<sup>2</sup>).

The ponar was deployed from the boat, which was positioned in slightly different locations for each replicate to prevent re-sampling the same area. After retrieving the ponar with a sediment sample, it was opened into a clean tub and the sediments were removed with a teflon spatula. The sediments were transferred to a 0.5 mm sieve that was agitated (by hand) in water to remove the small particles. The remaining contents in the sieve were transferred into 16-ounce plastic sample jars. The jars were filled up to one-half full with sediments, and buffered formalin solution (10 percent by weight) was added to the remainder of the jar to preserve the benthic macroinvertebrates contained in the sediments. A 100 percent cotton paper label, marked in pencil with the sample number, was placed inside the jar. The outside of the jar was labeled with the sample number using a black permanent marker to identify the sample containers.

After all the benthic macroinvertebrate sampling at the New River was completed, the sample jars were transported to RMC Environmental Services, Inc. for sample sorting and taxonomic identification of the benthic

macroinvertebrates.

#### Hadnot Creek

Benthic macroinvertebrates were collected in Hadnot Creek using the ponar grab deployed from the boat. The boat was not used at HC01 or HC04 because the water was too shallow. Benthic macroinvertebrates were collected using the same procedures used for collecting benthic macroinvertebrates in Webb Creek.

#### Holland Mill Creek

Benthic macroinvertebrates were collected in Holland Mill Creek using the ponar grab deployed from the boat. The boat was not used at HM01 because the water was too shallow. The same sample collection and sample processing procedures used in Webb Creek were conducted at the Holland Mill Creek stations.

#### Biological Tissue Sample Results

The analytical parameters included TCL VOCs, TCL SVOCs, TAL metals, and TCL pesticides/PCBs. Background fish fillet tissue were collected from Hadnot Creek and analyzed these results are discussed below.

##### *Hadnot Creek*

Several metals were detected in the Hadnot Creek fillet tissue samples. These metals included aluminum, arsenic, calcium, chromium, copper, magnesium, manganese, mercury, nickel, potassium, sodium and zinc in the fillet samples. The range of detected levels for these chemicals in the fish fillet tissue samples from Hadnot Creek are as follows:

	<u>Minimum (mg/kg)</u>	<u>Maximum (mg/kg)</u>
Aluminum	36.5	36.5
Arsenic	0.34	3.9
Calcium	154	1,170
Chromium	0.21	0.68
Copper	0.18	0.46
Magnesium	254	319
Manganese	0.008	0.38
Mercury	0.05	0.24
Nickel	0.45	0.45
Potassium	3,270	4,040
Sodium	505	1,060
Zinc	3.9	6.5

The maximum detect of manganese was in the southern flounder. The maximum detect of sodium was found in the red drum. Aluminum, calcium, chromium, magnesium, mercury, and potassium were detected at their highest concentrations in the largemouth bass. The maximum detects of arsenic, copper, nickel, and zinc were found in the longnose gar.

Two pesticides were detected in the fillet tissue samples, 4-4'-DDE and alpha-chlordane. 4,4'-DDE was detected twice, both in the longnose gar. Alpha-chlordane was detected once in the largemouth bass. The range of detected concentrations for these constituents were as follows:

	<u>Minimum (ug/kg)</u>	<u>Maximum (ug/kg)</u>
4-4'-DDE	9.7	12.0
alpha-Chlordane	0.17	0.17

Two VOCs and three SVOCs were detected in the fillet tissue samples. Common laboratory contaminants were the primary detections, which included methylene chloride, acetone, di-n-octyl phthalate and bis(2-ethylhexyl)phthalate. Phenol was also detected in the fillet tissue samples. The concentration ranges for these chemicals were the following:

	<u>Minimum (ug/kg)</u>	<u>Maximum (ug/kg)</u>
Methylene chloride	3.0	41.0
Acetone	16	130
di-n-octyl phthalate	61	500
bis(2-ethylhexyl) phthalate	820	17,000
Phenol	460	2,100

#### Field Chemistry Results

Samples from these surface water bodies were collected from the water surface and bottom.

##### *Webb Creek*

At Webb Creek, the salinity at station WC02 ranged from 0 to 7 ppt. Conductivity ranged from 850 to 10,500 micromhos/cm. Dissolved oxygen levels ranged from 4.4 to 9 mg/L. The pH at station WC02 in Webb Creek ranged from 6.85 to 7.48 S.U. in the surface water. The temperature of the water at WC02 ranged from 17.5 to 21 °C.

At WC03, the salinity ranged from 10 to 12.8 ppt. The conductivity ranged from 16,500 to 18,000 micromhos/cm. Dissolved oxygen levels ranged from 8.5 to 10 mg/L. The pH at WC03 in Webb Creek ranged from 7.33 to 7.56 S.U. in the surface water. The temperature of the water at WC03 ranged from 19 to 23 °C.

##### *Hadnot Creek*

In Hadnot Creek, the salinity at station HC01 was 0 ppt. The conductivity was 13.5 micromhos/cm. The dissolved oxygen level was 7.7 mg/L. The pH at HC01 was 6.89 S.U. in the surface water, and the temperature of the Hadnot Creek water was 17 °C.

At station HC02, the salinity ranged from 0 to 16.5 ppt. The conductivity ranged from 720 to 22,800 micromhos/cm. The dissolved oxygen levels ranged from 1 to 7.3 mg/L. The pH at HC02 ranged from 6.7 to 7.2 S.U. in the surface water. The temperature of the water at HC02 ranged from 15.5 to 22 °C.

At station HC03, the salinity ranged from 17 to 17.9 ppt. The conductivity ranged from 25,500 to 26,500 micromhos/cm. The dissolved oxygen level was 12 mg/L. The pH at HC03 ranged from 7.69 to 7.79 S.U. in the surface water. The temperature of the water at HC03 ranged from 17.5 to 17.8 °C.

At station HC04, the salinity was 0 ppt. The conductivity was 65 micromhos/cm, and the dissolved oxygen level was 5.3 mg/L. The pH at HC04 was 6.16 S.U. in the surface water, and the temperature of the water was 17.3 °C.

##### *Holland Mill Creek*

In Holland Mill Creek, the salinity was 0 ppt at station HM01. The conductivity was 140 micromhos/cm, and the dissolved oxygen level was 8.0 mg/L. The pH at station HM01 was 6.9 S.U. in the surface water, and the temperature of the water was 17.5 °C.

At station HM02, the salinity ranged from 1 to 25 ppt. The conductivity ranged from 2,490 to 38,000 micromhos/cm. The dissolved oxygen levels ranged from 5.0 to 11.8 mg/L. The pH at station HM02 ranged from 6.72 to 7.9 S.U. in the surface water. The temperature of the water at HM02 ranged from 15.2 to 20 °C.

At station HM03, the salinity ranged from 13.5 to 22 ppt. The conductivity ranged from 19,000 to 32,000 micromhos. The dissolved oxygen levels ranged from 3.4 to 10.8 mg/L. The pH at station HM03 ranged from 6.81 to 7.90 S.U. in the surface water. The temperature of the water at HM03 ranged from 17.5 to 17.8 °C.

**Statistical Summary of  
Analytical Results  
(Surface Water)**

## KEY TO STATISTICAL AND ANALYTICAL SUMMARY TABLES

U - Indicated analyte was analyzed for but not detected

J - Indicates an estimated value

UI - Not detected, quantitation limit may be inaccurate or imprecise

R - Result is rejected and unusable

B - Not detected substantially above the level reported in laboratory or field blanks (organics)

P - There is greater than 25 % difference for detected pesticide/PCB concentrations between the two GC columns, the lower of the two values is reported

L - Result is biased low

K - Result is biased high

ND - Analyte not detected

NZ - Analyte not analyzed

mg/L - Milligrams per liter

ug/L - Micrograms per liter

mg/kg - Milligrams per kilogram

ug/kg - Micrograms per kilogram

APPENDIX N (CONT.)  
 BACKGROUND - UPSTREAM  
 SURFACE WATER - METALS  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	LOG NORMAL UPPER 95% CONFIDENCE LEVE (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
Aluminum	692.00	692.00	*+ HC-SW04	333.17	858.66	265385.03	1	3	33%
Barium	9.00	49.00	*+ HM-SW01	25.67	60.76	15294.20	3	3	100%
Calcium	11600.00	27000.00	*+ HC-SW01	17566.67	31499.62	92587.15	3	3	100%
Iron	425.00	746.00	*+ HC-SW01	575.67	847.77	1123.79	3	3	100%
Magnesium	954.00	2830.00	*+ HM-SW01	1744.67	3383.48	22741.37	3	3	100%
Selenium	1.50	1.50	*+ HM-SW01	0.83	1.81	29.24	1	3	33%
Sodium	6090.00	16500.00	*+ HM-SW01	9830.00	19592.11	124185.69	3	3	100%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

## APPENDIX N (CONT.)

## BACKGROUND - UPSTREAM

## SURFACE WATER - PESTICIDES/PCBs

## STATISTICAL SUMMARY OF ANALYTICAL RESULTS

PARAMETER	MINIMUM DETECTED VALUE	MAXIMUM DETECTED VALUE	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE VALUE	RME	LOG NORMAL UPPER 95% CONFIDENCE LEVE	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
	(ug/L)	(ug/L)		(ug/L)	(ug/L)	(ug/L)			
No Pesticides/PCBs Were Detected									

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

## APPENDIX N (CONT.)

BACKGROUND - MIDSTREAM

SURFACE WATER - METALS

STATISTICAL SUMMARY OF ANALYTICAL RESULTS

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	LOG NORMAL UPPER 95% CONFIDENCE LEVE (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
Aluminum	535.00	535.00	*+ HM-SW02	270.67	657.22	47743.23	1	3	33%
Barium	20.00	38.00	*+ HM-SW02	29.00	44.17	75.32	3	3	100%
Calcium	36600.00	46900.00	*+ WC-SW02	41233.33	50045.63	52734.83	3	3	100%
Chromium	36.00	36.00	*+ HM-SW02	17.67	44.49	10009.89	1	3	33%
Iron	528.00	660.00	*+ WC-SW02	582.33	698.70	731.17	3	3	100%
Magnesium	29000.00	109000.00	*+ HM-SW02	60933.33	132363.27	2716191.24	3	3	100%
Potassium	10900.00	41100.00	*+ HM-SW02	22166.67	49975.40	1126700.03	3	3	100%
Sodium	202000.00	739000.00	*+ HM-SW02	441333.33	901929.85	17057165.85	3	3	100%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

APPENDIX N (CONT.)  
 BACKGROUND - MIDSTREAM  
 SURFACE WATER - PESTICIDE/PCBs  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS

PARAMETER	MINIMUM	MAXIMUM	SAMPLE No.	LOG NORMAL				No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
	DETECTED VALUE (ug/L)	DETECTED VALUE (ug/L)	OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	CONFIDENCE LEVE (ug/L)	No. OF TIMES DETECTED		
Aldrin	0.04	0.04	*+ WC-SW02	0.03	0.04	0.04	1	3	33%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

APPENDIX N (CONT.)  
 BACKGROUND - DOWNSTREAM  
 SURFACE WATER - METALS  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	LOG NORMAL UPPER 95% CONFIDENCE LEVE (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
Arsenic	20.00	20.00	+ HC-SW03	8.13	17.54	334.25	1	4	25%
Barium	20.00	27.00	*+ WC-SW03	24.25	27.89	29.23	4	4	100%
Calcium	405000.00	302000.00	+ HM-SW03	134025.00	269777.00	4658189.53	4	4	100%
Chromium	97.00	158.00	+ HM-SW03	127.50	156.90	176.16	4	4	100%
Iron	291.00	339.00	*+ HC-SW03	317.75	341.11	347.02	4	4	100%
Lead	58.10	58.10	+ HM-SW03	16.41	49.11	3685198.93	1	4	25%
Magnesium	44800.00	754000.00	*+ HM-SW03	511200.00	884281.78	5105380691.44	4	4	100%
Potassium	136000.00	288000.00	+ HM-SW03	207250.00	280542.42	373364.83	4	4	100%
Selenium	6.00	41.00	+ HM-SW03	12.50	35.01	330407465.54	2	4	50%
Silver	37.00	37.00	+ HM-SW03	19.13	33.16	80.27	1	4	25%
Sodium	895000.00	6750000.00	+ HM-SW03	3073750.00	6072745.87	108244495.85	4	4	100%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

## APPENDIX N (CONT.)

BACKGROUND - DOWNSTREAM

SURFACE WATER - PESTICIDES/PCBs

STATISTICAL SUMMARY OF ANALYTICAL RESULTS

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	LOG NORMAL UPPER 95% CONFIDENCE LEVE (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
No Pesticides/PCBs Were Detected									

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HADNOT CREEK  
 SURFACE WATER - METALS

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION	
Aluminum	692.00	692.00	+	HC-SW04	253.10	488.87	1019.72	1	5	20%
Arsenic	20.00	20.00	+	HC-SW03	5.30	13.35	3190.11	1	5	20%
Barium	9.00	26.00	+	HC-SW03	19.60	25.87	35.22	5	5	100%
Calcium	11600.00	107000.00	+	HC-SW03D	53760.00	92784.90	456379.04	5	5	100%
Chromium	125.00	130.00	+	HC-SW03	54.70	118.12	40374.07	2	5	40%
Iron	291.00	746.00	+	HC-SW01	492.00	666.33	793.41	5	5	100%
Magnesium	954.00	633000.00	+	HC-SW03	258640.80	576299.05	1.50E+16	5	5	100%
Potassium	14500.00	203000.00	+	HC-SW03	84234.00	187308.88	5.24E+12	3	5	60%
Selenium	6.00	6.00	+	HC-SW03	2.00	4.29	38.67	1	5	20%
Sodium	6090.00	2560000.00	+	HC-SW03D	1.01E+06	2.17E+06	4.80E+14	5	5	100%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HADNOT CREEK  
 SURFACE WATER - PESTICIDES/PCBs

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
NO PESTICIDES/PCBs WERE DETECTED									

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HADNOT CREEK  
 SURFACE WATER - SEMIVOLATILE ORGANIC COMPOUNDS

PARAMETER	MINIMUM DETECTED VALUE	MAXIMUM DETECTED VALUE	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE VALUE	RME	LOG NORMAL UPPER 95% CONFIDENCE LEVEL	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
	(ug/L)	(ug/L)		(ug/L)	(ug/L)	(ug/L)			
NO SEMIVOLATILE ORGANIC COMPOUNDS WERE DETECTED									

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HADNOT CREEK  
 SURFACE WATER - VOLATILE ORGANIC COMPOUNDS

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
NO VOLATILE ORGANIC COMPOUNDS WERE DETECTED									

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HOLLAND MILL CREEK  
 SURFACE WATER - METALS

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
Aluminum	535.00	535.00	*+ HM-SW02	269.50	657.32	48037.76	1	3	33%
Barium	20.00	49.00	*+ HM-SW01	35.67	60.35	204.30	3	3	100%
Calcium	14100.00	302000.00	*+ HM-SW03	118766.67	387190.45	4.42E+14	3	3	100%
Chromium	36.00	158.00	*+ HM-SW03	66.33	202.69	3.67E+12	2	3	67%
Iron	320.00	559.00	*+ HM-SW02	434.67	636.62	843.56	3	3	100%
Lead	58.10	58.10	*+ HM-SW03	19.95	75.65	1.70E+27	1	3	33%
Magnesium	2830.00	754000.00	*+ HM-SW03	288610.00	973947.76	1.02E+35	3	3	100%
Potassium	41100.00	288000.00	*+ HM-SW03	109978.33	372096.67	1.33E+36	2	3	67%
Selenium	1.50	41.00	*+ HM-SW03	15.00	52.97	8.42E+13	2	3	67%
Silver	37.00	37.00	*+ HM-SW03	16.83	46.42	284713.62	1	3	33%
Sodium	16500.00	6750000.00	*+ HM-SW03	2501833.33	8733985.25	1.96E+44	3	3	100%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HOLLAND MILL CREEK  
 SURFACE WATER - PESTICIDES/PCBs

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
NO PESTICIDES/PCBs WERE DETECTED									

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HOLLAND MILL CREEK  
 SURFACE WATER - SEMIVOLATILE ORGANIC COMPOUNDS

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
NO SEMIVOLATILE ORGANIC COMPOUNDS WERE DETECTED									

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HOLLAND MILL CREEK  
 SURFACE WATER - VOLATILE ORGANIC COMPOUNDS

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
NO VOLATILE ORGANIC COMPOUNDS WERE DETECTED									

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - WEBB CREEK  
 SURFACE WATER - METALS

PARAMETER	MINIMUM DETECTED VALUE	MAXIMUM DETECTED VALUE	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE	RME	LOG NORMAL UPPER 95% CONFIDENCE LEVEL	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
	(ug/L)	(ug/L)		(ug/L)	(ug/L)	(ug/L)			
Barium	27.00	29.00	*+ WC-SW02	28.00	34.31	32.19	2	2	100%
Calcium	40500.00	46900.00	*+ WC-SW02	43700.00	63904.80	58284.51	2	2	100%
Chromium	97.00	97.00	*+ WC-SW03	52.25	334.80	1.32E+20	1	2	50%
Iron	321.00	660.00	*+ WC-SW02	490.50	1560.72	14358.69	2	2	100%
Magnesium	29000.00	44800.00	*+ WC-SW03	36900.00	86780.60	133710.58	2	2	100%
Potassium	10900.00	136000.00	*+ WC-SW03	73450.00	468390.70	1.01E+23	2	2	100%
Sodium	202000.00	895000.00	*+ WC-SW03	548500.00	2736301.00	6.83E+11	2	2	100%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - WEBB CREEK  
 SURFACE WATER - PESTICIDES/PCBs

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
Aldrin	0.04	0.04 *+	WC-SW02	0.03	0.06	0.07	1	2	50%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - WEBB CREEK  
 SURFACE WATER - SEMIVOLATILE ORGANIC COMPOUNDS

PARAMETER	MINIMUM DETECTED	MAXIMUM DETECTED	SAMPLE No. OF MAXIMUM	ARITHMETIC DETECTED	LOG NORMAL UPPER 95%	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
	VALUE (ug/L)	VALUE (ug/L)	VALUE	AVERAGE (ug/L)	RME (ug/L)			
NO SEMIVOLATILE ORGANIC COMPOUNDS WERE DETECTED								

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*\* = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - WEBB CREEK  
 SURFACE WATER - VOLATILE ORGANIC COMPOUNDS

PARAMETER	MINIMUM DETECTED VALUE (ug/L)	MAXIMUM DETECTED VALUE (ug/L)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/L)	RME (ug/L)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/L)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
NO VOLATILE ORGANIC COMPOUNDS WERE DETECTED									

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

Statistical Summary of  
Analytical Results  
(Sediment)

## APPENDIX N (CONT.)

## BACKGROUND - UPSTREAM

## SEDIMENT - METALS

## STATISTICAL SUMMARY OF ANALYTICAL RESULTS

PARAMETER	MINIMUM DETECTED VALUE (mg/kg)	MAXIMUM DETECTED VALUE (mg/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (mg/kg)	RME (mg/kg)	LOG NORMAL UPPER 95% CONFIDENCE LEVE (mg/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION	
Aluminum	337.00	2940.00	+	HC-SD01-06	1165.57	1864.95	3139.37	7	7	100%
Arsenic	0.26	0.46	*+	HC-SD01-06	0.37	0.50	0.91	4	7	57%
Barium	4.10	16.30	+	HC-SD01-06	6.46	11.12	35.74	4	7	57%
Beryllium	0.14	0.14	+	HC-SD01-06	0.09	0.13	3228.49	1	4	25%
Cadmium	0.03	0.11		HM-SD01-06D	0.04	0.06	0.07	6	7	86%
Calcium	282.00	3620.00	+	HC-SD01-06	1967.14	2972.71	10565.45	7	7	100%
Chromium	1.10	3.20		HC-SD01-612	1.86	2.38	2.56	7	7	100%
Copper	0.66	1.10		HC-SD01-612	0.75	0.92	0.98	4	7	57%
Iron	225.00	648.00	+	HC-SD01-06	433.71	558.82	650.45	7	7	100%
Lead	0.62	1.00	*+	HM-SD01-612	0.79	1.11	1.34	3	3	100%
Magnesium	26.70	87.70		HC-SD01-06	45.25	64.25	76.34	5	7	71%
Manganese	1.30	6.90	+	HC-SD01-06	3.63	5.31	7.80	7	7	100%
Mercury	0.09	0.18	*+	HM-SD01-612	0.14	0.22	16712.85	3	3	100%
Selenium	0.21	0.34	+	HC-SD01-612	0.19	0.27	0.36	4	7	57%
Silver	0.49	0.49		HM-SD01-06	0.25	0.33	0.33	1	7	14%
Thallium	0.13	0.16		HC-SD01-612	0.10	0.13	0.15	3	7	43%
Vanadium	0.66	2.80	+	HC-SD01-612	1.52	2.23	4.68	6	7	86%
Zinc	6.70	9.70	+	HM-SD01-06	5.11	7.40	10.62	3	7	43%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

## APPENDIX N (CONT.)

BACKGROUND - UPSTREAM

SEDIMENT - PESTICIDES/PCBs

STATISTICAL SUMMARY OF ANALYTICAL RESULTS

PARAMETER	MINIMUM DETECTED VALUE (ug/kg)	MAXIMUM DETECTED VALUE (ug/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/kg)	RME (ug/kg)	LOG NORMAL UPPER 95%		No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
						CONFIDENCE LEVE (ug/kg)	No. OF TIMES			
beta-BHC	1.70	7.30	HM-SD01-06D	2.51	4.22	5.97	3	7	43%	
delta-BHC	0.64	0.64	*+ HC-SD01-06	1.09	1.26	1.34	1	7	14%	
Heptachlor	0.48	2.00	HC-SD04-612	1.18	1.51	1.79	2	7	29%	
Aldrin	0.56	0.72	*+ HM-SD01-612	1.05	1.27	1.41	2	7	29%	
Dieldrin	0.58	1.50	*+ HM-SD01-612	1.96	2.49	3.51	2	7	29%	
4,4'-DDE	1.00	4.30	HM-SD01-612	2.42	3.15	3.70	2	7	29%	
4,4'-DDD	0.87	3.10	HM-SD01-612	1.57	2.12	2.33	2	7	29%	
4,4'-DDT	1.70	1.70	*+ HM-SD01-612	2.20	2.44	2.47	1	7	14%	
Methoxychlor	0.94	0.94	*+ HC-SD04-06	10.21	13.34	49.17	1	7	14%	
Endrin aldehyde	0.59	0.59	*+ HC-SD01-06	2.01	2.51	3.54	1	7	14%	
alpha-Chlordane	1.30	1.30	+ HM-SD01-612	1.20	1.29	1.30	1	7	14%	
gamma-Chlordane	3.00	3.00	HM-SD01-612	1.44	1.95	1.98	1	7	14%	

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

## APPENDIX N (CONT.)

## BACKGROUND - MIDSTREAM

## SEDIMENT - METALS

## STATISTICAL SUMMARY OF ANALYTICAL RESULTS

PARAMETER	MINIMUM DETECTED VALUE (mg/kg)	MAXIMUM DETECTED VALUE (mg/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (mg/kg)	RME (mg/kg)	LOG NORMAL UPPER 95% CONFIDENCE LEVE (mg/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
Aluminum	7620.00	14800.00	WC-SD02-06	10726.33	13079.42	13758.44	6	6	100%
Arsenic	1.10	1.90	*+ HC-SD02-612	3.12	5.54	12.64	2	6	33%
Barium	8.70	28.20	WC-SD02-06	15.30	21.29	25.25	6	6	100%
Beryllium	0.25	0.32	*+ HC-SD02-612	0.29	0.51	137.32	2	2	100%
Cadmium	0.04	0.26	+ WC-SD02-06	0.11	0.17	0.26	6	6	100%
Calcium	1610.00	7860.00	HM-SD02-612	3845.00	5683.04	7847.17	6	6	100%
Chromium	6.00	38.40	+ HM-SD02-06	17.55	28.53	68.22	6	6	100%
Cobalt	3.50	4.40	+ HM-SD02-06	2.18	3.35	5.20	2	6	33%
Copper	0.81	1.50	*+ HC-SD02-06	2.24	3.54	5.85	2	5	40%
Iron	3660.00	32400.00	+ HM-SD02-612	13201.67	21966.51	53108.97	6	6	100%
Lead	6.00	16.90	+ WC-SD02-06	10.50	16.35	5555.00	4	4	100%
Magnesium	618.00	4940.00	+ HM-SD02-06	2123.00	3438.32	6614.15	6	6	100%
Manganese	4.90	67.20	+ HM-SD02-06	33.38	54.41	357.04	6	6	100%
Mercury	0.27	0.40	*+ WC-SD02-06	0.34	0.40	2.78	4	4	100%
Nickel	1.80	11.20	+ HM-SD02-06	5.58	8.93	24.88	5	6	83%
Potassium	623.00	1600.00	+ HM-SD02-612	759.00	1280.80	3459.84	3	6	50%
Selenium	0.40	0.60	*+ HC-SD02-06	0.44	0.62	0.85	3	6	50%
Sodium	1630.00	2750.00	*+ HC-SD02-06	2190.00	5725.84	1.53E+83	2	2	100%
Thallium	0.28	0.42	+ HC-SD02-06	0.27	0.38	0.56	4	6	67%
Vanadium	7.00	30.00	+ HM-SD02-612	17.57	25.67	41.76	6	6	100%
Zinc	27.80	52.00	+ WC-SD02-06	27.38	43.72	355.99	4	6	67%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

## APPENDIX N (CONT.)

BACKGROUND - MIDSTREAM

SEDIMENT - PESTICIDE/PCBs

STATISTICAL SUMMARY OF ANALYTICAL RESULTS

PARAMETER	MINIMUM DETECTED VALUE (ug/kg)	MAXIMUM DETECTED VALUE (ug/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/kg)	RME (ug/kg)	LOG NORMAL UPPER 95%		No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
						CONFIDENCE LEVE (ug/kg)	No. OF TIMES DETECTED			
delta-BHC	0.79	0.79	*+ WC-SD02-612	2.03	2.67	3.61	1	6	17%	
Aldrin	1.20	1.20	*+ WC-SD02-06	1.99	2.50	2.81	1	6	17%	
Dieldrin	3.70	3.70	*+ WC-SD02-06	4.10	4.92	5.23	1	6	17%	
4,4'-DDE	16.00	16.00	WC-SD02-06	6.15	10.20	13.96	1	6	17%	
4,4'-DDD	1.50	12.00	+ WC-SD02-06	4.84	8.05	19.53	2	6	33%	
4,4'-DDT	1.10	2.60	*+ WC-SD02-06	3.48	4.83	7.47	2	6	33%	
Endrin aldehyde	0.77	7.10	+ HC-SD02-06	4.50	6.25	18.84	2	6	33%	

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

## APPENDIX N (CONT.)

## BACKGROUND - DOWNSTREAM

## SEDIMENT - METALS

## STATISTICAL SUMMARY OF ANALYTICAL RESULTS

PARAMETER	MINIMUM DETECTED VALUE (mg/kg)	MAXIMUM DETECTED VALUE (mg/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (mg/kg)	RME (mg/kg)	LOG NORMAL UPPER 95% CONFIDENCE LEVE (mg/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION	
Aluminum	3120.00	14600.00	** WC-SD03-612	9864.29	12793.35	86863.15	7	7	100%	
Barium	10.20	19.20	+	WC-SD03-612	12.44	16.57	35.31	6	7	86%
Cadmium	0.03	0.66	+	HC-SD03-06	0.14	0.31	0.68	6	7	86%
Calcium	2000.00	3380.00	+	HC-SD03-06	2932.86	3371.27	3536.80	7	7	100%
Chromium	16.10	42.60	+	WC-SD03-612	30.87	38.48	43.57	7	7	100%
Cobalt	3.90	5.00	+	HC-SD03-612	3.20	4.30	5.76	4	7	57%
Iron	1700.00	20700.00	+	WC-SD03-612	12868.57	17941.12	46249.04	7	7	100%
Lead	3.70	9.20	+	HM-SD03-06	5.75	7.26	15.49	6	6	100%
Magnesium	4130.00	6540.00	HC-SD03-612	5081.43	5807.10	5894.11	7	7	100%	
Manganese	17.10	64.70	+	HC-SD03-612	45.66	57.52	71.47	7	7	100%
Mercury	0.23	0.42	HC-SD03-612	0.30	0.35	0.36	7	7	100%	
Nickel	5.50	14.20	HM-SD03-06	10.21	12.28	13.51	7	7	100%	
Potassium	1250.00	1840.00	HC-SD03-612	1570.00	1729.32	1753.19	7	7	100%	
Thallium	0.24	0.44	HC-SD03-612	0.25	0.32	0.34	4	7	57%	
Vanadium	18.40	36.90	HC-SD03-612	26.59	31.50	32.88	7	7	100%	
Zinc	20.80	40.00	HC-SD03-612	30.66	35.18	36.50	7	7	100%	

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

## APPENDIX N (CONT.)

BACKGROUND - DOWNSTREAM

SEDIMENT - PESTICIDES/PCBs

STATISTICAL SUMMARY OF ANALYTICAL RESULTS

PARAMETER	MINIMUM DETECTED VALUE (ug/kg)	MAXIMUM DETECTED VALUE (ug/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/kg)	RME (ug/kg)	LOG NORMAL UPPER 95% CONFIDENCE LEVE (ug/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
4,4'-DDD	1.10	4.00	*+ HC-SD03-612	3.38	4.56	6.24	4	7	57%
4,4'-DDT	0.76	1.20	*+ HC-SD03-06D	4.12	5.76	14.23	2	7	29%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HADNOT CREEK  
 SEDIMENT - METALS

PARAMETER	MINIMUM DETECTED VALUE (mg/kg)	MAXIMUM DETECTED VALUE (mg/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (mg/kg)	RME (mg/kg)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (mg/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION	
Aluminum	780.00	14000.00	+	HC-SD03-612	5467.78	8305.91	20353.32	9	9	100%
Arsenic	0.26	1.90	*+	HC-SD02-612	1.71	2.67	8.56	6	9	67%
Barium	4.10	17.20	+	HC-SD03-612	9.75	13.11	21.84	8	9	89%
Beryllium	0.14	0.32	+	HC-SD02-612	0.16	0.24	4.60	3	6	50%
Cadmium	0.03	0.66		HC-SD03-06	0.11	0.24	0.42	7	9	78%
Calcium	1030.00	3620.00	+	HC-SD01-06	2645.56	3233.82	3840.09	9	9	100%
Chromium	1.30	41.60	+	HC-SD03-612	10.81	18.97	53.55	9	9	100%
Cobalt	4.50	5.00		HC-SD03-612	1.87	2.91	4.01	2	9	22%
Copper	0.66	1.50	*+	HC-SD02-06	1.35	1.75	2.01	6	9	67%
Iron	382.00	11100.00	+	HC-SD03-06D	3396.56	5709.65	28323.00	9	9	100%
Lead	3.70	5.30	*+	HC-SD03-06	4.50	9.55	305.02	2	2	100%
Magnesium	77.10	6540.00	+	HC-SD03-612	1977.79	3486.31	1292043.17	7	9	78%
Manganese	3.50	64.70		HC-SD03-612	16.54	29.38	62.63	9	9	100%
Mercury	0.25	0.42	*+	HC-SD03-612	0.34	0.48	11.17	3	3	100%
Nickel	1.80	12.10	+	HC-SD03-612	3.77	6.49	17.25	4	9	44%
Potassium	623.00	1840.00	+	HC-SD03-612	671.39	1079.26	2769.97	4	9	44%
Selenium	0.21	0.60		HC-SD02-06	0.30	0.39	0.48	5	9	56%
Sodium	1630.00	2750.00	+	HC-SD02-06	845.25	1750.35	183541390882.91	2	6	33%
Thallium	0.14	0.44	+	HC-SD03-612	0.23	0.31	0.46	6	9	67%
Vanadium	1.50	36.90	+	HC-SD03-612	11.11	18.54	56.26	9	9	100%
Zinc	20.80	40.00	+	HC-SD03-612	12.71	22.07	63.76	3	9	33%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HADNOT CREEK  
 SEDIMENT - PESTICIDES/PCBs

PARAMETER	MINIMUM DETECTED VALUE (ug/kg)	MAXIMUM DETECTED VALUE (ug/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/kg)	RME (ug/kg)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
beta-BHC	1.70	1.70	*+ HC-SD04-612	1.93	2.39	2.58	1	9	11%
delta-BHC	0.64	0.64	*+ HC-SD01-06	1.82	2.35	2.91	1	9	11%
Heptachlor	0.48	2.00	*+ HC-SD04-612	1.89	2.42	3.26	2	9	22%
4,4'-DDD	1.50	4.00	HC-SD03-612	2.16	3.11	3.50	3	9	33%
4,4'-DDT	1.20	1.20	*+ HC-SD03-06D	3.23	4.23	5.08	1	9	11%
Methoxychlor	0.94	0.94	*+ HC-SD04-06	17.66	23.58	92.52	1	9	11%
Endrin aldehyde	0.59	7.10	+ HC-SD02-06	3.56	5.02	10.80	3	9	33%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HADNOT CREEK  
 SEDIMENT - SEMIVOLATILE ORGANIC COMPOUNDS

PARAMETER	MINIMUM DETECTED VALUE	MAXIMUM DETECTED VALUE	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/kg)	RME (ug/kg)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
	(ug/kg)	(ug/kg)							
NO SEMIVOLATILE ORGANIC COMPOUNDS WERE DETECTED									

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HADNOT CREEK  
 SEDIMENT - VOLATILE ORGANIC COMPOUNDS

PARAMETER	MINIMUM DETECTED VALUE (ug/kg)	MAXIMUM DETECTED VALUE (ug/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/kg)	RME (ug/kg)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
Acetone	70.00	70.00	HC-SD01-06	18.06	30.44	36.73	1	9	11%
Carbon Disulfide	14.00	19.00	HC-SD02-612	12.44	15.67	18.14	2	9	22%
2-Butanone	7.00	7.00	*+ HC-SD01-06	11.06	13.94	15.49	1	9	11%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HOLLAND MILL CREEK  
 SEDIMENT - METALS

PARAMETER	MINIMUM DETECTED VALUE (mg/kg)	MAXIMUM DETECTED VALUE (mg/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (mg/kg)	RME (mg/kg)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (mg/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION	
Aluminum	337.00	13600.00	+	HM-SD02-06	6181.29	10282.21	655067.62	7	7	100%
Barium	11.00	18.70	+	HM-SD02-06	8.71	13.92	68.49	4	7	57%
Cadmium	0.03	0.11		HM-SD01-06D	0.06	0.08	0.10	7	7	100%
Calcium	282.00	7860.00	+	HM-SD02-612	2952.86	4844.12	22431.34	7	7	100%
Chromium	1.10	38.40	+	HM-SD02-06	19.63	32.39	2021.73	7	7	100%
Cobalt	4.00	4.40	+	HM-SD02-06	2.02	3.18	6.18	2	7	29%
Iron	225.00	32400.00	+	HM-SD02-612	12262.43	21399.01	27918943.98	7	7	100%
Lead	0.62	9.20	+	HM-SD03-06	4.35	6.94	32.96	7	7	100%
Magnesium	26.70	5700.00	+	HM-SD03-06	2576.66	4422.69	136198282.35	7	7	100%
Manganese	1.30	67.20	+	HM-SD02-06	34.14	56.82	8851.72	7	7	100%
Mercury	0.09	0.35	+	HM-SD03-06	0.23	0.30	0.38	7	7	100%
Nickel	9.60	14.20	+	HM-SD03-06	6.76	11.07	359.48	4	7	57%
Potassium	1510.00	1760.00	+	HM-SD03-612	1007.00	1596.65	13233.89	4	7	57%
Selenium	0.25	0.40		HM-SD02-06	0.21	0.29	0.39	2	7	29%
Silver	0.49	0.49	*+	HM-SD01-06	0.39	0.49	0.60	1	7	14%
Thallium	0.13	0.37	+	HM-SD02-06	0.20	0.29	0.52	4	7	57%
Vanadium	0.66	30.00	+	HM-SD02-612	16.69	27.76	18094.26	6	7	86%
Zinc	6.70	43.10	+	HM-SD02-06	23.57	34.53	65.13	7	7	100%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HOLLAND MILL CREEK  
 SEDIMENT - PESTICIDES/PCBs

PARAMETER	MINIMUM DETECTED VALUE (ug/kg)	MAXIMUM DETECTED VALUE (ug/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/kg)	RME (ug/kg)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
beta-BHC	3.80	7.30	HM-SD01-06D	3.24	4.69	5.98	2	7	29%
Aldrin	0.56	0.72	*+ HM-SD01-612	1.84	2.60	4.20	2	7	29%
Dieldrin	0.58	1.50	*+ HM-SD01-612	3.55	5.13	12.37	2	7	29%
4,4'-DDE	1.00	4.30	*+ HM-SD01-612	4.01	5.37	8.82	2	7	29%
4,4'-DDD	0.87	3.10	*+ HM-SD01-612	2.85	4.16	6.44	4	7	57%
4,4'-DDT	1.70	1.70	*+ HM-SD01-612	3.79	5.13	6.75	1	7	14%
alpha-Chlordane	1.30	1.30	*+ HM-SD01-612	1.99	2.61	3.14	1	7	14%
gamma-Chlordane	3.00	3.00	+ HM-SD01-612	2.24	2.86	3.56	1	7	14%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HOLLAND MILL CREEK  
 SEDIMENT - SEMIVOLATILE ORGANIC COMPOUNDS

PARAMETER	MINIMUM	MAXIMUM	SAMPLE No.	LOG NORMAL					No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
	DETECTED VALUE (ug/kg)	DETECTED VALUE (ug/kg)	OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/kg)	RME (ug/kg)	CONFIDENCE LEVEL (ug/kg)	No. OF TIMES DETECTED			
Di-n-butylphthalate	534.00	619.00	+ HM-SD02-612	423.29	573.31	766.73	3	7	43%	
bis(2-Ethylhexyl)phthalate	454.00	454.00	*+ HM-SD03-612	378.64	500.04	607.73	1	7	14%	

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - HOLLAND MILL CREEK  
 SEDIMENT - VOLATILE ORGANIC COMPOUNDS

PARAMETER	MINIMUM DETECTED	MAXIMUM DETECTED	SAMPLE No.	LOG NORMAL			No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
	VALUE (ug/kg)	VALUE (ug/kg)	OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/kg)	RME (ug/kg)	CONFIDENCE LEVEL UPPER 95% (ug/kg)		
NO VOLATILE ORGANIC COMPOUNDS WERE DETECTED								

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - WEBB CREEK  
 SEDIMENT - METALS

PARAMETER	MINIMUM DETECTED VALUE (mg/kg)	MAXIMUM DETECTED VALUE (mg/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (mg/kg)	RME (mg/kg)	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (mg/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
Aluminum	8200.00	14800.00	*+ WC-SD02-06	12275.00	15932.10	19239.95	4	4	100%
Barium	13.30	28.20	+ WC-SD02-06	18.83	26.76	35.92	4	4	100%
Cadmium	0.06	0.26	+ WC-SD02-06	0.13	0.24	1.11	4	4	100%
Calcium	2190.00	4060.00	*+ WC-SD02-06	3222.50	4132.21	4914.08	4	4	100%
Chromium	8.70	42.60	+ WC-SD03-612	24.93	42.26	246.57	4	4	100%
Cobalt	3.50	3.90	*+ WC-SD03-612	2.44	4.16	21.71	2	4	50%
Iron	8120.00	20700.00	+ WC-SD03-612	13980.00	20133.62	29586.84	4	4	100%
Lead	5.10	16.90	+ WC-SD02-06	9.85	16.48	51.03	4	4	100%
Magnesium	618.00	6060.00	*+ WC-SD03-612	3197.00	6127.63	817766.37	4	4	100%
Manganese	26.00	47.80	*+ WC-SD03-612	39.35	50.44	60.95	4	4	100%
Mercury	0.23	0.40	*+ WC-SD02-06	0.31	0.41	0.48	4	4	100%
Nickel	3.80	11.40	+ WC-SD03-612	7.25	11.11	21.80	4	4	100%
Potassium	1410.00	1590.00	*+ WC-SD03-612	905.88	1719.51	81148.45	2	4	50%
Thallium	0.24	0.24	+ WC-SD03-06	0.16	0.23	0.31	1	4	25%
Vanadium	11.90	31.00	+ WC-SD03-612	21.33	30.50	45.84	4	4	100%
Zinc	27.20	52.00	+ WC-SD02-06	33.83	48.09	61.59	4	4	100%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - WEBB CREEK  
 SEDIMENT - PESTICIDES/PCBs

PARAMETER	MINIMUM DETECTED VALUE	MAXIMUM DETECTED VALUE	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE	RME	LOG NORMAL UPPER 95%	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
	(ug/kg)	(ug/kg)		(ug/kg)	(ug/kg)	(ug/kg)			
delta-BHC	0.79	0.79	*+ WC-SD02-612	1.99	3.02	9.99	1	4	25%
Aldrin	1.20	1.20	*+ WC-SD02-06	1.93	2.65	3.66	1	4	25%
Dieldrin	3.70	3.70	*+ WC-SD02-06	4.00	4.79	4.98	1	4	25%
4,4'-DDE	16.00	16.00	+ WC-SD02-06	7.08	14.12	97.81	1	4	25%
4,4'-UDD	12.00	12.00	+ WC-SD02-06	6.08	10.78	28.91	1	4	25%
4,4'-DDT	0.76	2.60	*+ WC-SD02-06	2.37	4.64	91.00	3	4	75%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - WEBB CREEK  
 SEDIMENT - SEMIVOLATILE ORGANIC COMPOUNDS

PARAMETER	MINIMUM DETECTED VALUE (ug/kg)	MAXIMUM DETECTED VALUE (ug/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/kg)	RME	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
Benzo(a)pyrene	544.00	544.00	*+ WC-SD03-612	436.25	554.81	635.17	1	4	25%

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

MARINE CORPS BASE CAMP LEJEUNE  
 STATISTICAL SUMMARY OF ANALYTICAL RESULTS  
 BACKGROUND - WEBB CREEK  
 SEDIMENT - VOLATILE ORGANIC COMPOUNDS

PARAMETER	MINIMUM DETECTED VALUE (ug/kg)	MAXIMUM DETECTED VALUE (ug/kg)	SAMPLE No. OF MAXIMUM DETECTED VALUE	ARITHMETIC AVERAGE (ug/kg)	RME	LOG NORMAL UPPER 95% CONFIDENCE LEVEL (ug/kg)	No. OF TIMES DETECTED	No. OF TIMES ANALYZED	FREQUENCY OF DETECTION
NO VOLATILE ORGANIC COMPOUNDS WERE DETECTED									

\* = THE RME IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

+ = THE LOG NORMAL 95% UCL IS GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

\*+ = BOTH THE RME AND LOG NORMAL 95% UCL ARE GREATER THAN THE MAXIMUM DETECTED VALUE; THEREFORE, THE MAXIMUM VALUE IS USED TO CALCULATE CHRONIC DAILY INTAKE

RME = REASONABLE MAXIMUM EXPOSURE

NA = NOT APPLICABLE

**Analytical Summary of Results  
(Surface Water)**

MARINE CORPS BASE CAMP LEJEUNE  
 ANALYTICAL SUMMARY OF RESULTS  
 BACKGROUND - HADNOT CREEK  
 SURFACE WATER - METALS

BAKER I.D.	HC-SW01	HC-SW02	HC-SW03	HC-SW03D	HC-SW04
LABORATORY I.D.	5167-16	5162	5166	5163	5152
DATE COLLECTED	08-MAY-1994	06-MAY-1994	06-MAY-1994	06-MAY-1994	08-MAY-1994
UNITS	UG/L	UG/L	UG/L	UG/L	UG/L
Aluminum	356 U	303 U	301 U	187 U	692
Arsenic	1 U	1 UJ	20	10 UJ	1 U
Barium	19 J	20 J	26 J	24 J	9 J
Calcium	27000	36600	86600	107000	11600
Chromium	9 U	19 U	130 J	125 J	9 U
Iron	746	528	339	291	556
Magnesium	1450	44800	633000	613000	954
Potassium	1670 U	14500	203000	202000	1670 U
Selenium	1 U	5 U	6 J	1 UJ	1 UJ
Sodium	6900	383000	2090000	2560000	6090

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - HADNOT CREEK  
SURFACE WATER PESTICIDES AND PCBs

BAKER I.D.	HC-SW01	HC-SW02	HC-SW03	HC-SW03D	HC-SW04
LABORATORY I.D.	5167-16	5162	5166	5163	5152
DATE COLLECTED	08-MAY-1994	06-MAY-1994	06-MAY-1994	06-MAY-1994	08-MAY-1994
UNITS	ug/l	ug/l	ug/l	ug/l	ug/l

NO PESTICIDES OR PCBs WERE DETECTED

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - HADNOT CREEK  
SURFACE WATER - SEMIVOLATILE ORGANIC COMPOUNDS

BAKER I.D.	HC-SW01	HC-SW02	HC-SW03	HC-SW03D	HC-SW04
LABORATORY I.D.	5167-16	5162	5166	5163	5152
DATE COLLECTED	08-MAY-1994	06-MAY-1994	06-MAY-1994	06-MAY-1994	08-MAY-1994
UNITS	ug/l	ug/l	ug/l	ug/l	ug/l

NO SEMIVOLATILE ORGANIC COMPOUNDS WERE DETECTED

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - HADNOT CREEK  
SURFACE WATER - VOLATILE ORGANIC COMPOUNDS

BAKER I.D.	HC-SW01	HC-SW02	HC-SW03	HC-SW03D	HC-SW04
LABORATORY I.D.	5167-16	5162	5166	5163	5152
DATE COLLECTED	08-MAY-1994	06-MAY-1994	06-MAY-1994	06-MAY-1994	08-MAY-1994
UNITS	ug/l	ug/l	ug/l	ug/l	ug/l

NO VOLATILE ORGANIC COMPOUNDS WERE DETECTED

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - HOLLAND MILL CREEK  
SURFACE WATER - METALS

BAKER I.D.	HM-SW01	HM-SW02	HM-SW03
LABORATORY I.D.	5167-18	5161	5160
DATE COLLECTED	08-MAY-1994	06-MAY-1994	06-MAY-1994
UNITS	UG/L	UG/L	UG/L
Aluminum	259 U	535 J	288 U
Barium	49 J	38 J	20 J
Calcium	14100	40200	302000
Chromium	10 U	36 J	158 J
Iron	425	559	320
Lead	1 U	2.5 U	58.1
Magnesium	2830	109000	754000
Potassium	1670 U	41100	288000
Selenium	1.5 J	5 U	41 J
Silver	10 U	17 U	37 J
Sodium	16500	739000	6750000

MARINE CORPS BASE CAMP LEJEUNE  
ANLAYTICAL SUMMARY OF RESULTS  
BACKGROUND - HOLLAND MILL CREEK  
SURFACE WATER - PESTICIDES AND PCBs

	HM-SW01	HM-SW02	HM-SW03
BAKER I.D.	5167-18	5161	5160
LABORATORY I.D.			
DATE COLLECTED	08-MAY-1994	06-MAY-1994	06-MAY-1994
UNITS	ug/l	ug/l	ug/l

NO PESTICIDES OR PCBs WERE DETECTED

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - HOLLAND MILL CREEK  
SURFACE WATER - SEMIVOLATILE ORGANIC COMPOUNDS

BAKER I.D.	HM-SW01	HM-SW02	HM-SW03
LABORATORY I.D.	5167-18	5161	5160
DATE COLLECTED	08-MAY-1994	06-MAY-1994	06-MAY-1994
UNITS	ug/l	ug/l	ug/l

NO SEMIVOLATILE ORGANIC COMPOUNDS WERE DETECTED

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - HOLLAND MILL CREEK  
SURFACE WATER - VOLATILE ORGANIC COMPOUNDS

BAKER I.D.	HM-SW01	HM-SW02	HM-SW03
LABORATORY I.D.	5167-18	5161	5160
DATE COLLECTED	08-MAY-1994	06-MAY-1994	06-MAY-1994
UNITS	ug/l	ug/l	ug/l

NO VOLATILE ORGANIC COMPOUNDS WERE DETECTED

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - WEBB CREEK  
SURFACE WATER - METALS

BAKER I.D.	WC-SW02	WC-SW03
LABORATORY I.D.	5167-8	5158
DATE COLLECTED	06-MAY-1994	06-MAY-1994
UNITS	UG/L	UG/L
Barium	29 J	27 J
Calcium	46900	40500
Chromium	15 U	97 J
Iron	660	321
Magnesium	29000	44800
Potassium	10900	136000
Sodium	202000	895000

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - WEBB CREEK  
SURFACE WATER - PESTICIDES AND PCBs

	WC-SW02	WC-SW03
BAKER I.D.	5167-8	5158
LABORATORY I.D.		
DATE COLLECTED	06-MAY-1994	06-MAY-1994
UNITS	ug/l	ug/l
Aldrin	0.035 J	0.05 U

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - WEBB CREEK  
SURFACE WATER - SEMIVOLATILE ORGANIC COMPOUNDS

BAKER I.D.	WC-SW02	WC-SW03
LABORATORY I.D.	5167-8	5158
DATE COLLECTED	06-MAY-1994	06-MAY-1994
UNITS	ug/l	ug/l

NO SEMIVOLATILE ORGANIC COMPOUNDS WERE DETECTED

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - WEBB CREEK  
SURFACE WATER - VOLATILE ORGANIC COMPOUNDS

BAKER I.D.	WC-SW02	WC-SW03
LABORATORY I.D.	5167-8	5158
DATE COLLECTED	06-MAY-1994	06-MAY-1994
UNITS	ug/l	ug/l

NO VOLATILE ORGANIC COMPOUNDS WERE DETECTED

**Analytical Summary of Results  
(Sediment)**

MARINE CORPS BASE CAMP LEJEUNE  
 ANALYTICAL SUMMARY OF RESULTS  
 BACKGROUND - HADNOT CREEK  
 SEDIMENT - METALS

BAKER I.D.	HC-SD01-06	HC-SD01-612	HC-SD02-06	HC-SD02-612	HC-SD03-06	HC-SD03-06D	HC-SD03-612	HC-SD04-06	HC-SD04-612
LABORATORY I.D	5050	5044	5057-2	5054	5238	5237	5236	5052	5051
DATE COLLECTED	8-MAY-1994	8-MAY-1994	6-MAY-1994	6-MAY-1994	07-MAY-1994	07-MAY-1994	07-MAY-1994	8-MAY-1994	8-MAY-1994
UNITS	MG/KG	MG/KG	MG/KG	MG/KG	MG/KG	MG/KG	MG/KG	MG/KG	MG/KG
Aluminum	2940 J	1880 J	7820 J	10100 J	3120 J	7310 J	14000 J	780 J	1260 J
Arsenic	0.46 J	0.28 J	1.1 J	1.9 J	7.5 U	6.5 U	7.9 U	0.45 J	0.26 J
Barium	16.3 J	14.6 J	9.2 J	8.7 J	3.9 U	10.2	17.2	4.1 J	5.5 J
Beryllium	0.14 J	0.16 U	0.25 J	0.32 J	0.95 R	0.92 R	1.3 R	0.13 U	0.15 U
Cadmium	0.03 J	0.03 J	0.1 J	0.04 J	0.66	0.08	0.04 U	0.03 J	0.03 UJ
Calcium	3620 J	3330 J	2030 J	1610 J	3380 J	3350 J	3310 J	1030 J	2150 J
Chromium	2.3	3.2	6	6	16.1	18.8	41.6	2	1.3
Cobalt	1.6 U	1.8 U	2.7 U	1.8 U	3.7 U	4.5	5	1.5 U	1.6 U
Copper	1	1.1	1.5	0.81	4.9 U	4.3 U	3.5 U	0.66	0.73
Iron	648	586	3660	4630	7280 J	11100 J	1700 J	382	583
Lead	0.77 R	0.88 R	1.1 R	7.1 R	5.3	3.7	8.6 R	1 R	1.1 R
Magnesium	87.7	77.1	1450	1040	4420	4130	6540	48.2 U	62.5 U
Manganese	6.9	6.5	6.5	4.9	17.1	35.1	64.7	3.7	3.5
Mercury	0.19 R	0.13 R	0.42 R	0.24 R	0.34	0.25	0.42	0.11 R	0.08 R
Nickel	1.6 U	1.8 U	2.7 U	1.8	9.9	5.5	12.1	1.5 U	1.6 U
Potassium	349 U	396 U	623	395 U	1420	1250	1840	324 U	355 U
Selenium	0.27 J	0.34 J	0.6 J	0.47 J	0.48 UJ	0.41 UJ	0.51 UJ	0.21 J	0.2 UJ
Sodium	339 U	385 U	2750	1630	14100 R	9860 R	6620 R	315 U	344 U
Thallium	0.14	0.16	0.42	0.28	0.34 U	0.29	0.44	0.13 U	0.15 U
Vanadium	2.6	2.8	8.4	7	20.5	18.4	36.9	1.5	1.9
Zinc	4.9 U	4.5 U	9.7 U	6.6 U	20.8	34.3	40	4.5 U	8.3 U

MARINE COPRS BASE CAMP LEJEUNE  
 ANALYTICAL SUMMARY OF RESULTS  
 BACKGROUND - HADNOT CREEK  
 SEDIMENT - PESTICIDES AND PCBs

BAKER I.D.	HC-SD01-06	HC-SD01-612	HC-SD02-06	HC-SD02-612	HC-SD03-06	HC-SD03-06D	HC-SD03-612	HC-SD04-06	HC-SD04-612
LABORATORY I.D.	5057-7	5044	5055	5054	5238	5237	5236	5052	5051
DATE COLLECTED	8-MAY-1994	8-MAY-1994	6-MAY-1994	6-MAY-1994	07-MAY-1994	07-MAY-1994	07-MAY-1994	8-MAY-1994	8-MAY-1994
UNITS	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg
beta-BHC	2.4 U	2.8 U	4.2 U	2.8 U	5.8 U	4.9 U	6.2 U	2.3 U	1.7 J
delta-BHC	0.64 J	2.8 U	4.2 U	2.8 U	5.8 U	4.9 U	6.2 U	2.3 U	2.5 U
Heptachlor	0.48 J	2.8 U	4.2 U	2.8 U	5.8 U	4.9 U	6.2 U	2.3 U	2 J
4,4'-DDD	2.4 U	2.8 U	1.5 J	2.8 U	11 U	2 J	4 J	2.3 U	2.5 U
4,4'-DDT	4.7 U	5.4 U	8.2 U	5.3 U	11 U	1.2 J	12 U	4.4 U	4.8 U
Methoxychlor	24 U	28 U	42 U	28 U	58 U	49 U	62 U	0.94 J	25 U
Endrin aldehyde	0.59 J	5.4 U	7.1 J	0.77 J	11 U	9.6 U	12 U	4.4 U	4.8 U

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - HADNOT CREEK  
SEDIMENT - SEMIVOLATILE ORGANIC COMPOUNDS

BAKER I.D.	HC-SD01-06	HC-SD01-612	HC-SD02-06	HC-SD02-612	HC-SD03-06	HC-SD03-06D	HC-SD03-612	HC-SD04-06	HC-SD04-612
LABORATORY I.D.	5057-7	5044	5055	5054	5238	5237	5236	5052	5051
DATE COLLECTED	8-MAY-1994	8-MAY-1994	6-MAY-1994	6-MAY-1994	07-MAY-1994	07-MAY-1994	07-MAY-1994	8-MAY-1994	8-MAY-1994
UNITS	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg

NO SEMIVOLATILE ORGANIC COMPOUNDS WERE DETECTED

MARINE CORPS BASE CAMP LEJEUNE  
 ANALYTICAL SUMMARY OF RESULTS  
 BACKGROUND - HADNOT CREEK  
 SEDIMENT - VOLATILE ORGANIC COMPOUNDS

BAKER I.D.	HC-SD01-06	HC-SD01-612	HC-SD02-06	HC-SD02-612	HC-SD03-06	HC-SD03-06D	HC-SD03-612	HC-SD04-06	HC-SD04-612
LABORATORY I.D.	5057-7	5044	5055	5054	5238	5237	5236	5052	5051
DATE COLLECTED	8-MAY-1994	8-MAY-1994	6-MAY-1994	6-MAY-1994	07-MAY-1994	07-MAY-1994	07-MAY-1994	8-MAY-1994	8-MAY-1994
UNITS	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg	ug/kg
Acetone	70 J	16 UJ	25 UJ	16 UJ	34 UJ	29 UJ	37 UJ	13 UJ	15 UJ
Carbon Disulfide	14 U	16 U	14	19 J	34 U	29 U	37 U	13 U	15 U
2-Butanone	7 J	16 UJ	25 UJ	16 UJ	34 UJ	29 UJ	37 UJ	13 UJ	15 UJ

MARINE CORPS BASE CAMP LEJEUNE  
 ANALYTICAL SUMMARY OF RESULTS  
 BACKGROUND - HOLLAND MILL CREEK  
 SEDIMENT - METALS

BAKER I.D. LABORATORY I.D. DATE COLLECTED UNITS	HM-SD01-06 5243-18 08-MAY-1994 MG/KG	HM-SD01-06D 5220 08-MAY-1994 MG/KG	HM-SD01-612 5219 08-MAY-1994 MG/KG	HM-SD02-06 5242 07-MAY-1994 MG/KG	HM-SD02-612 5241 07-MAY-1994 MG/KG	HM-SD03-06 5240 07-MAY-1994 MG/KG	HM-SD03-612 5239 07-MAY-1994 MG/KG
Aluminum	457 J	337 J	505 J	13600 J	9850 J	8760 J	9760 J
Barium	3.4 U	2.1 U	3.9 U	18.7	13.7	11	12.9
Cadmium	0.03	0.11	0.03	0.08	0.06	0.05	0.03
Calcium	282 J	508 J	2850 J	4250 J	7860 J	2920 J	2000 J
Chromium	1.6	1.1	1.5	38.4	28.1	30.7	36
Cobalt	1.3 U	1.4 U	1.4 U	4.4	3.5 U	3.9 U	4
Iron	262 J	225 J	350 J	15800 J	32400 J	16900 J	19900 J
Lead	0.62 J	0.74 J	1	6	7.2	9.2	5.7
Magnesium	35.5	26.7	34.4	4940	3000	5700	4300
Manganese	1.9	1.3	1.6	67.2	55.5	50.2	61.3
Mercury	0.09	0.16	0.18	0.27	0.32	0.35	0.27
Nickel	1.3 U	1.4 U	1.4 U	11.2	9.6	14.2	10.3
Potassium	297 U	304 U	317 U	1510	1600	1720	1760
Selenium	0.17 U	0.17 U	0.25 J	0.4 J	0.45 UJ	0.5 UJ	0.37 UJ
Silver	0.49	0.37 U	0.39 U	0.85 U	0.95 U	1.1 U	0.79 U
Thallium	0.12 U	0.12 U	0.13	0.37	0.32	0.35 U	0.27
Vanadium	0.84	0.62 U	0.66	27.1	30	28.4	29.5
Zinc	9.7	6.7	8.3	43.1	33.2	34.1	29.9

MARINE CORPS BASE CAMP LEJEUNE  
 ANALYTICAL SUMMARY OF RESULTS  
 BACKGROUND - HOLLAND MILL CREEK  
 SEDIMENT - PESTICIDES AND PCBs

BAKER I.D.	HM-SD01-06	HM-SD01-06D	HM-SD01-612	HM-SD02-06	HM-SD02-612	HM-SD03-06	HM-SD03-612
LABORATORY I.D.	5243-18	5220	5219	5242	5241	5240	5239
DATE COLLECTED	08-MAY-1994	08-MAY-1994	08-MAY-1994	07-MAY-1994	07-MAY-1994	07-MAY-1994	07-MAY-1994
UNITS	ug/kg						
beta-BHC	2.1 UJ	7.3 J	3.8	5.1 U	5.5 U	6 U	4.5 U
Aldrin	2.1 U	0.56 J	0.72 J	5.1 U	5.5 U	6 U	4.5 U
Dieldrin	4 U	0.58 J	1.5 J	9.8 U	11 U	12 U	8.8 U
4,4'-DDE	4 U	1 J	4.3	9.8 U	11 U	12 U	8.8 U
4,4'-DDD	4 U	0.87 J	3.1	9.8 U	11 U	2.5 J	1.1 J
4,4'-DDT	4 U	4.1 U	1.7 J	9.8 U	11 U	12 U	8.8 U
alpha-Chlordane	2.1 U	2.1 U	1.3 J	5.1 U	5.5 U	6 U	4.5 U
gamma-Chlordane	2.1 U	2.1 U	3	5.1 U	5.5 U	6 U	4.5 U

MARINE CORPS BASE CAMP LEJEUNE  
 ANALYTICAL SUMMARY OF RESULTS  
 BACKGROUND - HOLLAND MILL CREEK  
 SEDIMENT - SEMIVOLATILE ORGANIC COMPOUNDS

BAKER I.D.	HM-SD01-06	HM-SD01-06D	HM-SD01-612	HM-SD02-06	HM-SD02-612	HM-SD03-06	HM-SD03-612
LABORATORY I.D.	5243-18	5220	5219	5242	5241	5240	5239
DATE COLLECTED	08-MAY-1994	08-MAY-1994	08-MAY-1994	07-MAY-1994	07-MAY-1994	07-MAY-1994	07-MAY-1994
UNITS	ug/kg						
Di-n-butylphthalate	401 U	412 U	429 U	614 J	619 J	1150 U	534 J
bis(2-Ethylhexyl)phthalate	401 UJ	412 UJ	429 UJ	943 U	1058 U	1150 U	454 J

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - HOLLAND MILL CREEK  
SEDIMENT - VOLATILE ORGANIC COMPOUNDS

BAKER I.D.	HM-SD01-06	HM-SD01-06D	HM-SD01-612	HM-SD02-06	HM-SD02-612	HM-SD03-06	HM-SD03-612
LABORATORY I.D.	5243-18	5220	5219	5242	5241	5240	5239
DATE COLLECTED	08-MAY-1994	08-MAY-1994	08-MAY-1994	07-MAY-1994	07-MAY-1994	07-MAY-1994	07-MAY-1994
UNITS	ug/kg						

NO VOLATILE ORGANIC COMPOUNDS WERE DETECTED

MARINE CORPS BASE CAMP LEJEUNE  
 ANALYTICAL SUMMARY OF RESULTS  
 BACKGROUND - WEBB CREEK  
 SEDIMENT - METALS

BAKER I.D.	WC-SD02-06	WC-SD02-612	WC-SD03-06	WC-SD03-612
LABORATORY I.D.	5243-10	5232	5235	5234
DATE COLLECTED	06-MAY-1994	06-MAY-1994	07-MAY-1994	07-MAY-1994
UNITS	MG/KG	MG/KG	MG/KG	MG/KG
Aluminum	14800 J	8200	11500 J	14600 J
Barium	28.2	13.3	14.6	19.2
Cadmium	0.26	0.12	0.06	0.07
Calcium	4060 J	3260 J	2190 J	3380 J
Chromium	18.1	8.7	30.3	42.6
Cobalt	3.5	2.3 U	2.4 U	3.9
Iron	14600 J	8120	12500 J	20700 J
Lead	16.9	11.9	5.1	5.5
Magnesium	1690	618	4420	6060
Manganese	40.2	26	43.4	47.8
Mercury	0.4	0.36	0.23	0.26
Nickel	5.7	3.8	8.1	11.4
Potassium	739 U	508 U	1410	1590
Thallium	0.3 U	0.21 U	0.24	0.32 U
Vanadium	21	11.9	21.4	31
Zinc	52	27.8	28.3	27.2

MARINE CORPS BASE CAMP LEJEUNE  
 ANALYTICAL SUMMARY OF RESULTS  
 BACKGROUND - WEBB CREEK  
 SEDIMENT - PESTICIDES AND PCBs

BAKER I.D.	WC-SD02-06	WC-SD02-612	WC-SD03-06	WC-SD03-612
LABORATORY I.D.	5243-10	5232	5235	5234
DATE COLLECTED	06-MAY-1994	06-MAY-1994	07-MAY-1994	07-MAY-1994
UNITS	ug/kg	ug/kg	ug/kg	ug/kg
delta-BHC	5.2 U	0.79 J	3.7 U	5.4 U
Aldrin	1.2 J	3.9 U	3.7 U	5.4 U
Dieldrin	3.7 J	7.5 U	7.1 U	10 U
4,4'-DDE	16	7.5 U	7.1 U	10 U
4,4'-DDD	12	7.5 U	7.1 U	10 U
4,4'-DDT	2.6 J	1.1 J	0.76 J	10 U

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - WEBB CREEK  
SEDIMENT - SEMIVOLATILE ORGANIC COMPOUNDS

BAKER I.D.	WC-SD02-06	WC-SD02-612	WC-SD03-06	WC-SD03-612
LABORATORY I.D.	5243-10	5232	5235	5234
DATE COLLECTED	06-MAY-1994	06-MAY-1994	07-MAY-1994	07-MAY-1994
UNITS	ug/kg	ug/kg	ug/kg	ug/kg
Benzo(a)pyrene	1000 U	688 U	714 U	544 J

MARINE CORPS BASE CAMP LEJEUNE  
ANALYTICAL SUMMARY OF RESULTS  
BACKGROUND - WEBB CREEK  
SEDIMENT - VOLATILE ORGANIC COMPOUNDS

	WC-SD02-06	WC-SD02-612	WC-SD03-06	WC-SD03-612
BAKER I.D.	5243-10	5232	5235	5234
LABORATORY I.D.				
DATE COLLECTED	06-MAY-1994	06-MAY-1994	07-MAY-1994	07-MAY-1994
UNITS	ug/kg	ug/kg	ug/kg	ug/kg

NO VOLATILE ORGANIC COMPOUNDS WERE DETECTED

## Field Chemistry Results

**FIELD CHEMISTRY FROM BIOLOGICAL SAMPLES**  
**HADNOT CREEK, HOLLAND MILL CREEK, AND WEBB CREEK**  
**MCB CAMP LEJEUNE, NORTH CAROLINA**

Sample Identification	Sample Location	Salinity (ppt)	Conductivity (micromhos/cm)	DO (mg/L)	pH (S.U.)	Temperature (deg. C)
HC01-SW/SD-FS/BN	surface bottom	0 NA	13.5 NA	7.7 NA	6.89 NA	17 NA
HC02-SW/SD	surface bottom	0.8 15.5	1,810 21,900	5.9 1.0	6.71 6.73	16.1 18.2
HC02-FS/BN	surface bottom	0.3 13.1	1,200 20,900	NA NA	NA NA	20.5 22
	surface bottom	0 10.5	720 17,200	7.3 1	7.2 6.7	15.5 20
	surface bottom	0 16.5	1,050 22,800	NA NA	NA NA	20.5 21
HC03-SW/SD	surface bottom	17 NA	25,500 NA	12 NA	7.79 NA	17.5 NA
HC03-FS/BN	surface bottom	17.9 NA	26,500 NA	NA NA	7.69 NA	17.8 NA
HC04-SW/SD-FS/BN	surface bottom	0 NA	65 NA	5.3 NA	6.16 NA	17.3 NA
HM01-SW/SD-FS/BN	surface bottom	0 NA	140 NA	8.0 NA	6.9 NA	17.5 NA
HM02-SW/SD	surface bottom	24 25	36,000 38,000	11.8 11.6	7.9 7.6	17.2 17.6
HM02-FS/BN	surface bottom	21 19	29,000 27,000	7.75 7.75	NA NA	21 20
	surface bottom	2 3.75	3,810 6,000	NA NA	NA NA	19 19.5
	surface bottom	1 1.1	2,490 2,700	5.8 5.0	6.85 6.72	15.5 15.2
HM03-SW/SD	surface bottom	13.5 NA	19,000 NA	3.4 NA	6.81 NA	17.8 NA
HM03-FS/BN	surface bottom	22 NA	32,000 NA	10.8 NA	7.90 NA	17.5 NA

Sample Identification	Sample Location	Salinity (ppt)	Conductivity (micromhos/cm)	DO (mg/L)	pH (S.U.)	Temperature (deg. C)
WC02-SW/SD	surface bottom	4.5 5.5	9,000 9,000	9.0 7.0	7.48 7.48	21 20.5
	surface bottom	0 0	975 1,250	5.1 4.4	7.08 7.15	17.5 17.5
	surface bottom	0 7	850 10,500	5.5 6.1	6.98 6.85	20.5 21
	surface bottom	10 10	16,500 16,500	10 8.5	7.33 7.36	23 22.4
WC03-FS/BN	surface bottom	12 12.8	17,200 18,000	9.1 9.6	7.43 7.56	20 19

ppt = parts per thousand

S.U. = Standard Units

NA = Not Analyzed

Sample Location = Water surface or water bottom

DO = Dissolved Oxygen level

FS = Fish sample

BN = Benthic Macroinvertebrate sample

SW/SD = Surface water/sediment sample

**Positive Detection Summary  
Fish Fillet Tissue Analysis**

MARINE CORPS BASE CAMP LEJEUNE  
 BACKGROUND - HADNOT CREEK  
 POSITIVE DETECTIONS SUMMARY  
 FISH FILLET TISSUE SAMPLES

Parameter	HC1A-RD (Red Drum) (mg/kg)	HC1A-SF (Southern Flounder) (mg/kg)	HC1A-LBA (Largemouth Bass) (mg/kg)	HC1A-LBB (Largemouth Bass) (mg/kg)	HC1A-LBC (Largemouth Bass) (mg/kg)	HC1A-BCA (Blue Crab) (mg/kg)	HC1A-BCA (Blue Crab) (mg/kg)	HC1A-GA (Longnose Gar) (mg/kg)	HC1A-GB (Longnose Gar) (mg/kg)
Volatiles									
Acetone	0.13 J	0.056 J	0.077 J	0.07 J	0.037 J	0.11 J	0.099 J	0.028 J	0.016 J
Methylene Chloride	0.041	0.013 B	0.017 B	0.016 B	0.003 B	0.011 B	0.022 B	0.004 B	0.015 B
Semivolatiles									
Phenol	ND	0.46	ND	2.1	1.6	ND	ND	ND	ND
Di-n-octyl phthalate	ND	ND	0.061 J	ND	0.085	ND	ND	0.29 J	0.5 J
Bis(2-ethylhexyl)phthalate	1.1 B	0.82 B	3.6 B	3.2 B	4.8 B	ND	ND	11 J	17 J
Pesticides/PCBs									
4,4'-DDD	ND	ND	ND	ND	ND	0.0066	0.0056	ND	ND
4,4'-DDE	ND	ND	ND	ND	ND	0.0087	0.0046	0.012	0.0097
alpha-Chlordane	ND	ND	ND	ND	0.00017 P	0.0018	0.0012	ND	ND
Aroclor-1260	ND	ND	ND	ND	ND	ND	ND	ND	ND
Inorganics									
Aluminum	ND	ND	ND	36.5	ND	ND	ND	ND	ND
Arsenic	0.7 L	0.82	0.34 L	0.37 L	0.36 K	0.68	0.39	2.5	3.9 L
Barium	ND	ND	ND	ND	ND	ND	10.1	ND	ND
Cadmium	ND	ND	ND	ND	ND	0.14	0.11 J	ND	ND
Calcium	154	271	528	684	1170	4480	32200	493	520
Chromium	0.38 L	ND	0.23 L	0.68 L	0.63 L	ND	0.52 L	0.32 L	0.21 L
Copper	0.3 J	0.18 J	0.2 J	0.24 J	0.28 J	7.9	5.8	0.46 J	0.18 J
Iron	ND	ND	ND	ND	ND	ND	ND	ND	ND
Lead	ND	ND	ND	ND	ND	ND	ND	ND	ND
Magnesium	285	254	298	292	319	591	1800	286	300
Manganese	0.13	0.38	0.09 J	0.09 J	0.08 J	1.8	13.6	0.24 J	0.21 J
Mercury	0.07	0.05	0.22	0.24	0.17 K	0.08	0.02 J	0.22	0.14
Nickel	ND	ND	ND	ND	ND	ND	ND	0.45 L	ND
Potassium	3930	3700	3740	3610	4040	2170	1860	3410	3270
Sodium	1060	607	505	580	529	4060	4270	623	523
Zinc	5	5	3.9	4.4	4.6 L	25	17.9	6.5	4.6

## **Fish Distribution and Characterization**

FISH DISTRIBUTION AND CHARACTERIZATION  
BACKGROUND STATIONS - WEBB, HADNOT, AND HOLLAND MILL CREEKS

MCB CAMP LEJEUNE, NORTH CAROLINA

Common Name	Scientific Name	Length N.C. (cm)	Length Atlas (cm)	Water Type	Habitat	Spawning	Tolerance	Family	Sources
Atlantic Menhaden	<u>Brevoortia tyrannus</u>	20	46	Brackish or marine, enters freshwater	Rivers, streams	NA	Intermediate	Clupeidae	1,2,3,4
Spot	<u>Leiostomus xanthurus</u>	NA	NA	Brackish or marine, enters freshwater	NA	NA	NA	Sciaenidae	1
Stripped Mullet	<u>Mugil cephalus</u>	NA	23-35	Brackish or marine, enters freshwater	Rivers	NA	NA	Mugilidae	1,2
Pinfish	<u>Lagodon rhomboides</u>	NA	38	Marine, seldom enters freshwater	Shallow waters	NA	NA	Sparidae	1,2
Mud Catfish (Yellow Bullhead)	<u>Ictalurus natalis</u>	24	-38	Freshwater	Rivers Streams	April through May	Tolerant	Ictaluridae	1,2,3
Redbreast Sunfish	<u>Lepomis auritus</u>	18	6-15	Freshwater	Streams	April through June	NA	Centrarchidae	1,2,3
Atlantic Croaker	<u>Micropogonias undulatus</u>	NA	61	Estuaries, brackish-water or marine	NA	NA	NA	Sciaenidae	1,2
Pumpkinseed	<u>Lepomis gibbosus</u>	20	8-20	Freshwater	Streams Creeks	April through October	Moderately Tolerant	Centrarchidae	1,2,3,4
Longnose Gar	<u>Lepisosteus osseus</u>	80	-150	Freshwater; May enter brackish water	Rivers	April through May	Intermediate	Lepisosteidae	1,2,3
Summer Flounder	<u>Paralichthys dentatus</u>	NA	37	Brackish or marine, enters freshwater	Rivers	NA	NA	Bothidae	1
Flier	<u>Centrarchus macropterus</u>	12	7-19	Freshwater	Streams	April through May	NA	Centrarchidae	1,2,3
Chain Pickerel	<u>Esox niger</u>	44	38-45	Freshwater	Streams Creeks	February through March	Intermediate	Esocidae	1,2,3

FISH DISTRIBUTION AND CHARACTERIZATION  
 BACKGROUND STATIONS - WEBB, HADNOT, AND HOLLAND MILL CREEKS  
 REMEDIAL INVESTIGATION, CTO-0232  
 MCB CAMP LEJEUNE, NORTH CAROLINA

Common Name	Scientific Name	Length N.C. (cm)	Length Atlas (cm)	Water Type	Habitat	Spawning	Tolerance	Family	Sources
Redear Fish	<u>Lepomis microlophus</u>	18	14-25	Freshwater	Streams	May through August	Intermediate	Centrarchidae	1,2,3
Warmouth	<u>Lepomis gulosus</u>	16	8-26	Freshwater	Rivers Streams	May through August	Intermediate	Centrarchidae	1,2,3
White Perch	<u>Morone americana</u>	NA	to 48	Brackish water; Freshwater	Bays and estuaries; Rivers and lakes	NA	Intermediate	Percichthyidae	3,5
Bluefish	<u>Pomatomus saltatrix</u>	NA	NA	Coastal waters	Surface waters; Near shore and off shore	NA	NA	Pomatomidae	2
Bluegill	<u>Lepomis macrochirus</u>	25	18-20	Freshwater	Rivers Streams Creeks	May through October	Intermediate	Centrarchidae	1,2,3
White Catfish	<u>Ictalurus catus</u>	31	-46	Freshwater	Rivers	May through June	Intermediate	Ictaluridae	1,2,3
Largemouth Bass	<u>Micropterus salmoides</u>	48	12-70	Freshwater	Rivers Streams Creeks	May through June	Intermediate	Centrarchidae	1,2,3
Mummichog	<u>Fundulus heteroclitus</u>	7	8-10	Shallow coastal waters	Rivers Streams	April through August	NA	Cyprinodontidae	1,2,3
Redfin Pickerel	<u>Esox americanus</u>	23	25-30	Freshwater	Streams Creeks	February through March	NA	Esocidae	1,2,3
Hog Choker	<u>Trinectes maculatus</u>	5	7-12	Shallow coastal waters; Occasionally enters freshwater	Rivers Streams	March through April	NA	Soleidae	1,2,3

FISH DISTRIBUTION AND CHARACTERIZATION  
 BACKGROUND STATIONS - WEBB, HADNOT, AND HOLLAND MILL CREEKS  
 REMEDIAL INVESTIGATION, CTO-0232  
 MCB CAMP LEJEUNE, NORTH CAROLINA

Common Name	Scientific Name	Length N.C. (cm)	Length Atlas (cm)	Water Type	Habitat	Spawning	Tolerance	Family	Sources
Pirate Perch	<u>Aphredoderus sayanus</u>	9	7-14	Freshwater	Streams Creeks	January through March	Intermediate	Aphredoderidae	1,2,3
Eastern Mosquito (Mosquitofish)	<u>Gambusia affinis</u>	NA	NA	Fresh or brackish water	Ponds, lakes, ditches, backwaters, sluggish streams	NA	Intermediate	Poeciliidae	2,5

1 Menhinick, 1992.

2 Boschung, 1983.

3 USEPA, 1989d.

4 Raasch, 1991.

5 Kennish, 1986.

NA = Information not Available

**TOTAL NUMBER AND PERCENT OF AQUATIC SPECIES IDENTIFIED PER AREA  
WEBB CREEK AND HADNOT CREEK**

MCB CAMP LEJEUNE, NORTH CAROLINA

SPECIES	WEBB CREEK		Total Detected	HADNOT CREEK				Total Detected
	WC02	WC03		HC01	HC02	HC03	HC04	
<b>FISH SPECIES</b>								
Spot	4		4			12		12
Stripped Mullet	4		4			3		3
Pumpkinseed			0		3			3
Mudcat	3		3	3				3
Redbreast sunfish	1		1	2				2
Long-Nosed Gar	9	5	14					0
American flier			0	3				3
Chain pickerel			0	1				1
Redear fish			0	1				1
Atlantic croaker			0			5		5
Warmouth			0		1			1
Bluefish			0			3		3
Yellow Bullhead	3		3	2				2
Blue gill	4		4					0
White catfish	1		1					0
Largemouth bass	2		2					0
Summer flounder		1	1					0
Mummichog		3	3					0
Pinfish	25	24	49			5		5
Atlantic menhaden			0			2		2
Redfin pickerel			0				2	2
White perch			0			1		1
Hog choker			0			1		1
Pirate perch			0				8	8

**TOTAL NUMBER AND PERCENT OF AQUATIC SPECIES IDENTIFIED PER AREA  
WEBB CREEK AND HADNOT CREEK**

**MCB CAMP LEJEUNE, NORTH CAROLINA**

SPECIES	WEBB CREEK		Total Detected	HADNOT CREEK				Total Detected
	WC02	WC03		HC01	HC02	HC03	HC04	
NO. OF SPECIES	9	4	12	5	2	8	2	18
NO. OF INDIVIDUALS	53	33	86	10	4	32	10	56
OTHER AQUATIC SPECIES								
Grass shrimp		3	3					0
Crayfish			0				3	3
NUMBER OF SPECIES	0	1	1	0	0	0	1	1
NO. OF INDIVIDUALS	0	3	3	0	0	0	3	3

**TOTAL NUMBER AND PERCENT OF AQUATIC SPECIES IDENTIFIED PER AREA  
HOLLAND MILL CREEK**

**MCB CAMP LEJEUNE, NORTH CAROLINA**

SPECIES	HOLLAND MILL CREEK (CARTWHEEL BRANCH)			Total Detected
	HM01	HM02	HM03	
Spot			8	8
Stripped Mullet		11	3	14
Pumpkinseed	16	2		18
Chain pickerel	2			2
Swamp darter	6			6
Mud sunfish	1			1
Black drum		1		1
Ligar		3		3
Gizzard Shad		2		2
Spotted sunfish		2		2
Blue gill	2	1		3
Atlantic menhaden			199	199
Largemouth bass		1		1
Hog choker			2	2
Summer flounder		1	17	18
Mummichog		6		6
Pinfish		7	4	11
Goby, freshwater	1	1		2
<b>NUMBER OF SPECIES</b>	<b>6</b>	<b>12</b>	<b>6</b>	<b>18</b>
<b>NO. OF INDIVIDUALS</b>	<b>28</b>	<b>38</b>	<b>233</b>	<b>299</b>

**TOTAL NUMBER AND PERCENT OF AQUATIC SPECIES IDENTIFIED PER AREA  
HOLLAND MILL CREEK**

**MCB CAMP LEJEUNE, NORTH CAROLINA**

SPECIES	HOLLAND MILL CREEK (CARTWHEEL BRANCH)			Total Detected
	HM01	HM02	HM03	
<b>OTHER AQUATIC SPECIES</b>				
Unknown	1			1
Grass shrimp		13		13
Crayfish	3			3
<b>NUMBER OF SPECIES</b>	2	1	0	3
<b>NO. OF INDIVIDUALS</b>	4	13	0	17

## HADNOT CREEK - BACKGROUND STATIONS

SPECIES	COC SAMPLE NO.	HC01			HC02			HC03			HC04		
		Fish Length (cm)	Mass Weight (g)	Average Weight (g)	Fish Length (cm)	Mass Weight (g)	Average Weight (g)	Fish Length (cm)	Mass Weight (g)	Average Weight (g)	Fish Length (cm)	Mass Weight (g)	Average Weight (g)
Stripper Mullet	HC03							15.25	45	45			
								12.5	20	20			
								12.5	20	20			
	COUNT							3		3			
	AVERAGE							13.416666667		28.333333333			
	MAXIMUM							15.25		45			
	MINIMUM							12.5		20			
Atlantic Menhaden	HC03							+1 collected, no length or weight					
								5 <5		2.5			
	COUNT							2		2			
	AVERAGE							5		2.5			
	MAXIMUM							5		2.5			
	MINIMUM							5		2.5			
Blue Fish	HC03							7	7	7			
								11	17	17			
								8	8	8			
	COUNT							3		3			
	AVERAGE							8.6666666667		10.666666667			
	MAXIMUM							11		17			
	MINIMUM							7		7			
Spot	HC03							12.5	22	22			
								5.5 <5.0		2.5			
								5.75 <5.0		2.5			
								5 <5.0		2.5			
								3.5 <5.0		2.5			
								5.5 <5.0		2.5			
								14	40	40			
								13.5	35	35			
								12	35	35			
								14	35	35			
								5.5 <5.0		2.5			
								11.5	20	20			
	COUNT							12		12			
	AVERAGE							9.0208333333		16.833333333			
	MAXIMUM							14		40			
	MINIMUM							3.5		2.5			

#### HADNOT CREEK - BACKGROUND STATIONS

SPECIES	CO SAMPLE NO.	HC01			HC02			HC03			HC04		
		Fish Length (cm)	Mass Weight (g)	Average Weight (g)	Fish Length (cm)	Mass Weight (g)	Average Weight (g)	Fish Length (cm)	Mass Weight (g)	Average Weight (g)	Fish Length (cm)	Mass Weight (g)	Average Weight (g)
White Perch	HC03							18.5	105	105			
	COUNT							1		1			
	AVERAGE							18.5		105			
	MAXIMUM							18.5		105			
	MINIMUM							18.5		105			
Hogchoker	HC03							5.5	5	5			
	COUNT							1		1			
	AVERAGE							5.5		5			
	MAXIMUM							5.5		5			
	MINIMUM							5.5		5			
Pinfish	HC03							13	35	35			
								10.8	25	25			
								11	22	22			
								10.5	25	25			
								13	37	37			
	COUNT							5		5			
	AVERAGE							11.66		26.6			
	MAXIMUM							13		37			
	MINIMUM							10.5		22			
Atlantic Croaker	HC03							11.5	20	20			
								10.5	16	16			
								9	10	10			
								10.5	14	14			
								7.5 <5		2.5			
	COUNT							5		5			
	AVERAGE							9.8		12.5			
	MAXIMUM							11.5		20			
	MINIMUM							7.5		2.5			
Redbreast Sunfish	HC01	23.5 20	265 175	265 175									
	COUNT							2		2			
	AVERAGE							21.75	220				
	MAXIMUM							23.5	285				
	MINIMUM							20	175				
American Flyer	HC01	16.5 9.5	85 15	65 15									
	+1 collected, no length or weight												
	COUNT							3		3			
	AVERAGE							13	40				
	MAXIMUM							16.5	65				
	MINIMUM							9.5	15				

#### HADNOT CREEK - BACKGROUND STATIONS

#### HOLLAND MILL CREEK - BACKGROUND STATIONS

SPECIES	COO SAMPLE NO.	HM01			HM02			HM03		
		Fish Length (cm)	Mass Weight (g)	Average Weight (g)	Fish Length (cm)	Mass Weight (g)	Average Weight (g)	Fish Length (cm)	Mass Weight (g)	Average Weight (g)
Stripper Mullet	HM02				38.5	640	640			
					39.5	600	600			
					34.5	400	400			
					34.5	400	400			
					33.5	360	360			
					34	340	340			
					37	460	460			
					35	520	520			
					33.5	410	410			
					32	320	320			
					31	370	370			
	HM03							14.5	40	40
								6.5 <5		2.5
								+1 collected, no length or weight		
	COUNT			11		11		3		3
	AVERAGE			34.81818182		438.1818182		10.5		21.25
	MAXIMUM			39.5		640		14.5		40
	MINIMUM			31		320		6.5		2.5
Atlantic Menhaden	HM03							6	24	4
								6		4
								5.7		4
								5.4		4
								5.5		4
								5.6		4
								5.7	22	2.2
								5.5		2.2
								5		2.2
								5.5		2.2
								5.5		2.2
								5.2		2.2
								5.5		2.2
								5.5		2.2
								5.5		2.2
								5.6		2.2
								6.2		2.2
								6	25	2.5
								5.5		2.5
								5		2.5
								5.5		2.5
								5.5		2.5
								6		2.5
								5		2.5
								5.5		2.5
								5.5	20	2
								5.7		2
								5		2
								6		2
								5.5		2
								6		2
								5.5		2
								6		2
								5.5	27	1.8
								5.6		1.6
								5.5		1.8
								5.7		1.8
								6		1.8
								6		1.8
								6.5		1.8
								5.5		1.8
								6.5		1.8
								5.5		1.8
								6		1.8
								5.5		1.8
								6		1.8
								5.5	20	2
								4.5		2
								5		2
								5.5		2
								5.5		2
								5.5		2
								6		2
								5.5		2
								6		2
								6		2
								6		2

## HOLLAND MILL CREEK - BACKGROUND STATIONS

SPECIES	COC SAMPLE NO.	HM01			HM02			HM03		
		Fish Length (cm)	Mass Weight (g)	Average Weight (g)	Fish Length (cm)	Mass Weight (g)	Average Weight (g)	Fish Length (cm)	Mass Weight (g)	Average Weight (g)
Summer Flounder	HM02				29.5	250	250			
	HM03							33	400	400
							43	850	850	
							20.5	90	90	
							24	120	120	
					<u>+13 collected, no length or weight</u>					
	COUNT			1		1	17			4
	AVERAGE				29.5		250	30.125		365
	MAXIMUM				29.5		250	43		850
	MINIMUM				29.5		250	20.5		90
Black Drum	HM02				28	350	350			
	COUNT			1		1				
	AVERAGE				28		350			
	MAXIMUM				28		350			
	MINIMUM				28		350			
Spotted Sunfish	HM02				15.5	65	65			
					17	110	110			
	COUNT			2		2				
	AVERAGE				16.25		87.5			
	MAXIMUM				17		110			
	MINIMUM				15.5		65			
Largemouth Bass	HM02				34	540	540			
	COUNT			1		1				
	AVERAGE				34		540			
	MAXIMUM				34		540			
	MINIMUM				34		540			
Hogchoker	HM03							<u>+1 collected, no length or weight</u>		
	COUNT							6	10	10
	AVERAGE							2		1
	MAXIMUM							6		10
	MINIMUM							6		10
Spot	HM03							<u>+1 collected, no length or weight</u>		
	COUNT							5 <5	2.5	
	AVERAGE							12	25	25
	MAXIMUM							5.8	20	4
	MINIMUM							6		4
								6.2		4
								6.4		4
								6.4		4
					<u>+1 collected, no length or weight</u>					
	COUNT							8	7	
	AVERAGE							6.82857143	6.78571429	
	MAXIMUM							12	25	
	MINIMUM							5	2.5	
Blue Gill	HM02					17		105		
	HM01	10.5	10	10						
	COUNT	2			1	1		1		
	AVERAGE	10.5			10	17		105		
	MAXIMUM	10.5			10	17		105		
	MINIMUM	10.5			10	17		105		

## HOLLAND MILL CREEK - BACKGROUND STATIONS

SPECIES	COO SAMPLE NO.	HM01 Fish Length (cm)	Mass Weight (g)	Average Weight (g)	HM02 Fish Length (cm)	Mass Weight (g)	Average Weight (g)	HM03 Fish Length (cm)	Mass Weight (g)	Average Weight (g)
Pumpkinseed	HM02				15 11.5	50 30	50 30			
	HM01	7.5 6.5 7.5 7.5 6 6 4.5 8.5 8 5.5 8 8.5 6.5 8.5 11 7.5	45 4.5 4.5 4.5 4.5 4.5 4.5 4.5 4.5 4.5 8.3 8.3 8.3 8.3 8.3 8.3	4.5 4.5 4.5 4.5 4.5 4.5 4.5 4.5 4.5 4.5 8.3 8.3 8.3 8.3 8.3 8.3						
	COUNT	16		16	2		2			
	AVERAGE	7.34375		5.925	13.25		40			
	MAXIMUM	11		5.3	15		50			
	MINIMUM	4.5		4.5	11.5		30			
Long-nose Gar	HM02				73 83 72.5	1250 2000 1640	1250 2000 1640			
	COUNT			3		3				
	AVERAGE			76.16666667		1630				
	MAXIMUM			83		2000				
	MINIMUM			72.5		1250				
Pinfish	HM02 HM03			17.5	80	80		5 <5		2.5
	COUNT			7		1	4			1
	AVERAGE			17.5		80	5			2.5
	MAXIMUM			17.5		80	5			2.5
	MINIMUM			17.5		80	5			2.5
Gizzard Shad	HM02			33 34	480 460	480 460				
	COUNT			2		2				
	AVERAGE			33.5		470				
	MAXIMUM			34		480				
	MINIMUM			33		460				
Chain Pickerel	HM01	13 13.5	10	5						
	COUNT	2		2						
	AVERAGE	13.25		5						
	MAXIMUM	13.5		5						
	MINIMUM	13		5						
Unknown Fish	HM01	7.5 <5		2.5						
	COUNT	1		1						
	AVERAGE	7.5		2.5						
	MAXIMUM	7.5		2.5						
	MINIMUM	7.5		2.5						
Swamp Darter	HM01	6 6 6 6 6 6	18	3						
	COUNT	6		6						
	AVERAGE	6		3						
	MAXIMUM	6		3						
	MINIMUM	6		3						

#### HOLLAND MILL CREEK - BACKGROUND STATIONS

## WEBB CREEK - BACKGROUND STATIONS

SPECIES	COC SAMPLE NO.	WC02			WC03		
		Fish Length (cm)	Mass Weight (g)	Average Weight (g)	Fish Length (cm)	Mass Weight (g)	Average Weight (g)
Stripper Mullet	WC02	39.5 35.5 41.5 37	500 380 700 600	500 380 700 600			
		COUNT	4	4			
		AVERAGE	38.375	545			
		MAXIMUM	41.5	700			
		MINIMUM	35.5	380			
Summer Flounder	WC03				21	60	60
		COUNT			1		1
		AVERAGE			21		60
		MAXIMUM			21		60
		MINIMUM			21		60
Largemouth Bass	WC02	34 34	525 600	525 600			
		COUNT	2	2			
		AVERAGE	34	562.5			
		MAXIMUM	34	600			
		MINIMUM	34	525			
Redbreast Sunfish	WC02	16	60	60			
		COUNT	1	1			
		AVERAGE	16	60			
		MAXIMUM	16	60			
		MINIMUM	16	60			
White Catfish	WC02	37	750	750			
		COUNT	1	1			
		AVERAGE	37	750			
		MAXIMUM	37	750			
		MINIMUM	37	750			
Spot	WC02	14.5 13 13 <10	10 10 5	10 10 5			
		+1 collected, no length or weight					
		COUNT	4	4			
		AVERAGE	13.5	8.33333333			
		MAXIMUM	14.5	10			
		MINIMUM	13	5			
Blue Gill	WC02	23 23.5 21.5 16.75	300 300 250 85	300 300 250 85			
		COUNT	4	4			
		AVERAGE	21.1875	233.75			
		MAXIMUM	23.5	300			
		MINIMUM	16.75	85			

## WEBB CREEK - BACKGROUND STATIONS

SPECIES	COC SAMPLE NO.	WC02			WC03		
		Fish Length (cm)	Mass Weight	Average Weight (g)	Fish Length (cm)	Mass Weight	Average Weight (g)
Long-nose Gar	WC02	68	1100	1100			
		71.5	1220	1220			
		73.5	1350	1350			
		72.5	1220	1220			
		66.5	1120	1120			
		72.5	1260	1260			
		71.5	1340	1340			
		69.5	1240	1240			
		75	1420	1420			
	WC03				87	1900	1900
					83	1850	1850
					97	2850	2850
					71.5	1000	1000
					73	1580	1580
	COUNT	9		9	5		5
	AVERAGE	71.16667		1252.222	82.3		1836
	MAXIMUM	75		1420	97		2850
	MINIMUM	66.5		1100	71.5		1000
Pinfish	WC02	10.5	NA				
		+24 collected, no length or weight		24 collected, no length or weight			
		COUNT	25		24		
		AVERAGE	10.5				
		MAXIMUM	10.5				
		MINIMUM	10.5				
	WC02	38.5	900	900			
Yellow Bullhead Catfish		32.5	620	620			
		36.5	640	640			
	COUNT	3		3			
	AVERAGE	35.83333		720			
	MAXIMUM	38.5		900			
	MINIMUM	32.5		620			
Mudcat		3 fish collected at WC02, no length or weight					
Mummichog		3 fish collected at WC03, no length or weight					
Grass shrimp		3 collected at WC03, no length or weight					

**Benthic Macroinvertebrate  
Characterization and Statistics**

MARINE CORPS BASE CAMP LEJEUNE  
 BACKGROUND - WEBB CREEK  
 BENTHIC MACROINVERTEBRATES

	WC02-BN			WC03-BN		
	01	02	03	01	02	03
NEMERTEA						
Anopla						
Heteronemertea						
Lineidae						
Micrura leidyl				1	2	2
ANNELIDA						
Polychaeta						
Capitellida						
Capitellidae						
Heteromestus filiformis	2					
Phyllodocida						
Nereidae						
Nereis succinea				1		
Spionida						
Spionidae						
Scolecolepides viridis					1	
Terebellida						
Ampharetidae						
Hypaniola grayi	4	10				
ARTHROPODA						
Crustacea						
Amphipoda						
Gammaeridae						
Gammarus tigrinus	10			1	1	
Insecta						
Diptera						
Chironomidae						
Chironomus decorus gr.	8	24	13	38	17	6
Procladius sp.	1	3		2		1
Tanytarsus sp.		2	1			
MOLLUSCA						
Bivalvia						
Veneroida						
Corbiculidae						
Polymesoda caroliniana				1		
Tellinidae						
Macoma tenta				1		
Total Taxa	4	4	4	4	5	4
Total Specimens	21	33	25	42	22	10
Replicate Specimens Average		26.33			24.67	
Standard Deviation	4.42531	10.5317	6.18466	18.3394	7.05691	2.38048
Brillouin's Diversity		0.518			0.279	
SPECIES DENSITY (#/M <sup>2</sup> )	134	210	159	268	140	64
SPECIES DIVERSITY (Shannon-Wiener)	0.473	0.380	0.419	0.180	0.304	0.473

MARINE CORPS BASE CAMP LEJEUNE  
BACKGROUND - HADNOT CREEK  
BENTHIC MACROINVERTEBRATES

	HC01-BN			HC02-BN			HC03 BN			HC04-BN		
	01	02	03	01	02	03	01	02	03	01	02	03
NEMERTEA							6	5	3			
Anopla												
Heteronemertea												
Lineidae												
Micruro leidyi												
ANNELIDA												
Oligochaeta												
Lumbricida												
Lumbriculina												
Eclipidrilus sp.				1								
Tubificida												
Tubificidae												
Isotomidae freyi	77	42	36							21	21	6
Limnodrilus hoffmeisteri										1		
Spirosperma carolinense		3								1		3
Polychaeta												
Capitellidae												
Capitellidae												
Heteromyctes filiformis										14	9	
Phyllodocidae												
Nereidae										6	18	
Nereis succinea												
Phyllodocidae												
Eteone heteropoda												1
Terebellidae												
Ampharetidae												
Hyperia grayi (ampharetid worm)										18	6	46
ARTHROPODA												
Crustacea												
Amphipoda												
Ceropagidae												
Ceropagium lacustris												
Gammaridae												
Crangonyx pseudogracilis										1	1	
Gammarus tigrinus												
Tanidae												
Tanaididae												
Leptocheirus rapax												
Insects												
Coleoptera												
Dytiscidae												
Hydroorus sp.										1		
Elmidae												
Dubaephilus sp.												
Diptera												
Ceratopogonidae												
Palpomyia/sphaeromias sp. (biting midges)	5	7	4									
Chironomidae												
Ablepharomyia annulata	2	7	1									
Ablepharomyia namphe gr.	4	7	9									
Clinotanypus pinguis		2	3									
Cryptochironomus fulvus gr.												
Epaichidius sp.												
Glyptotendipes sp.												
Närotheuma sp.												
Paraculiciferornata nigrochakera	1	5	2									
Polypedilum illinoense	3	1										
Procladius sp. (midges)												
Tanytarsus sp.	2	9	2									
Tribolium lucidum	4	8	6									
Tipulidae												
Pseudolimnophila sp.												
Ephemeroptera												
Ephemeridae												
Hexagenia bilineata	3	3	1									
Megaloptera												
Sialidae												
Sialis sp.												
Odonata												
Coenagrionidae												
Argia sp.												
Libellulidae												
Pachydiplax longipennis												
Trichoptera												
Polycentropodidae												
Phylacanthopus sp.	1	5	7									
MOLLUSCA												
Bivalvia												
Mytilidae												
Mytilus demissus												
Veneridae												
Sphaeriidae												
Pisidium casertanum		2	1									
Tellinidae												
Macoma tenta												
Total Taxa	10	17	15	1	2	4	4	3	6	4	11	8
Total Specimens	102	106	76	18	7	34	30	31	183	44	69	52
Replicate Specimens Average	95.33333	98.614633	8.961824	NA	3.53534	21.79449	4.356898	8.062904	39.67241	9.321905	7.129687	6.047432
Standard Deviation	23.50792	8.614633	8.961824									
Brillouin's Diversity	0.755			0.072			0.673			0.757		
SPECIES DENSITY (#/M^2)	650	676	497	115	45	344	191	198	1166	260	440	331
SPECIES DIVERSITY (Shannon-Wiener)	0.463	0.956	0.831	0.000	0.178	0.230	0.534	0.364	0.449	0.458	0.603	0.763

MARINE CORPS BASE CAMP LEJEUNE  
BACKGROUND - HADNOT CREEK  
BENTHIC MACROINVERTEBRATES

	HM01-BN			HM02-BN			HM03-BN		
	01	02	03	01	02	03	01	02	03
NEMERTEA									
Anopla									
Heteronemertea									
Lineidae									
Micrura leidyi									
ANNELIDA									
Oligochaeta									
Tubificida									
Tubificidae									
Limnodrilus hoffmeisteri	3	1	3						
Polychaeta									
Aricida									
Orbiniidae									
Scoloplos fragilis									
Capitellida									
Capitellidae									
Heteromastus filiformis									
Phyllodocida									
Nereidae									
Nereis succinea									
Spionida									
Spionidae									
Streblospio benedicti									
Terebellida									
Ampharetidae									
Hypaniota grayi (ampharetid worm)									
ARTHROPODA									
Crustacea									
Decapoda									
Palaemonidae									
Palaemonetes pugic									
Insecta									
Coleoptera									
Dytiscidae									
Hydroporus sp.	1								
Elmidae									
Dubiraphis sp.									
Diptera									
Chaoboridae									
Chaoborus sp.									
Chironomidae									
Ablabesmyia mallochi	1								
Chironomus decorus gr.	2	2	2						
Dicrotendipes nervosus	5								
Larsia sp.									
Polypedilum illinoense	12								
Polypedilum scalaenum	18								
Tanytarsus sp.	11								
Tribelos lucundum	50	159	31						
Megaloptera									
Sialidae									
Sialis sp.	1								
MOLLUSCA									
Bivalvia									
Venerida									
Mactridae									
Mulinia lateralis									
Tellinidae									
Macoma tenta									
Total Taxa	10	3	10	3	2	4	7	4	4
Total Specimens	104	162	79	130	189	85	29	48	20
Replicate Specimens Average		115			134.667			32.3333	
Standard Deviation	15.0864	90.934	9.06091	66.4254	120.915	36.5639	5.75698	11.1056	4.08248
Brillouin's Diversity		0.5			0.122			0.497	
SPECIES DENSITY (#/M^2)	663	1033	504	829	1205	542	185	306	127
SPECIES DIVERSITY (Shannon-Wiener)	0.695	0.045	0.793	0.138	0.083	0.186	0.593	0.436	0.480

**SUMMARY STATISTICS OF BENTHIC MACROINVERTEBRATE SPECIES AT  
HADNOT CREEK, HOLLAND MILL CREEK, AND WEBB CREEK  
MCB CAMP LEJEUNE, NORTH CAROLINA**

Station	Number of Species	Number of Organisms	Species Density (#/m <sup>2</sup> )	Species Diversity (Shannon-Weiner)	Species Diversity (Brillouin's)	Macroinvertebrate Biotic Index
WC02	7	79	504	0.570	0.518	9.4
WC03	7	74	472	0.323	0.279	9.6
HC01	20	286	1,823	0.802	0.755	7.8
HC02	4	79	504	0.196	0.072	7.6
HC03	8	244	1,555	0.683	0.675	NA
HC04	13	165	1,052	0.807	0.757	7.6
HM01	13	345	2,199	0.525	0.500	6.9
HM02	4	404	2,575	0.128	0.122	9.6
HM03	7	97	618	0.538	0.497	9.6

WC = Webb Creek Stations

HC = Hadnot Creek Stations

HM = Holland Mill Creek Stations

BN = Benthic Macroinvertebrate Sample

NA = Not Applicable

Species Density (#m<sup>2</sup>) is based on a sample area of 0.0523 m<sup>2</sup>.

**SYSTEMATIC LIST OF BENTHIC MACROINVERTEBRATE SPECIES  
AT BACKGROUND STATIONS  
(WEBB, HADNOT, AND HOLLAND MILL CREEKS)  
MCB CAMP LEJEUNE, NORTH CAROLINA**

Species	USEPA <sup>(1)</sup> Metals
<b>NERMERTEA</b>	Phylum
Anopla	Class
Heteronemertea	Order
Lineidae	Family
<i>Micrura leidyl</i>	Genus Species
<b>ANNELIDA</b>	Phylum
Oligochaeta	Class
Lumbriculida	Order
Lumbriculiae	Family
<i>Eclipidrilus sp.</i>	Genus Species
Tubificida	Order
Tubificidae	Family
<i>Isochaetides freyi</i>	Genus Species
<i>Limnodrilus hoffmeisteri</i>	Genus Species
<i>Spirosperma carolinensis</i>	Genus Species
Polychaeta	Class
Ariciida	Order
Orbiniidae	Family
<i>Scoloplos fragilis</i>	Genus Species
Capitellida	Order
Capitellidae	Family
<i>Heteromestus filiformis</i>	Genus Species
Phyllodocida	Order
Nereidae	Family
<i>Nereis succinea</i>	Genus Species
Phyliodocidae	Family
<i>Eteone heteropoda</i>	Genus Species
Spionida	Order
Spionidae	Family
<i>Scolecolepides virdis</i>	Genus Species
<i>Streblospio benedicti</i>	Genus Species
Terebellida	Order

**SYSTEMATIC LIST OF BENTHIC MACROINVERTEBRATE SPECIES  
AT BACKGROUND STATIONS  
(WEBB, HADNOT, AND HOLLAND MILL CREEKS)  
MCB CAMP LEJEUNE, NORTH CAROLINA**

Species	USEPA <sup>(1)</sup> Metals
Ampharetidae	Family
<i>Hypaniola grayi</i>	Genus Species
<b>ARTHROPODA</b>	<b>Phylum</b>
Crustacea	Class
Amphipoda	Order
Corophiidae	Family
<i>Corophium lacuatre</i>	Genus Species
Gammaridae	Family
<i>Crangonyx pseudogracilllus</i>	Genus Species
<i>Gammarus tigrinus</i>	Genus Species
Tanaidacea	Order
Tanaidae	Family
<i>Leptocheilia rapox</i>	Genus Species
Decapoda	Order
Palaemonidae	Family
<i>Palaemonetes pugio</i>	Genus Species
Insecta	Class
Coleoptera	Order
Dytiscidae	Family
<i>Hydroporus sp.</i>	Genus Species
Elmidae	Family
<i>Dubiraphia sp.</i>	Genus Species
Diptera	Order
Ceratopogonidae	Family
<i>Palpomyia/sphaeromias sp.</i>	Genus Species
Chaoboridae	Family
<i>Chaoborus sp.</i>	Genus Species
Chironomidae	Family
<i>Ablabesmyia annulata</i>	Genus Species
<i>Ablabesmyia mallochi</i>	Genus Species
<i>Ablabesmyia ramphe gr.</i>	Genus Species
<i>Clinotanypus pinguis</i>	Genus Species
<i>Chironomus decorus gr.</i>	Genus Species

**SYSTEMATIC LIST OF BENTHIC MACROINVERTEBRATE SPECIES  
AT BACKGROUND STATIONS  
(WEBB, HADNOT, AND HOLLAND MILL CREEKS)  
MCB CAMP LEJEUNE, NORTH CAROLINA**

Species	USEPA <sup>(1)</sup> Metals
<i>Cryptochironomus fulvus gr</i>	Genus Species
<i>Dicrotendipes nervosus</i>	Genus Species
<i>Epoicladius sp.</i>	Genus Species
<i>Glyptotendipes sp.</i>	Genus Species
<i>Larsia sp.</i>	Genus Species
<i>Nilohauma sp.</i>	Genus Species
<i>Paraiauterborniella nigrohaite</i>	Genus Species
<i>Polypedilum illinoense</i>	Genus Species
<i>Polypedilum scalaenum</i>	Genus Species
<i>Procladius sp.</i>	Genus Species
<i>Tanytarsus sp.</i>	Genus Species
<i>Tribelos jucundum</i>	Genus Species
<i>Tribelos lucundum</i>	Genus Species
Tipulidae	Family
<i>Psuedolimnophila sp.</i>	Genus Species
Ephemeroptera	Order
Ephemeridae	Family
<i>Hexagenia billineata</i>	Genus Species
Megaloptera	Order
Sialidae	Family
<i>Sialis sp.</i>	Genus Species
Odonata	Order
Coenagrionidae	Family
<i>Argia sp.</i>	Genus Species
Libelluiidae	Family
<i>Pechydiplax longipennis</i>	Genus Species
Trichoptera	Order
Polycentropodidae	Family
<i>Phylacentropus sp.</i>	Genus Species
<b>MOLLUSCA</b>	Phylum
Bivalvia	Class
Mytiloida	Order
Mytilidae	Family

SYSTEMATIC LIST OF BENTHIC MACROINVERTEBRATE SPECIES  
 (AT BACKGROUND STATIONS  
 (WEBB, HADNOT, AND HOLLAND MILL CREEKS)  
 MCB CAMP LEJEUNE, NORTH CAROLINA

Species	USEPA <sup>(1)</sup> Metals
<i>Geukensia demissa</i>	Genus Species
Veneroida	Order
Corbiculidae	Family
<i>Polymesoda caroliniana</i>	Genus Species
Mactridae	Family
<i>Mullinia lateralis</i>	Genus Species
Sphaeriidae	Family
<i>Pisidium casertanum</i>	Genus Species
Tellinidae	Family
<i>Macoma tenta</i>	Genus Species

**USEPA SENSITIVITY TO METALS AND TOLERANCE TO ORGANIC WASTE AND BIOTIC INDEX  
FOR BENTHIC MACROINVERTEBRATE SPECIES AT BACKGROUND STATIONS  
(WEBB, HADNOT, AND HOLLAND MILL CREEKS)  
MCB CAMP LEJEUNE, NORTH CAROLINA**

Species	USEPA <sup>(1)</sup> Metals	Organics	NCDEHNR <sup>(2)</sup> Biotic Index
<b>NERMERTEA</b>			
Anopla			
Heteronemertea			
Lineidae			
<i>Micrura leidyl</i>	NA	NA	NA
<b>ANNELIDA</b>			
Oligochaeta			
Lumbriculida			
Lumbriculiae			
<i>Eclipidrillus sp.</i>	NA	NA	NA
Tubificida			
Tubificidae			
<i>Isochaetides freyi</i>	NA	NA	8.6
<i>Limnodrilus hoffmeisteri</i>	NA	5	9.4
<i>Spirosperma carolinensis</i>	NA	3	NA
Polychaeta			
Aricida			
Orbiniidae			
<i>Scoloplos fragilis</i>	NA	NA	NA
Capitellida			
Capitellidae			
<i>Heteromestus filiformis</i>	NA	NA	NA
Phyllodocida			
Nereidae			
<i>Nereis succinea</i>	NA	NA	NA
Phyliodocidae			
<i>Eteone heteropoda</i>	NA	NA	NA
Spionida			
Spionidae			
<i>Scolecolepides virdis</i>	NA	NA	NA
<i>Streblospio benedicti</i>	NA	NA	NA
Terebellida			

USEPA SENSITIVITY TO METALS AND TOLERANCE TO ORGANIC WASTE AND BIOTIC INDEX  
 FOR BENTHIC MACROINVERTEBRATE SPECIES AT BACKGROUND STATIONS  
 (WEBB, HADNOT, AND HOLLAND MILL CREEKS)  
 MCB CAMP LEJEUNE, NORTH CAROLINA

Species	USEPA <sup>(1)</sup> Metals	Organics	NCDEHNR <sup>(2)</sup> Biotic Index
Ampharetidae			
<i>Hypaniola grayi</i>	NA	NA	NA
ARTHROPODA			
Crustacea			
Amphipoda			
Corophiidae			
<i>Corophium lacuatre</i>	NA	NA	NA
Gammaridae			
<i>Crangonyx pseudogracilllus</i>	NA	NA	7.9
<i>Gammarus tigrinus</i>	NA	2	NA
Tanaidacea			
Tanaidae			
<i>Leptochelia rapox</i>	NA	NA	NA
Decapoda			
Palaemonidae			
<i>Palaemonetes pugio</i>	NA	NA	NA
Insecta			
Coleoptera			
Dytiscidae			
<i>Hydroporus sp.</i>	NA	NA	8.6
Elmidae			
<i>Dubiraphia sp.</i>	NA	NA	5.9
Diptera			
Ceratopogonidae			
<i>Palpomyia/sphaeromias sp.</i>	NA	NA	7.0
Chaoboridae			
<i>Chaoborus sp.</i>	NA	NA	8.5
Chironomidae			
<i>Ablabesmyia annulata</i>	NA	1	3.5
<i>Ablabesmyia mallochi</i>	S	2	7.2
<i>Ablabesmyia ramphe gr.</i>	NA	2	NA
<i>Clinotanypus pinguis</i>	S	3	8.7

**USEPA SENSITIVITY TO METALS AND TOLERANCE TO ORGANIC WASTE AND BIOTIC INDEXES  
FOR BENTHIC MACROINVERTEBRATE SPECIES AT BACKGROUND STATIONS  
(WEBB, HADNOT, AND HOLLAND MILL CREEKS)  
MCB CAMP LEJEUNE, NORTH CAROLINA**

Species	USEPA <sup>(1)</sup> Metals	Organics	NCDEHNR <sup>(2)</sup> Biotic Index
<i>Chironomus decorus</i> gr.	NA	NA	9.6
<i>Cryptochironomus fulvus</i> gr	NA	3	6.4
<i>Dicrotendipes nervosus</i>	S	2	9.7
<i>Epoicladius</i> sp.	NA	NA	0.0
<i>Glyptotendipes</i> sp.	NA	NA	9.4
<i>Larsia</i> sp.	NA	2	9.3
<i>Nilothauma</i> sp.	NA	NA	5.0
<i>Paraiauterborniella nigrohaite</i>	NA	NA	NA
<i>Polypedilum illinoense</i>	NA	3	9.0
<i>Polypedilum scalaenum</i>	NA	2	8.4
<i>Procladius</i> sp.	NA	NA	9.1
<i>Tanytarsus</i> sp.	NA	NA	6.7
<i>Tribelos jucundum</i>	S	1	6.3
<i>Tribelos lucundum</i>	NA	NA	6.3
Tipulidae			
<i>Pseudolimnophila</i> sp.	NA	NA	7.2
Ephemeroptera			
Ephemeridae			
<i>Hexagenia bilineata</i>	NA	2	NA
Megaloptera			
Sialidae			
<i>Sialis</i> sp.	T	4	7.2
Odonata			
Coenagrionidae			
<i>Argia</i> sp.	NA	NA	8.2
Libelluiidae			
<i>Pechydiplax longipennis</i>	NA	NA	NA
Trichoptera			
Polycentropodidae			
<i>Phylacentropus</i> sp.	NA	NA	6.2
MOLLUSCA			
Bivalvia			

**USEPA SENSITIVITY TO METALS AND TOLERANCE TO ORGANIC WASTE AND BIOTIC INDES  
FOR BENTHIC MACROINVERTEBRATE SPECIES AT BACKGROUND STATIONS  
(WEBB, HADNOT, AND HOLLAND MILL CREEKS)  
MCB CAMP LEJEUNE, NORTH CAROLINA**

Species	USEPA <sup>(1)</sup> Metals	Organics	NCDEHNR <sup>(2)</sup> Biotic Index
Mytiloida			
Mytilidae			
<i>Geukensia demissa</i>	NA	NA	NA
Veneroida			
Corbiculidae			
<i>Polymesoda caroliniana</i>	NA	NA	NA
Mactridae			
<i>Mullinia lateralis</i>	NA	NA	NA
Sphaeriidae			
<i>Pisidium casertanum</i>	NA	4	6.5
Tellinidae			
<i>Macoma tenta</i>	NA	NA	NA

<sup>(1)</sup> Macroinvertebrate Field and Laboratory Methods for Evaluating the Biological Integrity of Surface Waters

<sup>(2)</sup> Lenat, 1993

NA = Not Available

S = Sensitive to heavy metals


T = Tolerant to heavy metals

Organics Ranking = 0 to 5 with 0 being the least tolerant

**Sampling Station  
Location Maps**

## **REFERENCE**

Baker, 1994. Baker Environmental Inc., 1994. "Supplemental Aquatic Survey for Wallace Creek and Bearhead Creek". Prepared for the Department of the Navy, Naval Facilities Engineering Command, Atlantic Division, Norfork, Virginia.




FISH AND BENTHIC MACROINVERTEBRATE  
SAMPLING LOCATION IN HADNOT CREEK

MARINE CORPS BASE, CAMP LEJEUNE  
NORTH CAROLINA


SOURCE: N.C. DIVISION OF MARINE  
FISHERIES, REPORT AFC-9, NOV. 1975.

01580TB2Z



FISH AND BENTHIC MACROINVERTEBRATE  
SAMPLING LOCATION IN HOLLAND MILL CREEK

MARINE CORPS BASE, CAMP LEJEUNE  
NORTH CAROLINA



FISH AND BENTHIC MACROINVERTEBRATE  
SAMPLING LOCATION IN WEBB CREEK

MARINE CORPS BASE, CAMP LEJEUNE  
NORTH CAROLINA

**APPENDIX O**  
**SHOWER MODEL**

---

## SHOWER INHALATION MODEL

### INHALATION OF CONTAMINANTS VOLATILIZED FROM SHOWER WATER

In the model developed by Foster and Chrostowski (1986), inhalation exposures to volatile organic chemicals (VOCs) while showering are modeled by estimating the rate of chemical releases into the air (generation rate), the buildup of VOCs in the shower room air while the shower is on, and the decay of VOCs in the shower room air after the shower is turned off, and the quantity of airborne VOCs inhaled while the shower is both on and off.

Estimation of the rate of VOC release into the air is based upon Liss and Slater's (1974) adaptation of the two-film gas-liquid mass transfer theory. The two-film boundary theory provides the basis for estimating the overall mass transfer coefficient ( $K_L$ ) for each VOC of interest, according to the following equation:

$$K_L = \left( \frac{1}{k_1} + \frac{RT}{Hk_g} \right)^{-1} \quad (1)$$

where,

$K_L$  = overall mass transfer coefficient (centimeter per hour [cm/hr]),

H = Henry's law constant (atm-m<sup>3</sup>/mol-K),

RT =  $2.4 \times 10^{-2}$  atm-m<sup>3</sup>/mole (gas constant of  $8.2 \times 10^{-5}$  atm-m<sup>3</sup>/mole-K times absolute temperature of 293 K),

$k_g$  = gas-film mass transfer coefficient (cm/hr), and

$k_1$  = liquid-film mass transfer coefficient (cm/hr).

Equation 1 describes the mass transfer rate of a compound at an air-water interface where diffusion may be limited by both liquid- and gas-phase resistances.

Typical values of  $k_1$  (20 cm/hr) and  $k_g$  (3,000 cm/hr), which have been measured for CO<sub>2</sub> and H<sub>2</sub>O, respectively, may be used to estimate VOC-specific values for these parameters (Liss and Slater, 1974):

$$k_g(VOC) = k_g(H_2O(18/MW_{VOC})^{0.5}) \quad (2)$$

$$k_1(VOC) = k_1(CO_2(44/MW_{VOC})^{0.5}) \quad (3)$$

where,

MW = molecular weight (g/mol).

The mass transfer coefficient,  $K_L$ , is adjusted to the shower water temperature,  $T_s$ , according to a semi-empirical equation developed to estimate the effect of temperature on oxygen mass-transfer rate (O'Connor and Dobbins, 1956):

$$K_{aL} = K_L(T_{1\mu s}/T_{s\mu s})^{-0.5} \quad (4)$$

where,

$K_L$	=	adjusted overall mass transfer coefficient (cm/hr),
$T_1$	=	calibration water temperature of $K_L$ (K),
$T_s$	=	shower water temperature (K),
$\mu l$	=	water viscosity at $T_1$ (cp), and
$\mu s$	=	water viscosity at $T_s$ (cp).

The concentration leaving the shower droplet,  $C_{wd}$ , is obtained from an integrated rate equation based on a mass-balance approach:

$$C_{wd} = C_{w0}(1 - \exp[-K_{aL}t_s/60d]) \quad (5)$$

where,

$C_{wd}$	=	concentration leaving shower droplet after time $t_s$ ( $\mu g/l$ ),
$C_{w0}$	=	shower water concentration ( $\mu g/l$ ),
$d$	=	shower droplet diameter (mm), and
$t_s$	=	shower droplet drop time (sec).

The term  $K_{aL}/60d$  combines both the rate transfer and the available interfacial area across which volatilization can occur. The value  $1/60d$  equals the specific interfacial area,  $6/d$ , for a special

shower droplet of diameter "d" multiplied by conversion factors (hr/3,600 sec and 10 mm/cm).

The VOC generation rate in the shower room, S, can then be calculated by the equation:

$$S = C_{wd} (FR) / SV \quad (6)$$

where,

- $s$  = indoor VOC generation rate ( $\mu\text{g}/\text{m}^3\text{-min}$ ),  
 $FR$  = shower water flow rate (liter/min), and  
 $SV$  = shower room air volume ( $\text{m}^3$ ).

A simple one-box indoor air pollution model was used to estimate VOC air concentrations in the shower room. This model can be expressed as a differential equation describing the rate of change of the indoor pollutant concentration with time:

$$dC_a/dt = RC_a + S \quad (7)$$

where,

- $C_a$  = indoor VOC air concentration ( $\mu\text{g}/\text{m}^3$ , and  
 $R$  = air exchange rate ( $\text{min}^{-1}$ )).

When Equation 7 is integrated, the time-dependent indoor concentration can be estimated as follows:

$$C_a(t) = (S/R) (1 - \exp [-Rt]) \text{ for } t = D_s$$

and

$$C_a(t) = (S/R) (\exp [RD_s] - 1) \exp (-Rt) \quad t > D_s$$

where,

- $C_a(t)$  = indoor air VOC concentration at time  $t$  ( $\mu\text{g}/\text{m}^3$ ),  
 $D_s$  = shower duration (min), and  
 $t$  = time (min).

The inhalation exposure per shower can then be calculated according to the equation:

$$E_{inh} = [VR / (BW) (10^6)] \int_0^{D_t} C_a(t) dt$$

where,

- $E_{inh}$  = inhalation exposure per shower (mg/kg/shower),  
VR = ventilation rate (liter/min),  
BW = body weight (kg), and  
 $D_t$  = total duration in shower room (min.)

This equation can be solved as:

$$E_{inh} = (VR) (S) / [(BW) (R) (10^6)] [D_s - 1/R + \exp(-RD_s)/R]$$

for the duration of the shower, and as

$$E_{inh} = (VR) (S) / [(BW) (R) (10^6)] \times \left( [D_s + \frac{\exp(-RD_t)}{R}] - \frac{[\exp(R(D_s-D_t)}{R} \right)$$

for both the duration of the shower and the duration in the room after the shower is turned off.

Assuming that an individual showers daily,  $E_{inh}$  is then equivalent to the chronic daily intake.

Table C-1 lists the input parameters to the shower model.

Molecular weights and Henry's Law constants for the contaminants of concern are given in the respective toxicological profiles (see Appendix F).

## LIST OF REFERENCES

- Foster, S. A. and P. C. Chrostowski, 1986. Integrated Household Exposure Model for Use of Tap Water Contaminated with Volatile Organic Chemicals, presented at the 79th Annual Meeting of the Air Pollution Control Association, Minneapolis, Minnesota, June 22-17, 1986.
- Liss, P. S. and P. G. Slater, 1974. Flux of Gases Across the Air-Sea Interface. *Nature* 247:181-184.
- O'Connor, D. J. and W. Dobbins, 1956. The Mechanics of Reaeration in Natural Streams, J. Sanit. Eng. Div., ASCE 82:SA6, In Schroeder, E. D. *Water and Wastewater Treatment*, Chapter 4: Gas Transfer. McGraw-Hill, 1977.

## PARAMETERS USED IN THE CALCULATION OF SHOWER DOSE

PARAMETER	UNITS	VALUE
Calibration water temperature, $T_1$	K	293
Shower water temperature, $T_s$	K	318
Water viscosity at $T_1$ , $\mu_1$	Centipoise	1.002
Water viscosity at $T_s$ , $\mu_s$	Centipoise	0.596
Shower water droplet diameter, d	mm	1.0
Shower droplet drop time, t <sub>d</sub>	sec	2
Shower water flow rate, FR	liter/min	10
Shower room air volume, SV	m <sup>3</sup>	6
Air exchange rate, R	min <sup>-1</sup>	0.0083
Shower duration, D <sub>s</sub>	min	12
Total duration in shower room, D <sub>t</sub>	min	15
Ventilation rate, VR (adult)	liter/min	10
Ventilation rate, VR (child)	liter/min	13.3
Body weight, BW (Adult)	kg	70
Body weight, BW (Child)	kg	15

## APPENDIX O (cont.) SHOWER MODEL

Chemical-specific permeability constants (PCs or  $k_p$ ) were calculated using the following equation. (Reference: USEPA, 1992)

$$\text{Log } k_p = -2.72 + 0.71 \log k_{ow} - 0.0061 \text{ MW}$$

Where,

$k_p$  = permeability constant (cm/hr)

$k_{ow}$  = octanol/water coefficient (unitless)

MW = molecular weight (g/mole)

Parameter	Log $k_{ow}$	MW	$k_p$
acenaphthene	4	154	0.01
2-methylnaphthalene	3.86	142.2	$1.5 \times 10^{-3}$

**APPENDIX P**  
**LEAD UBK MODEL**

---

## **LEAD UBK MODEL**

Lead exposure at the project site was evaluated using USEPA's lead UBK model. This model incorporates exposure from six different media (air, soil, drinking water, diet, indoor dust, and paint) to estimate blood lead levels in infants and young children. The lead UBK model addresses the lowest age groups because children are exceptionally sensitive to the adverse effects of lead. Factors contributing to this sensitivity include: 1) an apparent intrinsic sensitivity of developing organ systems, 2) behavioral traits that result in increased contact with dust and soil, 3) certain physiological factors that result in greater deposition of lead in the respiratory tract and higher absorption rate from the gastrointestinal tract in children, and 4) sufficient transplacental transfer of lead to result in a fetal burden (USEPA, 1990). In contrast to typical exposure estimation techniques, the UBK model predicts blood lead levels in younger children.

The UBK model is flexible in that the user can apply site-specific exposure parameters to estimate blood lead levels. In evaluating exposure at the project site, it was assumed that infants and younger children are exposed to lead in soil, drinking water, diet, air, and indoor dust. The following is a discussion of lead exposure via these five pathways and a description of the exposure parameters used for each pathway.

### *Soil and Indoor Dust*

Soil concentrations of lead at the site were entered into the UBK model to establish the soil and indoor dust contribution to blood lead levels. The UBK model assumes that infants and younger children ingest both soil and indoor dust that is contaminated with lead. For the project site, both RME and average concentrations of lead in soil were modeled. These values are presented in Appendix D. The following discusses the exposure parameters used in estimating the blood lead levels from soil and indoor dust exposure.

The UBK model estimates the indoor lead dust concentration as a percentage of the outdoor soil lead concentration. A conversion factor of 0.28 is used in the model for establishing the indoor lead dust concentration from an outdoor soil source. This value is derived from a study by Davis et al. (1990) where they identified a 0.28 dust/soil ratio for aluminum and silicon. Because site-specific data are not available for the site, the default factor of 0.28 was utilized. The model also distributes the infant and young child's exposure between the soil and indoor dust. This soil/dust ingestion weighting factor is dependent on the amount of time spent indoors and outdoors. It is expected that the amount of time spent indoors is greater than the amount of time spent outdoors for infants and younger children (USEPA, 1990). As a result, the USEPA has selected a default value of 45 (i.e., 45% of lead intake from soil and dust is derived from ingestion of soil, 55% from ingestion of indoor dust).

Soil ingestion rates for the various age groups are required in the prediction of blood lead levels. The default values in Version 0.4 UBK model are 100 mg/day for each of the age groups. These values are unrealistic in that they assume a relatively high ingestion rate for the lower age groups (e.g., 0-1 and 1-2 yr.). As a result, the ingestion rate of 100 mg/day, especially in the lower age groups (0-2 years), is not representative for the site. USEPA's Exposure Factors Handbook (USEPA, 1989) presents values for soil ingestion for the various age groups. In particular, soil ingestion values that represent an intermediate tendency to ingest soil are presented. These values are more representative of the exposure at the site and were used in calculating the blood lead levels. Table K-1 presents the values for each of the age groups.

#### *Drinking Water*

Typically, the UBK model assumes that infants and younger children will be exposed to lead via consumption of drinking water at their place of residence. It was assumed that the younger children and infants will be exposed via direct ingestion of the groundwater at their place of residence. An ingestion rate of 1 L/day, which is consistent with the groundwater pathway, was used in estimating the blood lead contribution of surface water. Both the RME and average groundwater concentrations were used in estimating an overall blood lead level.

#### *Diet*

In creating the UBK model, the USEPA assumed a dietary contribution of lead based on Food and Drug Administration Market Basket Surveys and analysis of food lead content (USEPA, 1990). As a result, the default exposure parameters do not vary based on medium-specific concentrations (i.e., air, soil, water) at the site. The rationale is that foods are harvested from geographically diverse regions of the country and, for the most part, are not produced locally. Consequently, the lead concentrations in the food are not related to the lead levels in the local media. The default values supplied in the UBK model were used in estimating the blood lead contributions from dietary intake.

#### *Air*

Generation of wind blown dust is expected to occur to some extent at the project site. The modelled air concentrations were used as input parameters for the UBK model. These values were used in estimating the blood lead contribution of the air pathway at the project site. The indoor air concentration was calculated as a portion of the outdoor lead level. A default conversion factor of 0.3 was applied to the outdoor concentration to derive an indoor air lead level. Default values also were used for the amount of time spent indoors versus outdoors and the daily breathing volumes for each of the age groups.

**RISK ASSESSMENT REPORT FOR THE CROWN  
INDUSTRIES  
UBK SOIL INGESTION RATES**

AGE (Years)	SOIL INGESTION RATES (mg/day)	
	Default UBK Values	Site Specific Values*
0-1	100	0
1-2	100	50
2-3	100	200
3-4	100	200
4-5	100	50
5-6	100	10
6-7	100	10

\* Intermediate tendency ingestion rate values from the Exposure Factors Handbook (USEPA, 1989).

## REFERENCES

Davis, et. al., 1990. Davis, S., P. Waller, R. Buschborn, J. Ballou and P. White. Quantitative Estimates of Soil Ingestion in Normal Children Between the Ages of 2 and 7 years: Population-Based Estimates Using Aluminum, Silicon, and Titanium as Soil Tracer Elements. Arch. Environ. Health. 1990.

USEPA, 1990. U.S. Environmental Protection Agency. Technical Support Document on Lead. ECAO-CIN-757. Office of Health and Environmental Assessment. Cincinnati, Ohio: September, 1990.

USEPA, 1989. U.S. Environmental Protection Agency. Exposure Factors Handbook. EPA/600/8-89-043. Office of Health and Environmental Assessment. Washington, D.C.: July, 1989.

**APPENDIX Q**  
**SITE CONCEPTUAL MODEL**

---



bcc: APPajak/CF; WDTimbath/JWMentz/PRGF; RBonelli /PRJF; RWatras(ck);RSanPedro; Daily File  
SO# 62470-231  
Subfile: 2  
Initials RPW /

**Baker Environmental, Inc.**  
Airport Office Park, Building 3  
420 Rouser Road  
Coraopolis, Pennsylvania 15108

(412) 269-6000  
FAX (412) 269-2002

August 29, 1994

United States Environmental Protection Agency  
Region IV  
Waste Management Division  
345 Courtland Street  
Atlanta, GA 30365

Attn: Ms. Gena Townsend  
Project Manager

Re: Contract N62470-89-D-4814  
Navy CLEAN, District III  
Contract Task Order (CTO) 0231  
Preliminary Conceptual Evaluation Model  
Operable Unit No. 7 (Sites 1, 28 and 30)  
MCB, Camp Lejeune, North Carolina

Dear Ms. Townsend:

Please find enclosed a copy of the preliminary conceptual model of the human health risk assessment for Operable Unit No. 7, MCB Camp Lejeune. At your convenience, please provide a copy of the model for Mr. Kevin Koporec. The sites addressed in this model are Site 1 - French Creek Liquids Disposal Area; Site 28 - Hadnot Point Burn Dump; and Site 30 - Sneads Ferry Road Fuel Tank Sludge Area.

The model briefly describes the sites and presents the potential sources of contamination, constituents of potential concern, potential human receptors and exposure pathways at each site. In addition, a brief description of the risk assessment approach is provided.

Please review this document and provide comments as necessary. If you or Mr. Koporec have any questions and/or concerns, please feel free to call me at (412)269-2033 or Rachel San Pedro, risk assessment coordinator, at (412)269-6079.

Sincerely,

BAKER ENVIRONMENTAL, INC.

*Richard E. Bonelli*  
Richard E. Bonelli  
Project Manager

REB/lq  
Enclosure

cc: Ms. Lee Anne Rapp, Code 183 (w/o enclosure)  
Ms. Katherine Landman, Code 1823 (w/o enclosure)  
Mr. Neal Paul, MCB, Camp Lejeune (with enclosure)



A Total Quality Corporation

**PRELIMINARY HUMAN HEALTH  
RISK ASSESSMENT  
CONCEPTUAL EVALUATION MODEL FOR  
SITES 1, 28 AND 30**

**MARINE CORPS BASE  
CAMP LEJEUNE, NORTH CAROLINA**

**CONTRACT TASK ORDER 0231**

**AUGUST 29, 1994**

*Prepared for:*

**DEPARTMENT OF THE NAVY  
ATLANTIC DIVISION  
NAVAL FACILITIES  
ENGINEERING COMMAND**

*Norfolk, Virginia*

*Under:*

**LANTDIV CLEAN Program  
Contract N62470-89-D-4814**

*Prepared by:*

**BAKER ENVIRONMENTAL, INC.  
Coraopolis, Pennsylvania**

## TABLE OF CONTENTS

	<u>Page</u>
<b>INTRODUCTION .....</b>	<b>1</b>
<b>APPROACH .....</b>	<b>1</b>
<b>SITE 1 - FRENCH CREEK LIQUIDS DISPOSAL AREA .....</b>	<b>5</b>
Background .....	5
Current and Future Exposure Scenarios .....	5
<b>SITE 28 - HADNOT POINT BURN DUMP .....</b>	<b>6</b>
Background .....	6
Current and Future Exposure Scenarios .....	8
<b>SITE 30 - SNEADS FERRY ROAD FUEL TANK SLUDGE AREA .....</b>	<b>9</b>
Background .....	9
Current and Future Exposure Scenarios .....	9
<b>TABLES</b>	
1      Summary of Exposure Dose Input Parameters .....	2
<b>FIGURES</b>	
1      Flowchart of Potential Exposure Pathways and Receptors Site 1: French Creek Liquids Disposal Area .....	7
2      Flowchart of Potential Exposure Pathways and Receptors Site 28: Hadnot Point Burn Dump .....	10
3      Flowchart of Potential Exposure Pathways and Receptors Site 30: Sneads Ferry Road Fuel Tank Sludge Area .....	11

## INTRODUCTION

As part of the review of available site information for use in a risk assessment and feasibility study (RA/FS), a conceptual evaluation model of the sites has been formulated. Originally developed to aid in planning site activities for the remedial investigation (RI), the conceptual site model also can be used to identify potential exposure pathways, exposure points and data needs, including modeling needs, all of which are key elements in a risk assessment.

As part of the RA/FS for CTO-0231, which includes Sites 1, 28 and 30, a conceptual evaluation model was developed for each site. The models briefly describe the sites and present potential sources of contamination, the constituents present, potentially contaminated media, the routes of migration, and potential exposure pathways at each site. Ecological receptors were not be addressed in this model. The model was developed in accordance with the guidance provided in USEPA Data Quality Objectives for Remedial Response Activities Development Process (USEPA, 1987).

## APPROACH

For the baseline human health risk assessment, both current and future land use exposure scenarios will be assumed for each site. A reasonable maximum exposure (RME) case scenario will be utilized in the assessments. Consequently, the exposure scenarios presented will include RME assumptions for the input parameters in the exposure dose equations. Table 1 is a summary of these values.

The baseline risk assessment for each site will be conducted in concordance with the USEPA documents: Risk Assessment Guidance for Superfund: Human Health Evaluation Manual, Part A and "Supplemental Risk Assessment Guidance for Superfund, Volume II, Environmental Evaluation Manual."

The documents to be used in the assessment include, but are not limited to the following references: Risk Assessment Guidance for Superfund: Development of Risk-based Preliminary Remediation Goals, Part B; "Human Health Evaluation Manual, Supplemental Guidance: Standard Default Exposure Factors"; Exposure Factors Handbook; Dermal Exposure Assessment: Principles and Applications, Interim Report; and Superfund Exposure Assessment Manual (SEAM, 1988).

Toxicity values will be obtained primarily from the Region III Risk-Based Concentration Table, which is based on IRIS, HEAST and provisional and/or recommended USEPA toxicity values, in accordance with Region IV recommendations.

Several mathematical models to estimate exposure may be used. To estimate exposure from the inhalation of volatile contaminants in groundwater while showering, the "Integrated Household Exposure Model for Use of Tap Water Contaminated with Volatile Organic Chemicals" developed by S.A. Foster and P.C. Chrostowski will be applied. To evaluate the health effects of lead, the USEPA lead uptake/biokinetic model will be used. The model addresses the lowest age groups because children are exceptionally sensitive to the adverse effects of lead.

**TABLE 1**  
**SUMMARY OF EXPOSURE DOSE INPUT PARAMETERS**

Input Parameter	Units	Receptor			
		Child	Adult	Military Personnel	Construction Worker
<b>Soil (mg/kg)</b>					
Ingestion Rate, IR	mg/d	200	100	100	480
Fraction Ingested, FI	unitless	1	1	1	1
Exposure Frequency, EF	d/y	350	350	350	90
Exposure Duration, ED	y	6	24	4	1
Surface Area, SA	cm <sup>2</sup>	2,300	5,800	5,800	4,300
Absorption Factor, AF	mg/cm <sup>3</sup>	1	1	1	1
Averaging Time, Noncanc., ATnc	d	2,190	8,760	1,460	365
Averaging Time, Carc., ATcanc	d	25,550	25,550	25,550	25,550
Body Weight, BW	kg	15	70	70	70
Conversion Factor, CF	kg/mg	1x10 <sup>-6</sup>	1x10 <sup>-6</sup>	1x10 <sup>-6</sup>	1x10 <sup>-6</sup>
Absorbence Factor, ABS	unitless	Organics = 0.01; Inorganics = 0.001			
<b>Groundwater (mg/L)</b>					
Ingestion Rate, IR	L/d	1	2	NA	NA
Exposure Frequency, EF	d/y	350	350	NA	NA
Exposure Duration, ED	y	6	30	NA	NA
Exposure Time, ET	h/d	0.25	0.25	NA	NA
Surface Area, SA	cm <sup>2</sup>	10,000	23,000	NA	NA
Averaging Time, Noncanc., ATnc	d	2,190	10,950	NA	NA
Averaging Time, Carc., ATcanc	d	25,550	25,550	NA	NA
Conversion Factor, CF	L/cm <sup>3</sup>	0.001	0.001	NA	NA
Body Weight, BW	kg	15	70	NA	NA
<b>Sediment (mg/kg)</b>					
Ingestion Rate, IR	mg/d	200	100	100	NA
Fraction Ingested, FI	unitless	1	1	1	NA
Exposure Frequency, EF	d/y	350	350	350	NA
Exposure Duration, ED	y	6	30	4	NA
Surface Area, SA	cm <sup>2</sup>	2,300	5,800	5,800	NA
Absorption Factor, AF	mg/cm <sup>3</sup>	1	1	1	NA

**TABLE 1 (CONTINUED)**  
**SUMMARY OF EXPOSURE DOSE INPUT PARAMETERS**

Input Parameter	Units	Receptor			
		Child	Adult	Military Personnel	Construction Worker
Averaging Time, Noncanc., ATnc	d	2,190	10,950	1,460	NA
Averaging Time, Canc., ATcanc	d	25,550	25,550	25,550	NA
Body Weight, BW	kg	15	70	70	NA
Conversion Factor, CF	kg/mg	1x10 <sup>-6</sup>	1x10 <sup>-6</sup>	1x10 <sup>-6</sup>	NA
Absorbence Factor, ABS	unitless	Organics = 0.01; Inorganics = 0.001			
<b>Surface Water (mg/L)</b>					
Ingestion Rate, IR	L/h	0.005	0.005	0.005	NA
Exposure Time, ET	h/d	2.6	2.6	2.6	NA
Exposure Frequency, EF	d/y	45	45	45	NA
Exposure Duration, ED	y	6	30	4	NA
Surface Area, SA	cm <sup>2</sup>	2,300	5,800	5,800	NA
Averaging Time, Noncanc., ATnc	d	2,190	10,950	1,460	NA
Averaging Time, Canc., ATcanc	d	25,550	25,550	25,550	NA
Conversion Factor, CF	L/cm <sup>3</sup>	0.001	0.001	0.001	NA
<b>Air (mg/m<sup>3</sup>)</b>					
<b>Outdoor Air</b>					
Inhalation Rate, IR	m <sup>3</sup> /d	10	20	20	NA
Exposure Time, ET	h/d	0.25	0.25	0.25	NA
Exposure Frequency, EF	d/y	350	350	350	NA
Exposure Duration, ED	y	6	24	4	NA
Averaging Time, Noncanc., ATnc	d	2,190	8,760	1,460	NA
Averaging Time, Canc., ATcanc	d	25,550	25,550	25,550	NA
Body Weight, BW	kg	15	70	70	NA
<b>Shower Air</b>					
Inhalation Rate, IR	m <sup>3</sup> /h	0.6	0.6	NA	NA
Exposure Time, ET	h/d	0.25	0.25	NA	NA
Exposure Frequency, EF	d/y	350	350	NA	NA
Exposure Duration, ED	y	6	30	NA	NA
Averaging Time, Noncanc., ATnc	d	2,190	10,950	NA	NA
Averaging Time, Canc., ATcanc	d	25,550	25,550	NA	NA

**TABLE 1 (CONTINUED)**  
**SUMMARY OF EXPOSURE DOSE INPUT PARAMETERS**

Input Parameter	Units	Receptor			
		Child	Adult	Military Personnel	Construction Worker
Body Weight, BW	kg	15	70	NA	NA
<b>Fish (mg/kg)</b>					
Ingestion Rate, IR	kg/d	NA	0.54	NA	NA
Fraction Ingested, FI	unitless	NA	1	NA	NA
Exposure Frequency, EF	d/y	NA	250	NA	NA
Exposure Duration, ED	y	NA	30	NA	NA
Averaging Time, Noncanc., ATnc	d	NA	10,950	NA	NA
Averaging Time, Canc., ATcanc	d	NA	25,550	NA	NA
Body Weight, BW	kg	NA	70	NA	NA

The acceptable cancer risk range, as stipulated by the USEPA, is  $10^{-4}$  to  $10^{-6}$ . For these assessments, cancer risks exceeding  $10^{-6}$  will be considered unacceptable as protective of human health. The total noncarcinogenic acceptable risk level is a hazard index less than or equal to one. This value depicts a level at or below which adverse systemic effects are not expected to occur in the exposed population.

## **SITE 1 - FRENCH CREEK LIQUIDS DISPOSAL AREA**

### **Background**

Historically, waste liquids including petroleum, oil, lubricants and battery acids, were directly disposed into the ground at this site. The groundwater, surface water, and sediment of this site were initially investigated in 1984, 1986 and 1993 (groundwater only). Contaminants detected included VOCs, SVOCs, and metals. Specific constituents with elevated levels were phenol, cadmium, chromium and trichloroethene (TCE). In 1986, surface water and sediment were sampled. Results indicated elevated levels of phenol and chromium. In 1991, soils were analyzed for chemical constituents. Elevated levels of toluene, benzo(a)pyrene (B[a]P), chromium, lead, nickel and zinc were detected.

The most recent sampling event included investigation of the background surface and subsurface soil, the surface and subsurface soil from the northern and southern portions of the site, and the shallow and deep groundwater, including a potable water supply well. Surface water/sediment samples also were collected from a drainage ditch, but will not be evaluated for risk. The sediment samples will be included with the surface soils.

Although samples were collected from the northern and southern portions of the site and designated as such, they will be combined for this human health risk assessment and evaluated as a single data set since both historically and presently similar operations and processes occurred at both areas.

### **Current and Future Exposure Scenarios**

At present, the site is divided into a northern and southern portion. The northern portion is used for vehicle maintenance (Building FC-120) and also consists of an office building for the Landing Support Battalion. The southern portion of the site is primarily used for equipment, various waste and vehicle storage. There also are office buildings in this portion of the site which are associated with the Marine Corps 8th Engineers and Bridge Support Company. The majority of the site consists of a mixture of paved or coarse gravel road surfaces. Lawned and wooded areas border the site to the north, east and west. Consequently, current receptors will be on-site military personnel. The potential exposure pathways will be surface soil incidental ingestion, dermal contact and inhalation of fugitive dust from vehicular traffic.

At present, the groundwater at the site is not used for potable purposes. A water supply well, HP-638, is located on site, but was shut down in 1993 due to benzene contamination (2 ppb). Consequently, exposure to groundwater is not considered to be viable at the site. Exposure to subsurface soil is not expected for the on-site receptors. Surface water samples will not be evaluated as part of the assessment, since the surface water was collected from a drainage ditch, which is not considered to be a significant source of surface water exposure. The ditches receive surface water runoff from the nearby parking lots. Groundwater does not directly discharge into these ditches.

The sediment results from the ditches will be incorporated with the surface soil results and evaluated as part of the surface soil exposure pathway.

Trespassers are not considered to be a viable receptor population. The southern portion of this site is guarded by a sentry. The northern portion of the site is primarily surrounded by a perimeter fence and is capped with asphalt or concrete in most places. These security measures are deemed sufficient to discourage and minimize trespassing. It is important to note that there is a hot dog stand within the southern portion of the site, manned by a civilian attendant. However, it is assumed that this stand is frequented by the on-site military personnel and/or visitors. It is assumed that the time spent on-site by visitors will be minimal. Consequently, visitors will not be evaluated as part of the assessment of this site.

In the future case, it is expected that the site will remain a military restricted area. As stated previously, groundwater is currently not used for potable purposes. As a result, groundwater exposure will not be assessed for future military personnel.

It is unlikely that a future residence will be implemented at this site. However, to be conservative, groundwater exposure to a child and adult residential receptor will be assessed. It is assumed that a private well could be installed on-site in the future case. The potential exposure pathways are ingestion, dermal contact and inhalation while showering. Similarly, surface soil exposure via ingestion, dermal contact and inhalation will be evaluated for the future residential child and adult receptor.

Similarly, subsurface soil exposure as a result of excavation due to construction activities that may occur at the site in the future is to be assessed. A future construction worker will be evaluated for subsurface soil ingestion and dermal contact.

Figure 1 presents a flowchart of the potential exposure pathways and receptors at this site.


#### **SITE 28 - HADNOT POINT BURN DUMP**

##### **Background**

In the past, this site was used for solid waste disposal. This solid waste, which consisted of industrial solid waste, refuse, garbage, oil-based paints, was burned and covered with fill. The area is now grass-covered.

The site is located adjacent to the Mainside sewage treatment plant, which is enclosed with a six-foot high chain-link fence. Codgels Creek divides the site into east and west areas. The New River borders the site in the southwest. Currently, the site is a recreational/picnic area. The area is frequented by adults and children for recreational purposes. An on-site pond, Orde Pond, is stocked with fish and used for recreational fishing. Military personnel conduct physical training, equipment assembly and other related activities at this site.

**FIGURE 1**  
**FLOWCHART OF POTENTIAL EXPOSURE PATHWAYS AND RECEPTORS**  
**SITE 1: FRENCH CREEK LIQUIDS DISPOSAL AREA**



Field investigations conducted in 1984, 1986 and 1993 revealed specific contaminants in the groundwater, surface water, sediment and fish tissue. Of the four wells sampled, the monitoring well situated beside the New River indicated the most contamination. VOCs, pesticides and metals were prevalent. These contaminants included 1,2-dichloroethene, trichloroethene, vinyl chloride, 4,4'-DDD, 4,4'-DDE, dieldrin, arsenic, lead and mercury. The other three wells indicated the presence of SVOCs, pesticides and chromium. Chromium and zinc were detected at elevated levels in the surface water samples collected from the New River.

Recent field investigations included background surface and subsurface soils, site surface and subsurface soils, and shallow and deep groundwater; surface water, sediment, fish tissue and benthic macroinvertebrates were collected from Orde Pond, the New River and Cogdels Creek. Although media samples were collected from the eastern and western portions of the site and designated accordingly, the results will evaluated as a single data set per medium. Similar historical operations occurred at both the eastern and western portions of the site, and current activities do not differ between these two areas.

#### **Current and Future Exposure Scenarios**

At Site 28, on-site military personnel and recreational adults and children frequent the area. In addition, anglers mainly fish at Orde Pond and the New River. Fishing at Cogdels Creek is less frequent. No swimming has been observed at any of the surface water bodies, although wading is possible. Signs prohibiting swimming are posted at Orde Pond. However, it is important to note that the on-site military personnel conduct training exercises at this site, including activities in Orde Pond.

In the current exposure scenario, potential receptors include on-site military personnel, recreational adults and children, and fishermen. Potential exposure pathways include surface soil ingestion, dermal contact and inhalation for the military personnel and recreational receptor population. Recreational adults and children are expected to become exposed to surface water and sediment while playing/wading in the New River and possibly Cogdels Creek. Fishermen may become exposed to these media while fishing at Orde Pond and the New River, with less frequent exposure at Cogdels Creek. It is assumed that exposure to surface water and sediment at Cogdels Creek and the New River will be minimal for the on-site military personnel during physical training exercises. However, exposure to the surface water and sediment of Orde Pond via incidental ingestion and dermal contact will be evaluated for the on-site military personnel. In addition, fish ingestion for the adult receptor (fisherman) will be evaluated.

At present, groundwater is not utilized for potable purposes. As a result, current groundwater exposure will not be assessed. Exposure to subsurface soil in the current scenario is unlikely for the receptor population. Consequently, subsurface soil exposure is not considered to be viable.

In the future case, it is unlikely that a residential scenario will be implemented at the site. It is assumed that the present activities will continue into the foreseeable future. However, to be conservative, groundwater exposure to a residential child and adult receptor will be assessed. Surface soil, surface water, sediment and biota exposure, as calculated in the current scenario for the child and adult receptor, is expected to remain the same in the future case.

Similar to Site 1, groundwater exposure for future on-site military personnel will not be assessed. However, a construction worker will be evaluated in the future case. It is assumed that subsurface soil exposure may occur as a result of excavation for potential construction activities at the site.

Figure 2 presents a flowchart of the potential exposure pathways and receptors at this site.

## **SITE 30 - SNEADS FERRY ROAD FUEL TANK SLUDGE AREA**

### **Background**

Historically, sludge collected from storage tanks containing fuels such as leaded gasoline were disposed of in this area. In addition, cleaning solvents used as part of the sludge removal process were disposed.

In 1984, 1986 and 1993, the groundwater was sampled and analyzed for contaminants. Levels of methylene chloride, chloroform, cadmium, chromium, lead and mercury were detected. In the surface water and sediment samples collected from French Creek in 1986, no VOCs or metals were detected.

The most recent sampling event investigated the following media: background surface and subsurface soil, site surface and subsurface soil, groundwater, surface water and sediment.

### **Current and Future Exposure Scenarios**

At present, the site is not used by the on-site military personnel for training. However, the area north of the site is used for the training exercises by military personnel, and an access road is located on the northern border of the site. Access to the site by trespassers is not restricted. French Creek and Sneads Ferry Road border the site. The site is sandy with some wooded areas.

Potential receptors are the on-site military personnel. It is unlikely that trespassers, both adults and children, will be potential receptors because the area is primarily used for military training exercises. The potential exposure pathways include surface soil incidental ingestion, dermal contact and inhalation. Exposure to the surface water and sediment in French Creek via incidental ingestion and dermal contact are also viable for the current receptors.

The groundwater at the site is not potable. Hence, groundwater exposure will not be assessed. Subsurface soil exposure is not considered to be a viable pathway in the current scenario.

Similar to Sites 1 and 28, a future residential exposure scenario is not likely at Site 30. The site is a military restricted area. It is assumed that the present activities at this site will continue into the foreseeable future. However, to be conservative, groundwater exposure to a residential child and adult will be assessed. Similarly, subsurface soil exposure to a construction worker will be evaluated.

Figure 3 presents a flowchart of the potential exposure pathways and receptors at this site.

FIGURE 2  
FLOWCHART OF POTENTIAL EXPOSURE PATHWAYS AND RECEPTORS  
SITE 28: HADNOT POINT BURN DUMP





FIGURE 3

FLOWCHART OF POTENTIAL EXPOSURE PATHWAYS AND RECEPTORS  
SITE 30: SNEADS FERRY ROAD FUEL TANK SLUDGE AREA



---

**APPENDIX R**  
**CDI CALCULATIONS**

**EXAMPLE SOIL\* INGESTION CALCULATIONS  
OPERABLE UNIT NO. 7  
CONTRACT TASK ORDER 0231**

**Purpose:** Estimate intake/risk from ingestion of soil

$$\text{Intake (mg/kg-day)} = \frac{C \times CF \times EF \times ED \times IR}{BW \times AT}$$

Where:

C	=	Contaminant concentration in soil (mg/kg)
CF	=	Conversion factor (kg/mg)
EF	=	Exposure frequency (days/year)
ED	=	Exposure duration (years)
IR	=	Ingestion rate (mg/day)
BW	=	Body weight (kg)
AT <sub>c</sub>	=	Averaging time carcinogen (days)
AT <sub>nc</sub>	=	Averaging time noncarcinogen (days)

**Risks:**

$$\text{Carcinogens} = \text{Intake (mg/kg-day)} \times \text{CSF (mg/kg-day)}^{\dagger}$$

$$\text{Noncarcinogens} = \text{Intake (mg/kg-day)} / \text{RfD (mg/kg-day)}$$

**Example Carcinogen:** Arsenic

$$\begin{aligned} \text{Intake (mg/kg-day)} &= \frac{1.2 \text{ mg/kg} \times 100 \text{ mg/day} \times 350 \text{ days/yr} \times 30 \text{ yrs} \times 1.0E-6 \text{ kg/mg}}{70 \text{ kg} \times 25,550 \text{ days}} \\ &= 7E-07 \end{aligned}$$

$$\text{Risk} = 5E-09 \text{ mg/kg-day} \times 1.75 \text{ mg/kg-day}^{\dagger} = 1.2E-06$$

**Example Noncarcinogen:** Barium

$$\begin{aligned} \text{Intake (mg/kg-day)} &= \frac{9.1 \text{ mg/kg} \times 100 \text{ mg/day} \times 350 \text{ days/yr} \times 30 \text{ yrs} \times 1.0E-6 \text{ kg/mg}}{70 \text{ kg} \times 10,950 \text{ days}} \\ &= 1.2E-05 \\ \text{Risk} &= \frac{1.2E-05 \text{ mg/kg-day}}{7E-02 \text{ mg/kg-day}} = 1.8E-04 \end{aligned}$$

\* This example calculation also is applicable for sediment ingestion.  
Re: Site 1 Future Residential Adult

**EXAMPLE DERMAL CONTACT WITH SOIL\* CALCULATIONS  
OPERABLE UNIT NO. 7  
CONTRACT TASK ORDER 0231**

**Purpose:** Estimate intake/risk from dermal contact with soil

$$\text{Intake (mg/kg-day)} = \frac{C \times CF \times SA \times AF \times ABS \times EF \times ED}{BW \times AT}$$

Where:

C	=	Contaminant concentration in soil (mg/kg)
CF	=	Conversion factor (kg/mg)
SA	=	Surface available for contact (cm <sup>2</sup> /event)
AF	=	Soil to skin adherence factor (mg/cm <sup>2</sup> )
ABS	=	Fraction absorbed (percent) - 0.01 organics, 0.001 inorganics
EF	=	Exposure frequency (days/year)
ED	=	Exposure duration (years)
IR	=	Ingestion rate (mg/day)
BW	=	Body weight (kg)
AT <sub>c</sub>	=	Averaging time carcinogen (days)
AT <sub>nc</sub>	=	Averaging time noncarcinogen (days)

**Risks:**

Carcinogens = Intake (mg/kg·day) x dermally - adjusted CSF (mg/kg·day)<sup>-1</sup>

Noncarcinogens = Intake (mg/kg·day)/ dermally - adjusted RfD (mg/kg·day)

**Example Carcinogen: Arsenic**

$$\begin{aligned} \text{Intake (mg/kg-day)} &= \frac{1.2 \text{ mg/kg} \times 1.0E-06 \text{ kg/mg} \times 5,800 \text{ cm}^2/\text{event} \times 0.001 \times 1 \text{ mg/cm}^2 \times 350 \text{ days/yr} \times 30 \text{ yrs}}{70 \text{ kg} \times 25,550 \text{ days}} \\ &= 4.1E-08 \end{aligned}$$

$$\text{Risk} = 4.1E-08 \text{ mg/kg-day} \times 8.8 \text{ mg/kg-day}^{-1} = 3.6E-07$$

**Example Noncarcinogen: Cadmium**

$$\begin{aligned} \text{Intake (mg/kg-day)} &= \frac{0.32 \text{ mg/kg} \times 1.0E-06 \text{ kg/mg} \times 5,800 \text{ cm}^2/\text{event} \times 1 \text{ mg/cm}^2 \times 0.001 \times 350 \text{ days/yr} \times 30 \text{ yrs}}{70 \text{ kg} \times 10,950 \text{ days}} \\ &= 2.5E-08 \end{aligned}$$

$$\text{Risk} = \frac{2.5E-08 \text{ mg/kg-day}}{1E-04 \text{ mg/kg-day}} = 2.5E-04$$

\* This example calculation also is applicable for sediment dermal contact.  
Re: Site 1 Future Residential Adult

**EXAMPLE INHALATION OF PARTICULATES CALCULATIONS  
OPERABLE UNIT NO. 7  
CONTRACT TASK ORDER 0231**

**Purpose:** Estimate intake/risk from the inhalation of soil particulates

$$\text{Intake (mg/kg-day)} = \frac{C \times IR \times EF \times ED \times 1/\text{PEF}}{BW \times AT}$$

Where:

C	=	Contaminant concentration in soil (mg/kg)
IR	=	Inhalation rate ( $\text{m}^3/\text{day}$ )
EF	=	Exposure frequency (days/year)
ED	=	Exposure duration (years)
PEF	=	Particulate Emission Factor ( $\text{m}^3/\text{kg}$ )
BW	=	Body weight (kg)
AT <sub>c</sub>	=	Averaging time carcinogen (days)
AT <sub>nc</sub>	=	Averaging time noncarcinogen (days)

**Risks:**

$$\text{Carcinogens} = \text{Intake (mg/kg-day)} \times \text{CSF (mg/kg-day)}^{-1}$$

$$\text{Noncarcinogens} = \text{Intake (mg/kg-day)} / \text{RfD (mg/kg-day)}$$

**Example Carcinogen: Arsenic**

$$\text{Intake (mg/kg-day)} = \frac{1.2 \text{ mg/kg} \times 20 \text{ m}^3/\text{day} \times 350 \text{ days/yr} \times 30 \text{ yrs} \times 1/4.6E+09 \text{ m}^3/\text{kg}}{70 \text{ kg} \times 25,550 \text{ days}}$$

$$= 3.1\text{E-}11$$

$$\text{Risk} = 3.1\text{E-}11 \text{ mg/kg-day} \times 15.1 \text{ mg/kg-day}^{-1} = 4.6\text{E-}10$$

**Example Noncarcinogen: Manganese**

$$\text{Intake (mg/kg-day)} = \frac{8.5 \text{ mg/kg} \times 20 \text{ m}^3/\text{day} \times 350 \text{ days/yr} \times 30 \text{ yrs} \times 1/4.6E+09 \text{ m}^3/\text{kg}}{70 \text{ kg} \times 10,950 \text{ days}}$$

$$= 5.1\text{E-}10$$

$$\text{Risk} = \frac{5.1\text{E-}10 \text{ mg/kg-day}}{1.4E-05 \text{ mg/kg-day}} = 3.6E-05$$

**EXAMPLE GROUNDWATER INGESTION CALCULATIONS  
OPERABLE UNIT NO. 7  
CONTRACT TASK ORDER 0231**

**Purpose:** Estimate intake/risk from ingestion of groundwater

$$\text{Intake (mg/kg-day)} = \frac{C \times IR \times EF \times ED}{BW \times AT}$$

Where:

C	=	Contaminant concentration in groundwater (mg/L)
IR	=	Daily intake ingestion rate (L/day)
EF	=	Exposure frequency (days/year)
ED	=	Exposure duration (years)
BW	=	Body weight (kg)
AT <sub>c</sub>	=	Averaging time carcinogen (days)
AT <sub>nc</sub>	=	Averaging time noncarcinogen (days)

Risks:

$$\text{Carcinogens} = \text{Intake (mg/kg-day)} \times \text{CSF (mg/kg-day)}^{-1}$$

$$\text{Noncarcinogens} = \text{Intake (mg/kg-day)} / \text{RfD (mg/kg-day)}$$

**Example Carcinogen: Arsenic**

$$\text{Intake (mg/kg-day)} = \frac{2.7E-03 \text{ mg/L} \times 1 \text{ L/day} \times 350 \text{ days/yr} \times 6 \text{ yrs}}{15 \text{ kg} \times 25,550 \text{ days}}$$

$$= 1.5E-05$$

$$\text{Risk} = 1.5E-05 \text{ mg/kg-day} \times 1.75 \text{ mg/kg-day}^{-1} = 2.6E-05$$

**Example Noncarcinogen: 4,4'-DDT**

$$\text{Intake (mg/kg-day)} = \frac{1.8E-04 \text{ mg/L} \times 1 \text{ L/day} \times 350 \text{ days/yr} \times 6 \text{ yrs}}{15 \text{ kg} \times 2,190 \text{ days}}$$

$$= 1.2E-05$$

$$\text{Risk} = \frac{1.2E-05 \text{ mg/kg-day}}{5E-04 \text{ mg/kg-day}} = 2.3E-02$$

**EXAMPLE DERMAL CONTACT WITH GROUNDWATER CALCULATIONS  
OPERABLE UNIT NO. 7  
CONTRACT TASK ORDER 0231**

**Purpose:** Estimate intake/risk from dermal contact with groundwater

$$\text{Intake (mg/kg-day)} = \frac{C \times CF \times SA \times PC \times ET \times EF \times ED}{BW \times AT}$$

Where:

C	=	Contaminant concentration in groundwater (mg/L)
CF	=	Conversion factor (1 L/1,000 cm <sup>3</sup> )
SA	=	Exposed skin surface available for contact (cm <sup>2</sup> )
PC	=	Chemical-specific dermal permeability constant (cm/hr)
ET	=	Exposure time (hr/day)
EF	=	Exposure frequency (days/year)
ED	=	Exposure duration (years)
IR	=	Ingestion rate (L/day)
BW	=	Body weight (kg)
AT <sub>c</sub>	=	Averaging time carcinogen (days)
AT <sub>nc</sub>	=	Averaging time noncarcinogen (days)

Risks:

$$\text{Carcinogens} = \text{Intake (mg/kg-day)} \times \text{CSF (mg/kg-day)}^{\text{l}}$$

$$\text{Noncarcinogens} = \text{Intake (mg/kg-day)} / \text{RfD (mg/kg-day)}$$

**Example Carcinogen: Arsenic**

$$\text{Intake (mg/kgday)} = \frac{2.7E-03 \text{ mg/L} \times 1.0E-03 \text{ L/cm}^3 \times 10,000 \text{ cm}^2/\text{event} \times 1.0E-03 \text{ cm/hr} \times 0.25 \text{ hr/day} \times 350 \text{ days/yr} \times 6 \text{ yrs}}{15 \text{ kg} \times 25,550 \text{ days}}$$

$$= 3.7E-08$$

$$\text{Risk} = 3.7E-08 \text{ mg/kg-day} \times 8.8 \text{ mg/kg-day}^{\text{l}} = 3.3E-07$$

**Example Noncarcinogen: Mercury**

$$\text{Intake (mg/kg-day)} = \frac{5E-04 \text{ mg/L} \times 1.0E-03 \text{ L/cm}^3 \times 10,000 \text{ cm}^2/\text{event} \times 1.0E-03 \text{ cm/hr} \times 0.25 \text{ hr/day} \times 350 \text{ days/yr} \times 6 \text{ yrs}}{15 \text{ kg} \times 2,190 \text{ days}}$$

$$= 8E-08$$

$$\text{Risk} = \frac{8E-08 \text{ mg/kg-day}}{6E-05 \text{ mg/kg-day}} = 1.3E-03$$

**EXAMPLE INHALATION OF GROUNDWATER VOLATILES CALCULATIONS  
OPERABLE UNIT NO. 7  
CONTRACT TASK ORDER 0231**

**Purpose:** Estimate intake/risk from the inhalation of groundwater volatiles

$$\text{Intake (mg/kg-day)} = \frac{C \times EF \times ED}{AT}$$

Where:	C	=	Contaminant concentration in shower air (mg/kg/shower) - Foster Model (Appendix O)
	EF	=	Exposure frequency (days/year)
	ED	=	Exposure duration (years)
	AT <sub>c</sub>	=	Averaging time carcinogen (days)
	AT <sub>nc</sub>	=	Averaging time noncarcinogen (days)

**Risks:**

Carcinogens = Intake (mg/kg·day) x CSF (mg/kg·day)<sup>-1</sup>

Noncarcinogens = Intake (mg/kg·day)/RfD (mg/kg·day)

**Example Carcinogen:** Trichloroethene

$$\text{Intake (mg/kg-day)} = \frac{4E-05 \text{ mg/kg/shower} \times 350 \text{ days/yr} \times 24 \text{ yrs}}{25,550 \text{ days}}$$

$$= 1.7E-05$$

$$\text{Risk} = 1.7E-05 \text{ mg/kg-day} \times 6E-03 \text{ mg/kg-day}^{-1} = 9.9E-08$$

**Example Noncarcinogen:** None identified as a COPC

**EXAMPLE SURFACE WATER INGESTION CALCULATIONS  
OPERABLE UNIT NO. 7  
CONTRACT TASK ORDER 0231**

**Purpose:** Estimate intake/risk from ingestion of surface water

$$\text{Intake (mg/kg-day)} = \frac{C \times IR \times EF \times ED \times ET}{BW \times AT}$$

Where:

C	=	Contaminant concentration in surface water (mg/L)
IR	=	Daily intake ingestion rate (kg/meal)
EF	=	Exposure frequency (meal/year)
ED	=	Exposure duration (years)
ET	=	Exposure time (hrs/day)
BW	=	Body weight (kg)
AT <sub>c</sub>	=	Averaging time carcinogen (days)
AT <sub>nc</sub>	=	Averaging time noncarcinogen (days)

Risks:

$$\text{Carcinogens} = \text{Intake (mg/kg-day)} \times \text{CSF (mg/kg-day)}^{-1}$$

$$\text{Noncarcinogens} = \text{Intake (mg/kg-day)} / \text{RfD (mg/kg-day)}$$

**Example Carcinogen: Arsenic**

$$\begin{aligned} \text{Intake (mg/kgday)} &= \frac{4.3E-03 \text{ mg/L} \times 0.005 \text{ L/day} \times 48 \text{ days/yr} \times 30 \text{ yrs} \times 2.6 \text{ hrs/day}}{70 \text{ kg} \times 25,550 \text{ days}} \\ &= 4.5E-08 \end{aligned}$$

$$\text{Risk} = 4.5E-08 \text{ mg/kg-day} \times 1.75 \text{ mg/kg-day}^{-1} = 7.9E-08$$

**Example Noncarcinogen: Cadmium**

$$\begin{aligned} \text{Intake (mg/kgday)} &= \frac{4.2E-03 \text{ mg/L} \times 0.005 \text{ L/day} \times 48 \text{ days/yr} \times 30 \text{ yrs} \times 2.6 \text{ hrs/day}}{70 \text{ kg} \times 10,950 \text{ days}} \\ &= 1E-7 \end{aligned}$$

$$\text{Risk} = \frac{1E-7 \text{ mg/kgday}}{5E-04 \text{ mg/kgday}} = 2.1E-04$$

**EXAMPLE SURFACE WATER DERMAL CONTACT CALCULATIONS  
OPERABLE UNIT NO. 7  
CONTRACT TASK ORDER 0231**

**Purpose:** Estimate intake/risk from dermal contact with surface water

$$\text{Intake (mg/kgday)} = \frac{C \times CF \times EF \times ED \times SA \times ET}{BW \times AT}$$

Where:

C	=	Contaminant concentration in surface water (mg/L)
SA	=	Skin surface area (cm <sup>2</sup> )
CF	=	Conversion factor (1 L/1,000 cm <sup>3</sup> )
EF	=	Exposure frequency (days/year)
ED	=	Exposure duration (years)
ET	=	Exposure time (hrs/day)
IR	=	Ingestion rate (L/day)
PC	=	Chemical-specific dermal permeability constant (cm/hr)
BW	=	Body weight (kg)
AT <sub>c</sub>	=	Averaging time carcinogen (days)
AT <sub>nc</sub>	=	Averaging time noncarcinogen (days)

**Risks:**

$$\text{Carcinogens} = \text{Intake (mg/kg·day)} \times \text{CSF (mg/kg·day)}^{-1}$$

$$\text{Noncarcinogens} = \text{Intake (mg/kg·day)} / \text{RfD (mg/kg·day)}$$

**Example Carcinogen: 4,4'-DDD**

$$\begin{aligned} \text{Intake (mg/kgday)} &= \frac{5E-05 \text{ mg/L} \times 5,800 \text{ cm}^2 \times 48 \text{ days/yr} \times 30 \text{ yrs} \times 2.6 \text{ hrs/day} \times 1.0E-3 \text{ L/cm}^3 \times 0.001 \text{ cm/hr}}{70 \text{ kg} \times 25,550 \text{ days}} \\ &= 1.75E-07 \end{aligned}$$

$$\text{Risk} = 1.7E-7 \text{ mg/kg·day} \times 0.48 \text{ mg/kg·day}^{-1} = 8.2E-08$$

**Example Noncarcinogen: Arsenic**

$$\begin{aligned} \text{Intake (mg/kgday)} &= \frac{4.3E-03 \text{ mg/L} \times 5,800 \text{ cm}^2 \times 48 \text{ days/yr} \times 30 \text{ yrs} \times 2.6 \text{ hrs/day} \times 1.0E-3 \text{ L/cm}^3 \times 1E-03 \text{ cm/hr}}{70 \text{ kg} \times 10,950 \text{ days}} \\ &= 1.2E-07 \end{aligned}$$

$$\text{Risk} = \frac{1.2E-07 \text{ mg/kgday}}{6E-05 \text{ mg/kgday}} = 2E-03$$

**EXAMPLE FISH INGESTION CALCULATIONS  
OPERABLE UNIT NO. 7  
CONTRACT TASK ORDER 0231**

**Purpose:** Estimate intake/risk from ingestion of soil

$$\text{Intake (mg/kg/day)} = \frac{C \times FI \times EF \times ED \times IR}{BW \times AT}$$

Where:

C	=	Contaminant concentration in fish tissue (mg/kg)
FI	=	Fraction ingested
EF	=	Exposure frequency (meal/year)
ED	=	Exposure duration (years)
IR	=	Ingestion rate (kg/meal)
BW	=	Body weight (kg)
AT <sub>c</sub>	=	Averaging time carcinogen (days)
AT <sub>nc</sub>	=	Averaging time noncarcinogen (days)

**Risks:**

$$\text{Carcinogens} = \text{Intake (mg/kg/day)} \times \text{CSF (mg/kg/day)}^1$$

$$\text{Noncarcinogens} = \text{Intake (mg/kg/day)} / \text{RfD (mg/kg/day)}$$

**Example Carcinogen: 4,4'-DDD**

$$\begin{aligned} \text{Intake (mg/kg/day)} &= \frac{2.3E-02 \text{ mg/kg} \times 0.284 \text{ kg/meal} \times 48 \text{ days/yr} \times 30 \text{ yrs} \times 1.0}{70 \text{ kg} \times 25,550 \text{ days}} \\ &= 5.3E-06 \end{aligned}$$

$$\text{Risk} = 5.3E-06 \text{ mg/kg-day} \times 2.45E-01 \text{ mg/kg-day}^1 = 1.3E-06$$

**Example Noncarcinogen: Antimony**

$$\begin{aligned} \text{Intake (mg/kg/day)} &= \frac{0.24 \text{ mg/kg} \times 0.284 \text{ kg/meal} \times 48 \text{ days/yr} \times 30 \text{ yrs} \times 1.0}{70 \text{ kg} \times 10,950 \text{ days}} \\ &= 1.3E-04 \end{aligned}$$

$$\text{Risk} = \frac{1.3E-04 \text{ mg/kg/day}}{4E-04 \text{ mg/kg/day}} = 0.32$$

**SITE 1**

---

GROUNDWATER INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 1 - SHALLOW AND DEEP GROUNDWATER)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL CHILD

Intake from drinking water is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * IRw * EF * ED/BW * AT \text{ or } ATnc * DY$$

Risk = Intake \* CSF or /RfD

Where:

	INPUTS
C = contaminant concentration in water (mg/l)	
IRw = child daily water ingestion rate (L/Day)	1
EF = child exposure frequency (days/yr)	350
ED = child exposure duration (yr)	6
BW = child body weight (kg)	15
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	6
DY = days per year (day/year)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RfD = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

SHALLOW AND DEEP GROUNDWATER

COPC	Concentration (mg/L)	Ingestion Rate (L/day) Child	Exposure Frequency (day/year) Child	Exposure Duration (year) Child	Body Weight (kg) Child	Average Carc Time (years) Child	Carc Dose (mg/kg-day)-1 Child	Slope Factor (mg/kg-day)-1	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc Time (years)	Noncarc Dose (mg/kg-day) Child	Reference Dose (mg/kg-day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child
Arsenic	8.5E-03	1	350	6	15	25550	4.7E-05	1.75E+00	8.2E-06	100%	2190	5.4E-04	3.00E-04	1.8E+00	10%
Barium	4.5E-02	1	350	6	15	25550	2.5E-04	0.00E+00	0.0E+00	0%	2190	2.9E-03	7.00E-02	4.1E-02	0%
Manganese (water)	1.2E+00	1	350	6	15	25550	6.5E-03	0.00E+00	0.0E+00	0%	2190	7.7E-02	5.00E-03	1.5E+01	88%
Mercury	1.0E-03	1	350	6	15	25550	5.8E-08	0.00E+00	0.0E+00	0%	2190	6.6E-05	3.00E-04	2.2E-01	1%
1,2-Dichloroethene	2.6E-03	1	350	6	15	25550	1.4E-05	0.00E+00	0.0E+00	0%	2190	1.7E-04	8.00E-03	1.8E-02	0%
Trichloroethene	4.1E-03	1	350	6	15	25550	2.2E-05	1.10E-02	2.5E-07	0%	2190	2.6E-04	6.00E-03	4.4E-02	0%
TOTAL								8.2E-05						1.7E+01	

GROUNDWATER DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 1 - SHALLOW AND DEEP GROUNDWATER)  
 REMEDIAL INVESTIGATION CTD-0291  
 MC8 CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL CHILD

Dermal Contact from groundwater is calculated as follows:

$$\text{Intake (mg/kg-day)} = \text{CW} * \text{SA} * \text{PC} * \text{ET} * \text{EF} * \text{ED} * \text{CF/BW} * \text{ATc or ATnc} * \text{DY}$$

Risk = Intake \* CSF or RID

Where:

CW = contaminant concentration in water (mg/l)  
 SA = child skin surface available for contact (cm<sup>2</sup>)  
 PC = contaminant specific dermal permeability (cm/h)  
 ET = child exposure time (hours/day)  
 EF = child exposure frequency (days/yr)  
 ED = child exposure duration (years)  
 CF = volumetric conversion factor for water (liter/1000 cm<sup>3</sup>)  
 BW = child body weight (kg)  
 ATc = averaging time for carcinogen (yr)  
 ATnc = averaging time for noncarcinogen (yr)  
 DY = days per year (days)

INPUTS

Specific  
 10000  
 0.25  
 0.25  
 365  
 6  
 0.001  
 15  
 70  
 6  
 365

Note: Inputs are site and scenario specific.

SHALLOW AND DEEP GROUNDWATER

COPC	Concentration (mg/l)	Surface Area (cm <sup>2</sup> )	Dermal Permeability (cm/h)	Exposure Time (hours/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	Volumetric Conversion (l/m <sup>3</sup> )	Body Weight (kg)	Averaging Carb Time (years)	Carc Dose (mg/kg-day)	Dermat Adjust Slope Factor (mg/kg-day) <sup>-1</sup>	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc Time (years)	Noncarc Dose (mg/kg-day)	Dermat Adjust Reference Dose (mg/kg-day)	Noncarc Risk Child	Percent Noncarcinogenic Risk Child
Arsenic	4.0E-03	10000	1.00E-03	0.25	365	6	0.001	15	25600	1.0E-07	8.7E+00	1.0E+00	99%	2160	7.4E-06	8.0E-03	4.3E-02	10%
Barium	4.0E-02	10000	1.00E-03	0.25	365	6	0.001	15	25600	8.1E-07	0.0E+00	0.0E+00	0%	2160	7.1E-06	1.4E-02	6.1E-04	0%
Manganese (water)	1.2E+03	10000	1.00E-03	0.25	365	6	0.001	15	25600	1.0E-08	0.0E+00	0.0E+00	0%	2160	1.0E-04	1.0E-03	1.0E-07	87%
Mercury	1.0E-03	10000	1.00E-03	0.25	365	6	0.001	15	25600	1.4E-08	0.0E+00	0.0E+00	0%	2160	1.0E-05	8.0E-05	2.7E-03	1%
1,2-Dichloroethene	2.0E-03	10000	1.00E-02	0.25	365	6	0.001	15	25600	3.0E-07	0.0E+00	0.0E+00	0%	2160	4.2E-05	7.2E-03	6.6E-04	0%
Tetrachloroethene	4.1E-03	10000	1.00E-02	0.25	365	6	0.001	15	25600	9.0E-07	1.3E-02	1.2E-08	1%	2160	1.0E-05	4.0E-03	2.2E-03	1%
<b>TOTAL</b>																		2.2E-01

CTO-231 SITE 1 GROUNDWATER INHALATION

01-Jun-95

FILE: SHOW.WQ1

GROUNDWATER INHALATION - RME CASE

CHILD CHEMICAL	C rme mg/kg/shw y	ED shwr/y	EF d	AT d	ATC d	CDI mg/kg/d	CDIC mg/kg/d	RFD mg/kg/d	PF (mg/kg/d)^-1	HI	CR	% CONTRIB NC RISK	% CONTRIB CARC RISK
1, 2-Dichloroethene	1.73E-04	6	350	3285	25550	1.1E-04	1.4E-05	0.0E+00	0.0E+00	---	0.0E+00	0%	0%
Trichloroethene	2.44E-04	6	350	3285	25550	1.6E-04	2.0E-05	0.0E+00	6.0E-03	---	1.2E-07	0%	100%
TOTAL											0.0E+00	1.2E-07	

SHOWMOD.WQ1  
07-Jun-96

## SHOWER EXPOSURE MODEL

CAS No.	TARGET COMPOUND LIST CONSTITUTE	OVERALL MASS TRANSF COEFFICIENT (KL)	HENRY'S LAW CONSTAN (H)	GAS CONSTAN (F)	ABSOLUTE TEMP. (T1)	GAS CONST. * ABS. TEMP. (RT)	GAS-FLM MASS TRANSF. COEFFICIENT (kg)	LIQUID-FLM MASS TRANSF. COEFFICIENT (kg)	CARBON DIOXIDE TRANS. COEFF. (kg-CO2)	WATER TRANS. COEFF. (kg-H2O)	MOLECULAR WEIGHT (gram/mole)	RECIPROCAL OF (K)	(RT)/(H)*(kg)	OVERALL MASS TRANSF. COEFF. (Kal)	WATER VISCOSITY (AT T1) (U1)	WATER VISCOSITY (AT T1) (U1)	SHOWER WATER TEMP. (Ts)	(T1)*(U1)) / (Ts)*(U1))
		(cm/hr)	(atm-m3/mol-k)	(at-m3/mol-k)	(K)	(RT)	(cm/hr)	(cm/hr)	(cm/hr)	(cm/hr)	(gram/mole)	(K)	(cm/hr)	(cp)	(cp)	(K)		
<b>VOLATILE ORGANICS:</b>																		
ADULT																		
540-59-0	1,2-DICHLOROETHENE (total)	1.30E+01	7.58E-03	8.20E-05	293.0	2.40E-02	1.29E+03	1.35E+01	20.0	3000.0	97.00	7.42E-02	2.45E-03	1.76E+01	0.596	1.002	318.0	5.48E-01
79-01-6	TRICHLOROETHENE (B2)	1.13E+01	9.10E-03	8.20E-05	293.0	2.40E-02	1.11E+03	1.16E+01	20.0	3000.0	131.40	8.64E-02	2.38E-03	1.52E+01	0.596	1.002	318.0	5.48E-01
540-59-0	1,2-DICHLOROETHENE (total)	1.30E+01	7.58E-03	8.20E-05	293.0	2.40E-02	1.29E+03	1.35E+01	20.0	3000.0	97.00	7.42E-02	2.45E-03	1.76E+01	0.596	1.002	318.0	5.48E-01
79-01-6	TRICHLOROETHENE (B2)	1.13E+01	9.10E-03	8.20E-05	293.0	2.40E-02	1.11E+03	1.16E+01	20.0	3000.0	131.40	8.64E-02	2.38E-03	1.52E+01	0.596	1.002	318.0	5.48E-01

CAS No.	TARGET COMPOUND LIST CONSTITUENT	CONCENTRATION	SHOWER	SHOWER	SHOWER	KEL (%)	60% (60%)	-10% (-10%)	IMP [KUL/m <sup>3</sup> ] [STC]	INDOOR VOC GENERATION RATE (g/h <sup>2</sup> m <sup>3</sup> )	SHOWER	SHOWER	INDOOR AIR	SHOWER	TOTAL TIME DURATION	IMP/(%)	IMP/(%)	IMPACTION	VENTILATION	BODY	IMP/ADU	DURATION	DA+	IMP/ADU M	DA-D <sub>0</sub>	M + M <sub>0</sub> (M <sub>0</sub> /M)	
		LEAVING WATER	DROPLET CONCENTRATION	DROPLET DIAMETER	DROPLET DROP TIME	d	w				WATER FLOW RATE (L) (m <sup>3</sup> /min)	ROOM AIR VOLUME (m <sup>3</sup> )	VOC CONC (mg/m <sup>3</sup> )	AIR EXCHANGE RATE n (h <sup>-1</sup> )	SHOWER RATE R (min <sup>-1</sup> )	I (hrs)	IMP/(%)	IMP/(%)	PER SHOWER VR (m <sup>3</sup> )	VR (m <sup>3</sup> )	SW (kg)	IMP/ADU SW	IN THE SHOWER ROOM	DA	IMP/ADU DA	DA-D <sub>0</sub>	M + M <sub>0</sub> (M <sub>0</sub> /M)
		Out (M <sub>0</sub> )	In (M <sub>0</sub> )																								
<b>VOLATILE ORGANIC COMPOUNDS</b>																											
<b>ADULT</b>																											
80-04-0	1,2-DICHLOROETHENE (80)	1.00E+00	2.00E+00	1.00	2.00	2.00E+01	6.00E+01	-4.07E-01	6.0000E-01	1.00E+00	10	0	0.00E+01	0.0000	12	10	4.00E-01	8.00E-01	2.00E-05	10	70	8.00E-01	1.194E+02	1.179E+02	-3	2.31E-05	
78-01-4	TRICHLOROETHENE (82)	1.00E+00	4.10E+00	1.00	2.00	3.04E+01	6.00E+01	-1.07E-01	6.0221E-01	2.70E+00	10	0	1.10E+02	0.0000	12	10	4.00E-01	8.00E-01	4.02E-05	10	70	8.00E-01	1.194E+02	1.179E+02	-3	4.60E-05	
<b>CHILD</b>																											
80-04-0	1,2-DICHLOROETHENE (80)	1.00E+00	2.00E+00	1.00	2.00	2.00E+01	6.00E+01	-4.07E-01	6.0000E-01	1.00E+00	10	0	0.00E+01	0.0000	12	10	4.00E-01	8.00E-01	1.70E-04	10	10	8.00E-01	1.194E+02	1.179E+02	-3	2.01E-04	
78-01-4	TRICHLOROETHENE (82)	1.00E+00	4.10E+00	1.00	2.00	3.04E+01	6.00E+01	-1.07E-01	6.0221E-01	2.70E+00	10	0	1.10E+02	0.0000	12	10	4.00E-01	8.00E-01	2.44E-04	10	10	8.00E-01	1.194E+02	1.179E+02	-3	2.94E-04	

SUBSURFACE SOIL INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 1)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL CHILD

Intake from ingestion of soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * CF * EF * ED * IR/BW * ATc \text{ or } ATnc * DY$$

$$\text{Risk} = \text{Intake} * CSF \text{ or } RfD$$

Where:  
 C = contaminant concentration in soil (mg/kg)  
 CF = conversion for kg to mg  
 EF = child exposure frequency (days/yr)  
 ED = child exposure duration (yr)  
 IR = child soil ingestion rate (mg/day)  
 BW = child body weight (kg)  
 ATc = averaging time for carcinogen (yr)  
 ATnc = averaging time for noncarcinogen (yr)  
 DY = days per year (days/year)  
 CSF = cancer slope factor (mg/kg-day)-1  
 RfD = reference dose (mg/kg-day)

INPUTS

specific  
specific

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Exposure Frequency (days/yr) Child	Exposure Duration (yr) Child	Conversion Factor (kg/mg)	Ingestion Rate (mg/day) Child	Body Weight (kg) Child	Average Carc Time (years)	Carc Dose (mg/kg/day)-1 Child	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc Time (years)	Noncarc Dose (mg/kg/day) Child	Reference Dose (mg/kg/day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child
Aluminum	4.6E+03	360	6	1E-06	200	15	25550	5.1E-03	0.00E+00	0.0E+00	0%	2190	6.9E-02	1.00E+00	5.8E-02	44%
Arsenic	1.2E+00	360	6	1E-06	200	15	25550	1.3E-06	1.75E+00	2.3E-06	100%	2190	1.6E-05	3.00E-04	6.2E-02	38%
Barium	9.1E+00	360	6	1E-06	200	15	25550	1.0E-06	0.00E+00	0.0E+00	0%	2190	1.2E-04	7.00E-02	1.7E-03	1%
Cadmium (soil)	3.2E-01	360	6	1E-06	200	15	25550	3.6E-07	0.00E+00	0.0E+00	0%	2190	4.1E-06	1.00E-03	4.1E-03	3%
Chromium	7.0E+00	360	6	1E-06	200	15	25550	7.7E-06	0.00E+00	0.0E+00	0%	2190	9.0E-05	1.00E+00	9.0E-05	0%
Cobalt	4.7E-01	360	6	1E-06	200	15	25550	6.1E-07	0.00E+00	0.0E+00	0%	2190	5.9E-06	6.00E-02	9.9E-05	0%
Copper	1.3E+00	360	6	1E-06	200	15	25550	1.4E-06	0.00E+00	0.0E+00	0%	2190	1.7E-05	3.71E-02	4.5E-04	0%
Lead	5.6E+00	360	6	1E-06	200	15	25550	6.1E-06	0.00E+00	0.0E+00	0%	2190	7.1E-06	0.00E+00	0.0E+00	0%
Manganese (soil)	6.6E+00	360	6	1E-06	200	15	25550	9.3E-06	0.00E+00	0.0E+00	0%	2190	1.1E-04	1.40E-01	7.8E-04	1%
Nickel	1.6E+00	360	6	1E-06	200	15	25550	1.6E-06	0.00E+00	0.0E+00	0%	2190	1.9E-05	2.00E-02	9.4E-04	1%
Vanadium	8.3E+00	360	6	1E-06	200	15	25550	9.1E-06	0.00E+00	0.0E+00	0%	2190	1.1E-04	7.00E-03	1.5E-02	11%
Zinc	5.4E+00	360	6	1E-06	200	15	25550	8.0E-06	0.00E+00	0.0E+00	0%	2190	4.9E-05	3.00E-01	2.3E-04	0%
Bis(2-ethylhexyl)phthalate	4.4E-01	360	6	1E-06	200	15	25550	4.8E-07	1.40E-02	6.7E-09	0%	2190	5.6E-06	2.00E-02	2.8E-04	0%
TOTAL								2.3E-06						1.3E-01		

SUBSURFACE SOIL DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 1)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL CHILD

Dermal contact with soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RfD$$

Where:

C = contaminant concentration in soil (mg/kg)  
 CF = conversion factor (kg/mg)  
 SA = child exposed skin surface area (cm<sup>2</sup>)  
 AF = soil to skin adherence factor (mg/cm<sup>2</sup>)  
 Abs = fraction absorbed (unitless)  
 EF = child exposure frequency (events/yr)  
 ED = child exposure duration (years)  
 BW = child body weight (kg)  
 ATc = averaging time for carcinogen (yr)  
 ATnc = averaging time for noncarcinogen (yr)  
 DY = day per year (day/yr)  
 CSF = cancer slope factor (mg/kg-day)-1  
 RfD = reference dose (mg/kg-day)

INPUTS

1E-06  
 2300  
 1  
 Specific  
 350  
 6  
 15  
 70  
 6  
 365  
 specific  
 specific

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> )	Averaging Factor (mg/cm <sup>2</sup> )	Fraction Absorbed (%)	Exposure Frequency (events/yr)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (years)	Carc Dose (mg/kg/day)	Dermal Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc Time (years)	Noncarc Dose (mg/kg/day)	Dermal Adjust Reference Dose (mg/kg-day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child
Aluminum	4.6E+03	1E-06	2300	1	0.001	350	6	15	25550	8.9E-05	0.0E+00	0%	2190	8.8E-04	2.0E-01	3.4E-03	44%	
Arsenic	1.2E+00	1E-06	2300	1	0.001	350	6	15	25550	1.6E-06	8.8E+00	1.3E-07	99%	2190	1.8E-07	6.0E-05	3.0E-03	38%
Barium	9.1E+00	1E-06	2300	1	0.001	350	6	15	25550	1.1E-07	0.0E+00	0.0E+00	0%	2190	1.3E-08	1.4E-02	9.6E-05	1%
Cadmium (soil)	3.2E-01	1E-06	2300	1	0.001	350	6	15	25550	4.0E-09	0.0E+00	0.0E+00	0%	2190	4.7E-08	2.0E-04	2.3E-04	3%
Chromium	7.0E+00	1E-06	2300	1	0.001	350	6	15	25550	8.8E-08	0.0E+00	0.0E+00	0%	2190	1.0E-08	2.0E-01	5.2E-08	0%
Dobalt	4.7E-01	1E-06	2300	1	0.001	350	6	15	25550	5.9E-09	0.0E+00	0.0E+00	0%	2190	6.8E-08	1.2E-02	6.7E-08	0%
Copper	1.3E+00	1E-06	2300	1	0.001	350	6	15	25550	1.6E-08	0.0E+00	0.0E+00	0%	2190	1.9E-07	7.4E-03	2.6E-05	0%
Lead	5.5E+00	1E-06	2300	1	0.001	350	6	15	25550	7.0E-08	0.0E+00	0.0E+00	0%	2190	8.1E-07	0.0E+00	0.0E+00	0%
Manganese (soil)	8.6E+00	1E-06	2300	1	0.001	350	6	15	25550	1.1E-07	0.0E+00	0.0E+00	0%	2190	1.2E-06	2.8E-02	4.6E-05	1%
Nickel	1.6E+00	1E-06	2300	1	0.001	350	6	15	25550	1.9E-08	0.0E+00	0.0E+00	0%	2190	2.2E-07	4.0E-03	5.4E-05	1%
Vanadium	8.3E+00	1E-06	2300	1	0.001	350	6	15	25550	1.0E-07	0.0E+00	0.0E+00	0%	2190	1.2E-06	1.4E-03	8.7E-04	11%
Zinc	5.4E+00	1E-06	2300	1	0.001	350	6	15	25550	6.8E-08	0.0E+00	0.0E+00	0%	2190	8.0E-07	6.0E-02	1.3E-05	0%
Bis(2-ethylhexyl)phthalate	4.4E-01	1E-06	2300	1	0.01	350	6	15	25550	5.5E-08	2.8E-02	1.5E-09	1%	2190	8.4E-07	1.0E-02	8.4E-05	1%
TOTAL										1.4E-07						7.8E-03		

SUBSURFACE SOIL PARTICULATE INHALATION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 1)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL CHILD

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (C * EF * ED * IR * 1/PEF) / (BW * ATc or ATnc * DY)$$

$$\text{Risk} = \text{Intake} * \text{CSF or RfD}$$

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	Calculated
CSF = carcinogenic slope factor	Specific
RfD = reference dose for noncarcinogen	Specific
IR = inhalation rate (m3)	10
EF = child exposure frequency (days)	350
ED = child exposure duration (years)	6
BW = child body weight (kg)	15
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	6
DY = day per year (day/yr)	365
PEF = particulate emission factor (m3/kg)	4.63E+09

Note: Inputs are scenario and site specific

COPC	Concentration Carcinogen (mg/kg)	Particulate Emission Factor (m3/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m3/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (years)	Carc Dose (mg/kg-day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncarc Time (years)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Aluminum	4.6E+03	4.6E+09	350	10	6	15	25550	6.6E-08	0.0E+00	0%	2190	6.6E-08	0.0E+00	0.0E+00	0%	
Arsenic	1.2E+00	4.6E+09	350	10	6	15	25550	1.4E-11	1.5E+01	2.2E-10	100%	2190	1.4E-11	0.0E+00	0.0E+00	0%
Barium	9.1E+00	4.6E+09	350	10	6	15	25550	1.1E-10	0.0E+00	0%	2190	1.1E-10	1.4E-04	7.5E-07	100%	
Cadmium (soil)	3.2E-01	4.6E+09	350	10	6	15	25550	3.8E-12	0.0E+00	0%	2190	3.8E-12	0.0E+00	0.0E+00	0%	
Chromium	7.0E+00	4.6E+09	350	10	6	15	25550	8.3E-11	0.0E+00	0%	2190	8.3E-11	0.0E+00	0.0E+00	0%	
Cobalt	4.7E-01	4.6E+09	350	10	6	15	25550	5.5E-12	0.0E+00	0%	2190	5.5E-12	0.0E+00	0.0E+00	0%	
Copper	1.3E+00	4.6E+09	350	10	6	15	25550	1.6E-11	0.0E+00	0%	2190	1.6E-11	0.0E+00	0.0E+00	0%	
Lead	5.5E+00	4.6E+09	350	10	6	15	25550	6.5E-11	0.0E+00	0%	2190	6.5E-11	0.0E+00	0.0E+00	0%	
Manganese (soil)	8.5E+00	4.6E+09	350	10	6	15	25550	1.0E-10	0.0E+00	0%	2190	1.0E-10	0.0E+00	0.0E+00	0%	
Nickel	1.6E+00	4.6E+09	350	10	6	15	25550	1.7E-11	0.0E+00	0%	2190	1.7E-11	0.0E+00	0.0E+00	0%	
Vanadium	8.3E+00	4.6E+09	350	10	6	15	25550	9.8E-11	0.0E+00	0%	2190	9.8E-11	0.0E+00	0.0E+00	0%	
Zinc	5.4E+00	4.6E+09	350	10	6	15	25550	6.4E-11	0.0E+00	0%	2190	6.4E-11	0.0E+00	0.0E+00	0%	
Bis(2-ethylhexyl)phthalate	4.4E-01	4.6E+09	350	10	6	15	25550	5.2E-12	0.0E+00	0%	2190	5.2E-12	0.0E+00	0.0E+00	0%	
<b>TOTAL</b>									2.2E-10						7.5E-07	

GROUNDWATER INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 1 - SHALLOW AND DEEP GROUNDWATER)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL ADULT

Intake from drinking water is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * IRw * EF * ED/BW * AT \text{ or } ATnc * DY$$

$$\text{Risk} = \text{Intake} * CSF \text{ or } RfD$$

Where:

	INPUTS
C = contaminant concentration in water (mg/l)	
IRw = adult daily water ingestion rate (L/Day)	2
EF = adult exposure frequency (days/yr)	350
ED = adult exposure duration (yr)	30
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	30
DY = days per year (day/year)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RfD = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

SHALLOW AND DEEP GROUNDWATER

COPC	Concentration (mg/l)	Ingestion Rate (L/day) Adult	Exposure Frequency (day/year) Adult	Exposure Duration (year) Adult	Body Weight (kg) Adult	Average Carc Time (years)	Carc Dose (mg/kg-day)-1 Adult	Slope Factor (mg/kg-day)-1 Adult	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (years)	Noncarc Dose (mg/kg-day) Adult	Reference Dose (mg/kg-day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Arsenic	8.5E-03	2	350	30	70	25550	1.0E-04	1.7E+00	1.7E-04	100%	10950	2.3E-04	3.00E-04	7.8E-01	10%
Barium	4.5E-02	2	350	30	70	25550	5.3E-04	0.00E+00	0.0E+00	0%	10950	1.2E-03	7.00E-02	1.8E-02	0%
Manganese (water)	1.2E+00	2	350	30	70	25550	1.4E-02	0.00E+00	0.0E+00	0%	10950	3.3E-02	5.00E-03	6.8E+00	88%
Mercury	1.0E-03	2	350	30	70	25550	1.2E-05	0.00E+00	0.0E+00	0%	10950	2.8E-05	3.00E-04	9.4E-02	1%
1,2-Dichloroethene	2.6E-03	2	350	30	70	25550	3.1E-05	0.00E+00	0.0E+00	0%	10950	7.1E-06	8.00E-03	7.9E-03	0%
Trichloroethene	4.1E-03	2	350	30	70	25550	4.8E-05	1.10E-02	5.3E-07	0%	10950	1.1E-04	6.00E-03	1.9E-02	0%
<b>TOTAL</b>								1.8E-04						7.8E+00	

GROUNDWATER DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 1 (SITE 1 - SHALLOW AND DEEP GROUNDWATER)  
 REMEDIAL INVESTIGATION OT-0201  
 MC3 CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL ADULT

Dermal Contact from groundwater is calculated as follows:

$$\text{Intake (mg/kg-day)} = \text{CW} * \text{SA} * \text{PC} * \text{ET} * \text{EF} * \text{ED} * \text{CF/BW} * \text{ATc or ATnc} * \text{DY}$$

Risk = Intake \* CSF or RfD

Where:

CW = contaminant concentration in water (mg/l)  
 SA = adult skin surface available for contact (cm<sup>2</sup>)  
 PC = contaminant specific dermal permeability (cm/h)  
 ET = adult exposure time (hours/day)  
 EF = adult exposure frequency (days/yr)  
 ED = adult exposure duration (years)  
 CF = volumetric conversion factor for water (liter/1000 cm<sup>3</sup>)  
 BW = adult body weight (kg)  
 ATc = averaging time for carcinogen (yr)  
 ATnc = averaging time for noncarcinogen (yr)  
 DY = days per year (days)

INPUTS

23000  
 Specific  
 0.25  
 360  
 30  
 0.001  
 70  
 70  
 30  
 365

Note: Inputs are site and scenario specific

SHALLOW AND DEEP GROUNDWATER

Contaminant	Concentration (mg/l)	Surface Area (cm <sup>2</sup> )	Dermal Permeability (cm/h)	Exposure Time (hours/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	Volumetric Conversion (l/mg)	Body Weight (kg)	Averaging Time (years)	Dose (mg/kg-day)	Dermal Adjust Slope Factor (mg/kg-day <sup>-1</sup> )	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc. Time (years)	Noncarc. Dose (mg/kg-day)	Dermal Adjust Reference Dose (mg/kg-day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Asbestos	1.0E-03	23000	1.00E-02	0.25	360	30	0.0001	70	26500	2.00E-07	2.00E-07	0%	0%	10000	3.5E-09	1.40E-02	2.5E-04	0%
Boron	4.0E-02	23000	1.00E-03	0.25	360	30	0.001	70	26500	1.00E-08	0.00E+00	0%	0%	10000	3.5E-09	1.40E-02	2.5E-04	0%
Manganese (water)	1.0E+00	23000	1.00E-03	0.25	360	30	0.001	70	26500	4.1E-05	0.00E+00	0%	0%	10000	9.5E-05	1.00E-03	9.5E-02	87%
Mercury	1.0E-03	23000	1.00E-02	0.25	360	30	0.001	70	26500	3.00E-08	0.00E+00	0%	0%	10000	8.1E-08	8.00E-06	1.4E-03	1%
2,0-Chloroethene	2.0E-03	23000	1.00E-02	0.25	360	30	0.001	70	26500	8.00E-07	0.00E+00	0%	0%	10000	2.0E-08	7.20E-03	2.5E-04	0%
Trichloroethane	4.1E-03	23000	1.00E-02	0.25	360	30	0.001	70	26500	2.2E-08	1.30E-02	3.0E-08	1%	10000	6.2E-08	4.80E-03	1.1E-03	1%
<b>TOTAL</b>																		

CTO-231 SITE 1 GROUNDWATER INHALATION  
01-Jun-95  
FILE: SHOW.WQ1

GROUNDWATER INHALATION - RME CASE

ADULT CHEMICAL	C <sub>rme</sub> mg/kg/shw	E <sub>D</sub> y	E <sub>F</sub> shwr/y	A <sub>T</sub> d	A <sub>TC</sub> d	C <sub>DI</sub> mg/kg/d	C <sub>DIC</sub> mg/kg/d	R <sub>FD</sub> mg/kg/d	P <sub>F</sub> (mg/kg/d) <sup>-1</sup>	H <sub>I</sub>	C <sub>R</sub>	% CONTRIB NC RISK	% CONTRIB CARC RISK
1,2-Dichloroethene	2.8E-05	30	350	10950	25550	2.7E-05	1.2E-05	0.0E+00	0.0E+00	---	0.0E+00	0%	0%
Trichloroethene	4.0E-05	30	350	10950	25550	3.9E-05	1.7E-05	0.0E+00	6.0E-03	---	9.9E-08	0%	100%
										TOTAL	0.0E+00	9.9E-08	

SHOWMOD.WQ1  
07-Jun-95

## SHOWER EXPOSURE MODEL

CAS No.	TARGET COMPOUND LIST CONSTITUTE	SHOWER EXPOSURE MODEL																
		OVERALL MASS TRANSF COEFFICIENT (M1) (cm/hr)	HENRY'S LAW CONSTAN (H) (atm-m3/mol-h)	GAS CONSTAN (F) (x10^-3 mol/h)	ABSOLUTE TEMP. (T1)	GAS CONST. ABS. TEMP. (RT)	GAS-film MASS TRANSF. COEFFICIENT (kg/cm^2 hr)	Liquid-film MASS TRANSF. COEFFICIENT (kg/cm^2 hr)	CARBON DIOXIDE TRANS. COEFF. (M-CO2) (cm/hr)	WATER TRANS. COEFF. (Vg-H2O) (cm/hr)	MOLECULAR WEIGHT (gram/mole)	RECIPROCAL OF (M)	(RT)/(H)*(kq))	OVERALL MASS TRANSF. COEFF. TEMP. COMP. (KAL) (cm/hr)	WATER VISCOSITY (U1) (cp)	WATER VISCOSITY (AT T1) (U1) (cp)	WATER TEMP. (T1) (K)	SHOWER' ((T1)*(U1))/((Tg)*(U1))
<b>/VOLATILE ORGANICS:</b>																		
	ADULT																	
540-59-0	1,2-DICHLOROETHENE (total)	1.30E+01	7.58E-03	8.20E-05	293.0	2.40E-02	1.29E+03	1.35E+01	20.0	3000.0	97.00	7.42E-02	2.45E-03	1.70E+01	0.596	1.002	318.0	5.48E-01
79-01-6	TRICHLOROETHENE (B2)	1.13E+01	9.10E-03	8.20E-05	293.0	2.40E-02	1.11E+03	1.16E+01	20.0	3000.0	131.40	8.64E-02	2.38E-03	1.52E+01	0.596	1.002	318.0	5.48E-01
	CHILD																	
540-59-0	1,2-DICHLOROETHENE (total)	1.30E+01	7.58E-03	8.20E-05	293.0	2.40E-02	1.29E+03	1.35E+01	20.0	3000.0	97.00	7.42E-02	2.45E-03	1.70E+01	0.596	1.002	318.0	5.48E-01
79-01-6	TRICHLOROETHENE (B2)	1.13E+01	9.10E-03	8.20E-05	293.0	2.40E-02	1.11E+03	1.16E+01	20.0	3000.0	131.40	8.64E-02	2.38E-03	1.52E+01	0.596	1.002	318.0	5.48E-01

SHOWMOOD  
07.JUN.98

CASE No.	TARGET COMPOUND/LENT CONSTITUENT	CONCENTRATION		SHOWER DURATION	SHOWER DROPLET DIA	SHOWER DROPLET DROP TIME	Kd, %	k <sub>d</sub> , %	-k <sub>d</sub> (t) /k <sub>d</sub> (0)	exp HOL, %	exp HOL, %	INDOOR VOC GENERATION RATE	SHOWER WATER FLOW RATE	SHOWER ROOM AIR VOLUME	INDOOR AIR EXCHANGE RATE	AIR CHg, %	SHOWER DURATION	TOTAL TIME DURATION	exp(D+I)	exp(I)	INHALATION EXPOSURE PER SHOWER	VENTILATION RATE	BODY WEIGHT	exp(RPD)	DURATION IN THE SHOWER ROOM	Di (hrs)	exp(D+I) (%)		(M) (M/H)/M <sub>0</sub>	
		Leaving Water Concen. (mg/l)	Leaving Water Concen. (mg/l)																											
VOLATILE ORGANIC	ADULT																													
60-090	1,2-DICHLOROETHENE (80%)	1.18E+00	2.40E+00	1.00	2.00	2.00E+01	6.00E+01	-6.97E-01	8.000E-01	1.20E+00	10	0	8.00E+01	0.0000	12	10	4.00E-01	0.00E-01	2.00E-05	10	70	6.00E-01	1.170E+02	1.170E+02	-4	3.01E-05				
70-014	TRICHLOROETHENE (82%)	1.09E+00	4.10E+00	1.00	2.00	2.00E+01	6.00E+01	-6.07E-01	0.0221E+00	0.72E+00	10	0	1.10E+02	0.0000	12	10	4.00E-01	0.00E-01	4.00E-05	10	70	6.00E-01	1.170E+02	1.170E+02	-3	4.00E-05				
CHILD	ADULT																													
60-090	1,2-DICHLOROETHENE (80%)	1.18E+00	2.40E+00	1.00	2.00	2.00E+01	6.00E+01	-6.97E-01	8.000E-01	1.20E+00	10	0	8.00E+01	0.0000	12	10	4.00E-01	0.00E-01	1.70E-05	10	15	1.00E+02	1.170E+02	1.170E+02	-3	2.01E-04				
70-014	TRICHLOROETHENE (82%)	1.09E+00	4.10E+00	1.00	2.00	2.00E+01	6.00E+01	-6.07E-01	0.0221E+00	0.72E+00	10	0	1.10E+02	0.0000	12	10	4.00E-01	0.00E-01	2.40E-04	10	15	1.00E+02	1.170E+02	1.170E+02	-3	4.00E-04				

SUBSURFACE SOIL INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 1)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL ADULT

Intake from ingestion of soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * CF * EF * ED * IR/BW * ATc \text{ or } ATnc * DY$$

$$\text{Risk} = \text{Intake} / \text{CSF or } RD$$

Where:

C = contaminant concentration in soil (mg/kg)  
 CF = conversion for kg to mg  
 EF = adult exposure frequency (days/yr)  
 ED = adult exposure duration (yr)  
 IR = adult soil ingestion rate (mg/day)  
 BW = adult body weight (kg)  
 ATc = averaging time for carcinogen (yr)  
 ATnc = averaging time for noncarcinogen (yr)  
 DY = days per year (days/year)  
 CSF = cancer slope factor (mg/kg-day)  
 RD = reference dose (mg/kg-day)

INPUTS

specific  
 specific

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Exposure Frequency (days/yr)	Exposure Duration (yr)	Conversion Factor (kg/mg)	Ingestion Rate (mg/day)	Body Weight (kg)	Average Carc Time (years)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (years)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Antimony	4.6E+03	360	30	1E-06	100	70	25550	2.7E-03	0.0E+00	0.0E+00	0%	10950	6.4E-03	1.0E+00	6.4E-03	44%
Arsenic	1.2E+00	360	30	1E-06	100	70	25550	7.1E-07	1.8E+00	1.2E-08	100%	10950	1.7E-08	3.0E-04	5.5E-03	38%
Barium	8.1E+00	360	30	1E-06	100	70	25550	5.4E-06	0.0E+00	0.0E+00	0%	10950	1.2E-06	7.0E-02	1.8E-04	1%
Cadmium (soil)	3.2E-01	360	30	1E-06	100	70	25550	1.9E-07	0.0E+00	0.0E+00	0%	10950	4.4E-07	1.0E-03	4.4E-04	3%
Chromium	7.0E+00	360	30	1E-06	100	70	25550	4.1E-06	0.0E+00	0.0E+00	0%	10950	9.6E-06	1.0E+00	9.6E-06	0%
Cobalt	4.7E-01	360	30	1E-06	100	70	25550	2.7E-07	0.0E+00	0.0E+00	0%	10950	6.4E-07	6.0E-02	1.1E-05	0%
Copper	1.3E+00	360	30	1E-06	100	70	25550	7.6E-07	0.0E+00	0.0E+00	0%	10950	1.8E-06	3.7E-02	4.8E-05	0%
Lead	6.6E+00	360	30	1E-06	100	70	25550	3.2E-06	0.0E+00	0.0E+00	0%	10950	7.6E-06	0.0E+00	0.0E+00	0%
Manganese (soil)	8.6E+00	360	30	1E-06	100	70	25550	5.0E-06	0.0E+00	0.0E+00	0%	10950	1.2E-06	1.4E-01	8.3E-05	1%
Nickel	1.5E+00	360	30	1E-06	100	70	25550	8.7E-07	0.0E+00	0.0E+00	0%	10950	2.0E-06	2.0E-02	1.0E-04	1%
Vanadium	9.3E+00	360	30	1E-06	100	70	25550	4.9E-06	0.0E+00	0.0E+00	0%	10950	1.1E-05	7.0E-03	1.8E-03	11%
Zinc	5.4E+00	360	30	1E-06	100	70	25550	3.2E-06	0.0E+00	0.0E+00	0%	10950	7.4E-06	3.0E-01	2.8E-05	0%
Bis(2-ethylhexyl)phthalate	4.4E-01	360	30	1E-06	100	70	25550	2.6E-07	1.4E-02	3.6E-09	0%	10950	8.0E-07	2.0E-02	3.0E-05	0%
TOTAL								1.2E-06							1.4E-02	

SUBSURFACE SOIL DERMA-CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 1)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL ADULT

Dermal contact with soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * CF * SA * AF * Abs * EF * ED/BW * ATc or ATnc * DY$$

Risk = Intake \* CSF or /RfD

Where:

C = contaminant concentration in soil (mg/kg)	1E-05
CF = conversion factor (kg/mg)	5800
SA = adult exposed skin surface area (cm <sup>2</sup> )	5800
AF = soil to skin adherence factor (mg/cm <sup>2</sup> )	1
Abs = fraction absorbed (units)	Specific
EF = adult exposure frequency (events/yr)	350
ED = adult exposure duration (years)	30
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	30
DY = day per year (day/yr)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RfD = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> ) Adult	Adherence Factor (mg/cm <sup>2</sup> )	Fraction Absorbed (%)	Exposure Frequency (events/yr) Adult	Exposure Duration (yrs) Adult	Body Weight (kg) Adult	Average Carc Time (years)	Carc Dose (mg/kg/day) Adult	Dermal Adjust. Slope Factor (mg/kg-day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (years)	Noncarc Dose (mg/kg/day) Adult	Dermal Adjust Reference Dose (mg/kg-day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	4.6E+03	1E-05	5800	1	0.001	350	30	70	25550	1.6E-04	0.0E+00	0%	10950	3.7E-04	2.0E-01	1.8E-03	44%	
Arsenic	1.2E+00	1E-05	5800	1	0.001	350	30	70	25550	4.1E-08	8.8E+00	98%	10950	9.8E-08	6.0E-06	1.6E-03	38%	
Barium	9.1E+00	1E-05	5800	1	0.001	350	30	70	25550	3.1E-07	0.0E+00	0%	10950	7.2E-07	1.4E-02	5.2E-05	1%	
Cadmium (soil)	3.2E-01	1E-05	5800	1	0.001	350	30	70	25550	1.1E-08	0.0E+00	0%	10950	2.5E-08	2.0E-04	1.3E-04	3%	
Chromium	7.0E+00	1E-05	5800	1	0.001	350	30	70	25550	2.4E-07	0.0E+00	0%	10950	5.6E-07	2.0E-01	2.8E-06	0%	
Cobalt	4.7E-01	1E-05	5800	1	0.001	350	30	70	25550	1.6E-08	0.0E+00	0%	10950	3.7E-08	1.2E-02	3.1E-05	0%	
Copper	1.3E+00	1E-05	5800	1	0.001	350	30	70	25550	4.4E-08	0.0E+00	0%	10950	1.0E-07	7.4E-03	1.4E-05	0%	
Lead	5.6E+00	1E-05	5800	1	0.001	350	30	70	25550	1.9E-07	0.0E+00	0%	10950	4.4E-07	0.0E+00	0.0E+00	0%	
Manganese (soil)	8.5E+00	1E-05	5800	1	0.001	350	30	70	25550	2.9E-07	0.0E+00	0%	10950	8.7E-07	2.8E-02	2.4E-05	1%	
Nickel	1.6E+00	1E-05	5800	1	0.001	350	30	70	25550	5.0E-08	0.0E+00	0%	10950	1.2E-07	4.0E-03	2.8E-05	1%	
Vanadium	8.3E+00	1E-05	5800	1	0.001	350	30	70	25550	2.8E-07	0.0E+00	0%	10950	6.8E-07	1.4E-03	4.7E-04	11%	
Zinc	5.4E+00	1E-05	5800	1	0.001	350	30	70	25550	1.8E-07	0.0E+00	0%	10950	4.3E-07	8.0E-02	7.2E-06	0%	
Bis(2-ethylhexyl)phthalate	4.4E-01	1E-05	5800	1	0.01	350	30	70	25550	1.6E-07	2.8E-02	4.2E-09	1%	10950	3.8E-07	1.0E-02	3.6E-06	1%
<b>TOTAL</b>										3.7E-07						4.2E-03		

## SUBSURFACE SOIL PARTICULATE INHALATION EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 1)

REMEDIAl INVESTIGATION CTO-0231

MCB CAMP LEJEUNE, NORTH CAROLINA

FUTURE RESIDENTIAL ADULT

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (C \times EF \times ED \times IR \times 1/\text{PEF}) / (\text{BW} \times ATc \text{ or } ATnc \times DY)$$

$$\text{Risk} = \text{Intake} \times \text{CSF} \text{ or } \text{RfD}$$

Where:

C = contaminant concentration in soil (mg/kg)

CSF = carcinogenic slope factor

RfD = reference dose for noncarcinogen

IR = Inhalation rate (m3)

## INPUTS

Calculated

Specific

Specific

20

350

30

70

70

30

365

4.63E+09

EF = adult exposure frequency (days)

ED = adult exposure duration (years)

BW = adult body weight (kg)

ATc = averaging time for carcinogen (yr)

ATnc = averaging time for noncarcinogen (yr)

DY = day per year (day/yr)

PEF = particulate emission factor (m3/kg)

TOTAL

COPC	Concentration Carcinogen (mg/kg)	Particulate Emission Factor (m3/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m3/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (years)	Carc Dose (mg/kg-day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncarc Time (years)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Aluminum	4.6E+03	4.6E+09	350	20	30	70	25550	1.2E-07	0.0E+00	0%	10950	2.9E-07	0.0E+00	0.0E+00	0%	
Arsenic	1.2E+00	4.6E+09	350	20	30	70	25550	3.1E-11	1.5E+01	4.6E-10	100%	10950	7.2E-11	0.0E+00	0.0E+00	0%
Barium	9.1E+00	4.6E+09	350	20	30	70	25550	2.3E-10	0.0E+00	0.0E+00	0%	10950	5.4E-10	1.4E-04	3.8E-06	100%
Cadmium (soil)	3.2E-01	4.6E+09	350	20	30	70	25550	9.1E-12	0.0E+00	0.0E+00	0%	10950	1.9E-11	0.0E+00	0.0E+00	0%
Chromium	7.0E+00	4.6E+09	350	20	30	70	25550	1.8E-10	0.0E+00	0.0E+00	0%	10950	4.2E-10	0.0E+00	0.0E+00	0%
Cobalt	4.7E-01	4.6E+09	350	20	30	70	25550	1.2E-11	0.0E+00	0.0E+00	0%	10950	2.8E-11	0.0E+00	0.0E+00	0%
Copper	1.3E+00	4.6E+09	350	20	30	70	25550	3.3E-11	0.0E+00	0.0E+00	0%	10950	7.7E-11	0.0E+00	0.0E+00	0%
Lead	6.6E+00	4.6E+09	350	20	30	70	25550	1.4E-10	0.0E+00	0.0E+00	0%	10950	3.3E-10	0.0E+00	0.0E+00	0%
Manganese (soil)	8.6E+00	4.6E+09	350	20	30	70	25550	2.2E-10	0.0E+00	0.0E+00	0%	10950	5.0E-10	0.0E+00	0.0E+00	0%
Nickel	1.6E+00	4.6E+09	350	20	30	70	25550	3.7E-11	0.0E+00	0.0E+00	0%	10950	8.7E-11	0.0E+00	0.0E+00	0%
Vanadium	8.3E+00	4.6E+09	350	20	30	70	25550	2.1E-10	0.0E+00	0.0E+00	0%	10950	4.9E-10	0.0E+00	0.0E+00	0%
Zinc	5.4E+00	4.6E+09	350	20	30	70	25550	1.4E-10	0.0E+00	0.0E+00	0%	10950	3.2E-10	0.0E+00	0.0E+00	0%
Bis(2-ethylhexyl)phthalate	4.4E-01	4.6E+09	350	20	30	70	25550	1.1E-11	0.0E+00	0.0E+00	0%	10950	2.6E-11	0.0E+00	0.0E+00	0%
<b>TOTAL</b>										4.6E-10					3.8E-06	

SURFACE SOIL INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 1)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT MILITARY PERSONNEL

Intake from ingestion of soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times EF \times IR/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RID$$

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	1E-06
CF = conversion for kg to mg	250
EF = adult exposure frequency (days/yr)	4
ED = adult exposure duration (yr)	100
IR = adult soil ingestion rate (mg/day)	70
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	4
DY = days per year (days/year)	365
CSF = cancer slope factor (mg/kg/day)-1	specific
RID = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Exposure Frequency (days/yr)	Exposure Duration (yr)	Conversion Factor (kg/mg)	Ingestion Rate (mg/day)	Body Weight (kg)	Average Carc Time (years)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (years)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	3.8E+03	250	4	1E-06	100	70	25550	2.1E-04	0.0E+00	0.0E+00	0%	1460	3.7E-03	1.00E+00	3.7E-03	15%
Antimony	6.2E+00	250	4	1E-06	100	70	25550	3.5E-07	0.0E+00	0.0E+00	0%	1460	6.1E-06	4.00E-04	1.5E-02	63%
Arsenic	1.1E+00	260	4	1E-08	100	70	25550	6.2E-08	1.75E+00	1.1E-07	100%	1460	1.1E-06	3.00E-04	3.6E-03	15%
Cadmium (soil)	7.1E-01	250	4	1E-06	100	70	25550	4.0E-09	0.0E+00	0.0E+00	0%	1460	6.9E-07	1.00E-03	6.9E-04	3%
Chromium	6.4E+00	250	4	1E-06	100	70	25550	3.6E-07	0.0E+00	0.0E+00	0%	1460	6.9E-06	1.00E+00	6.9E-06	0%
Manganese (soil)	1.0E+01	250	4	1E-06	100	70	25550	5.8E-07	0.0E+00	0.0E+00	0%	1460	1.0E-05	1.40E-01	7.2E-05	0%
Vanadium	6.2E+00	250	4	1E-06	100	70	25550	3.4E-07	0.0E+00	0.0E+00	0%	1460	6.0E-06	7.00E-03	8.6E-04	4%
Zinc	2.6E+01	250	4	1E-06	100	70	25550	1.6E-06	0.0E+00	0.0E+00	0%	1460	2.6E-05	3.00E-01	8.5E-05	0%
6,4'-DDE	2.6E-03	250	4	1E-06	100	70	25550	1.6E-10	3.40E-01	5.0E-11	0%	1460	2.6E-09	0.00E+00	0.0E+00	0%
1,4'-DDT	5.0E-03	250	4	1E-06	100	70	25550	2.8E-10	3.40E-01	9.4E-11	0%	1460	4.9E-09	5.00E-04	9.7E-06	0%
TOTAL									1.1E-07						2.4E-02	

SURFACE SOIL DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 1)  
 REMEDIAL INVESTIGATION CTC-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT MILITARY PERSONNEL

Dermal contact with soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RID$$

Where:

C = contaminant concentration in soil (mg/kg)

INPUTS

CF	1E-08
SA	4300
AF	1
Abs	Fraction absorbed (unitless)
EF	250
ED	adult exposure duration (years)
BW	adult body weight (kg)
ATc	averaging time for carcinogen (yr)
ATnc	averaging time for noncarcinogen (yr)
DY	day per year (day/yr)
CSF	cancer slope factor (mg/kg-day)-1
RID	reference dose (mg/kg-day)

Specific	
Specific	
specific	
specific	

Note: Inputs are scenario and site specific

CP/C	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> )	Adherence Factor (mg/cm <sup>2</sup> )	Fraction Absorbed (%)	Exposure Frequency (events/yr)	Exposure Duration (ms)	Body Weight (kg)	Average Carc Time (years)	Carc Dose (mg/kg-day)	Dermal Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (years)	Noncarc Dose (mg/kg/day)	Dermal Adjust Reference Dose (mg/kg-day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	3.8E+03	1E-08	4300	1	0.001	250	4	70	25550	9.2E-08	0.00E+00	0.0E+00	0%	1460	1.8E-04	2.00E-01	1.0E-04	15%
Antimony	6.2E+00	1E-08	4300	1	0.001	250	4	70	25550	1.8E-08	0.00E+00	0.0E+00	0%	1460	2.6E-07	8.00E-05	3.3E-03	63%
Arsenic	1.1E+00	1E-08	4300	1	0.001	250	4	70	25550	2.7E-09	0.75E+00	2.3E-08	99%	1460	4.7E-08	6.00E-05	7.8E-04	15%
Cadmium (soil)	7.1E-01	1E-08	4300	1	0.001	250	4	70	25550	1.7E-09	0.00E+00	0.0E+00	0%	1460	3.0E-08	2.00E-04	1.5E-04	3%
Chromium	6.4E+00	1E-08	4300	1	0.001	250	4	70	25550	1.8E-08	0.00E+00	0.0E+00	0%	1460	2.7E-07	2.00E-01	1.3E-08	0%
Manganese (soil)	1.0E+01	1E-08	4300	1	0.001	250	4	70	25550	2.6E-08	0.00E+00	0.0E+00	0%	1460	4.4E-07	2.00E-02	1.8E-05	0%
Vanadium	6.2E+00	1E-08	4300	1	0.001	250	4	70	25550	1.6E-08	0.00E+00	0.0E+00	0%	1460	2.6E-07	1.40E-03	1.8E-04	4%
Zinc	2.8E+01	1E-08	4300	1	0.001	250	4	70	25550	8.8E-09	0.00E+00	0.0E+00	0%	1460	1.1E-08	6.00E-02	1.8E-05	0%
U-DOE	2.6E-03	1E-08	4300	1	0.01	250	4	70	25550	8.0E-11	6.80E-01	4.3E-11	0%	1460	1.1E-08	0.00E+00	0.0E+00	0%
U-DOT	6.0E-03	1E-08	4300	1	0.01	250	4	70	25550	1.2E-10	6.80E-01	8.1E-11	0%	1460	2.1E-08	2.80E-04	8.4E-08	0%
TOTAL										2.3E-08							5.2E-03	

## SURFACE SOIL PARTICULATE INHALATION EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 1)

REMEDIAl INVESTIGATION CTO-0231

MCB CAMP LEJEUNE, NORTH CAROLINA

CURRENT MILITARY PERSONNEL

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (\text{C} * \text{EF} * \text{ED} * \text{IR} * 1/\text{PEF}) / (\text{BW} * \text{ATc or ATnc} * \text{DY})$$

$$\text{Risk} = \text{Intake} * \text{CSF or RID}$$

Where:

C = contaminant concentration in soil (mg/kg)  
 CSF = carcinogenic slope factor  
 RID = reference dose for noncarcinogen  
 IR = inhalation rate (m³)  
 30  
 EF = adult exposure frequency (days)  
 250  
 ED = adult exposure duration (years)  
 4  
 BW = adult body weight (kg)  
 70  
 ATc = averaging time for carcinogen (yr)  
 70  
 ATnc = averaging time for noncarcinogen (yr)  
 4  
 DY = day per year (day/yr)  
 365  
 PEF = particulate emission factor (m³/kg)  
 4.63E+09

INPUTS  
 Calculated  
 Specific  
 Specific  
 Specific  
 30  
 250  
 4  
 70  
 70  
 4  
 365  
 4.63E+09

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Particulate Emission Factor (m³/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m³/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (years)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncarc Time (years)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Aluminum	3.8E+03	4.6E+09	250	30	4	70	25550	1.4E-08	0.00E+00	0%	1460	2.4E-07	0.00E+00	0.0E+00	0%	
Antimony	6.2E+00	4.6E+09	250	30	4	70	25550	2.3E-11	0.00E+00	0%	1460	4.0E-10	0.00E+00	0.0E+00	0%	
Arsenic	1.1E+00	4.6E+09	250	30	4	70	25550	4.0E-12	1.51E+01	6.1E-11	100%	1460	7.0E-11	0.00E+00	0.0E+00	0%
Cadmium (soil)	7.1E-01	4.6E+09	250	30	4	70	25550	2.6E-12	0.00E+00	0%	1460	4.5E-11	0.00E+00	0.0E+00	0%	
Chromium	6.4E+00	4.6E+09	250	30	4	70	25550	2.3E-11	0.00E+00	0%	1460	4.1E-10	0.00E+00	0.0E+00	0%	
Manganese (soil)	1.0E+01	4.6E+09	250	30	4	70	25550	3.7E-11	0.00E+00	0%	1460	6.6E-10	0.00E+00	0.0E+00	0%	
Titanium	6.2E+00	4.6E+09	250	30	4	70	25550	2.2E-11	0.00E+00	0%	1460	3.9E-10	0.00E+00	0.0E+00	0%	
Zinc	2.6E+01	4.6E+09	250	30	4	70	25550	9.5E-11	0.00E+00	0%	1460	1.7E-09	0.00E+00	0.0E+00	0%	
1,4-DDE	2.6E-03	4.6E+09	250	30	4	70	25550	9.6E-15	0.00E+00	0%	1460	1.7E-13	0.00E+00	0.0E+00	0%	
1,4-DDT	5.0E-03	4.6E+09	250	30	4	70	25550	1.8E-14	3.40E-01	6.1E-15	0%	1460	3.2E-13	0.00E+00	0.0E+00	0%
<b>TOTAL</b>									6.1E-11							0.0E+00

SUBSURFACE SOIL INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 1)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE CONSTRUCTION WORKER

Intake from ingestion of soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * CF * EF * ED * IR/BW * ATc \text{ or } ATnc * DY$$

Risk = Intake \* CSF or /RfD

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	1E-08
CF = conversion for kg to mg	90
EF = adult exposure frequency (days/yr)	1
ED = adult exposure duration (yr)	480
IR = adult soil ingestion rate (mg/day)	70
BW = adult body weight (kg)	1
ATc = averaging time for carcinogen (yr)	365
ATnc = averaging time for noncarcinogen (yr)	70
DY = days per year (days/year)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RfD = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Exposure Frequency (days/yr) Adult	Exposure Duration (yr) Adult	Conversion Factor (kg/mg)	Ingestion Rate (mg/day) Adult	Body Weight (kg) Adult	Average Carc Time (years)	Carc Dose (mg/kg/day)-1 Adult	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (years)	Noncarc Dose (mg/kg/day) Adult	Heterogeneity Dose (mg/kg/day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	4.6E+03	90	1	1E-08	480	70	25550	1.1E-04	0.00E+00	0.0E+00	0%	365	1.1E-04	1.00E+00	1.1E-04	44%
Arsenic	1.2E+00	90	1	1E-08	480	70	25550	2.9E-08	1.75E+00	5.1E-08	100%	365	2.9E-08	3.00E-04	9.8E-05	38%
Barium	9.1E+00	90	1	1E-08	480	70	25550	2.2E-07	0.00E+00	0.0E+00	0%	365	2.2E-07	7.00E-02	3.1E-06	1%
Cadmium (soil)	3.2E-01	90	1	1E-08	480	70	25550	7.7E-09	0.00E+00	0.0E+00	0%	365	7.7E-09	1.00E-03	7.7E-06	3%
Chromium	7.0E+00	90	1	1E-08	480	70	25550	1.7E-07	0.00E+00	0.0E+00	0%	365	1.7E-07	1.00E+00	1.7E-07	0%
Cobalt	4.7E-01	90	1	1E-08	480	70	25550	1.1E-06	0.00E+00	0.0E+00	0%	365	1.1E-06	6.00E-02	1.9E-07	0%
Copper	1.3E+00	90	1	1E-08	480	70	25550	3.1E-08	0.00E+00	0.0E+00	0%	365	3.1E-08	3.71E-02	8.8E-07	0%
Lead	6.5E+00	90	1	1E-08	480	70	25550	1.3E-07	0.00E+00	0.0E+00	0%	365	1.3E-07	0.00E+00	0.0E+00	0%
Manganese (soil)	8.6E+00	90	1	1E-08	480	70	25550	2.1E-07	0.00E+00	0.0E+00	0%	365	2.1E-07	1.40E-01	1.8E-06	1%
Nickel	1.5E+00	90	1	1E-08	480	70	25550	3.6E-08	0.00E+00	0.0E+00	0%	365	3.6E-08	2.00E-02	1.8E-06	1%
Vanadium	8.3E+00	90	1	1E-08	480	70	25550	2.0E-07	0.00E+00	0.0E+00	0%	365	2.0E-07	7.00E-03	2.9E-05	11%
Zinc	5.4E+00	90	1	1E-08	480	70	25550	1.3E-07	0.00E+00	0.0E+00	0%	365	1.3E-07	3.00E-01	4.4E-07	0%
Bis(2-ethylhexyl)phthalate	4.4E-01	90	1	1E-08	480	70	25550	1.1E-08	1.40E-02	1.6E-10	0%	365	1.1E-08	2.00E-02	5.3E-07	0%
<b>TOTAL</b>									<b>6.1E-08</b>						<b>2.5E-04</b>	

SUBSURFACE SOIL DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 1)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCS CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE CONSTRUCTION WORKER

Dermal contact with soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RfD$$

Where:  
 C = contaminant concentration in soil (mg/kg)

INPUTS

CF = conversion factor (kg/mg)	1E-06
SA = adult exposed skin surface area (cm <sup>2</sup> )	4300
AF = soil to skin adherence factor (mg/cm <sup>2</sup> )	1
Abs = fraction absorbed (unitless)	Specific
EF = adult exposure frequency (events/yr)	90
ED = adult exposure duration (years)	1
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	1
DY = day per year (day/yr)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RfD = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

POPC	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> ) Adult	Adherence Factor (mg/cm <sup>2</sup> )	Fraction Absorbed (%)	Exposure Frequency (events/yr) Adult	Exposure Duration (yrs) Adult	Body Weight (kg) Adult	Average Carc Time (years) Adult	Carc Dose (mg/kg/day) Adult	Dermal Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (years)	Noncarc Dose (mg/kg/day) Adult	Dermal Adjust Reference Dose (mg/kg-day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	4.8E+03	1E-06	4300	1	0.001	90	1	70	25550	1.0E-06	0.00E+00	0%	365	7.0E-06	2.00E-01	3.6E-04	44%	
Arsenic	1.2E+00	1E-06	4300	1	0.001	90	1	70	25550	2.6E-10	8.75E+00	2.3E-09	99%	365	1.8E-06	6.00E-05	3.1E-04	38%
Barium	9.1E+00	1E-06	4300	1	0.001	90	1	70	25550	2.0E-09	0.00E+00	0.0E+00	0%	365	1.4E-07	1.40E-02	9.9E-06	1%
Cadmium (soil)	3.2E-01	1E-06	4300	1	0.001	90	1	70	25550	8.9E-11	0.00E+00	0.0E+00	0%	365	4.8E-09	2.00E-04	2.4E-05	3%
Chromium	7.0E+00	1E-06	4300	1	0.001	90	1	70	25550	1.6E-09	0.00E+00	0.0E+00	0%	365	1.1E-07	2.00E-01	5.3E-07	0%
Cobalt	4.7E-01	1E-06	4300	1	0.001	90	1	70	25550	1.0E-10	0.00E+00	0.0E+00	0%	365	7.0E-09	1.20E-02	5.9E-07	0%
Copper	1.3E+00	1E-06	4300	1	0.001	90	1	70	25550	2.6E-10	0.00E+00	0.0E+00	0%	365	2.0E-06	7.42E-03	2.7E-06	0%
Lead	5.6E+00	1E-06	4300	1	0.001	90	1	70	25550	1.2E-09	0.00E+00	0.0E+00	0%	365	8.4E-06	0.00E+00	0.0E+00	0%
Manganese (soil)	8.6E+00	1E-06	4300	1	0.001	90	1	70	25550	1.8E-09	0.00E+00	0.0E+00	0%	365	1.3E-07	2.80E-02	4.6E-06	1%
Nickel	1.5E+00	1E-06	4300	1	0.001	90	1	70	25550	3.2E-10	0.00E+00	0.0E+00	0%	365	2.2E-06	4.00E-03	5.6E-06	1%
Vanadium	8.3E+00	1E-06	4300	1	0.001	90	1	70	25550	1.8E-09	0.00E+00	0.0E+00	0%	365	1.3E-07	1.40E-03	9.0E-05	11%
Zinc	8.4E+00	1E-06	4300	1	0.001	90	1	70	25550	1.2E-09	0.00E+00	0.0E+00	0%	365	8.2E-06	6.00E-02	1.4E-06	0%
Bis(2-ethylhexyl)phthalate	4.4E-01	1E-06	4300	1	0.01	90	1	70	25550	9.4E-10	2.80E-02	2.6E-11	1%	365	8.6E-06	1.00E-02	6.6E-06	1%
<b>TOTAL</b>											2.0E-09						8.0E-04	

## SUBSURFACE SOIL PARTICULATE INHALATION EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 1)

REMEDIATION INVESTIGATION CTO-0231

MCB CAMP LEJEUNE, NORTH CAROLINA

FUTURE CONSTRUCTION WORKER

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (\text{C} * \text{EF} * \text{ED} * \text{IR} * 1/\text{PEF}) / (\text{BW} * \text{ATc} * \text{ATnc} * \text{DY})$$

Risk = Intake \* CSF or /RfD

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	Calculated
CSF = carcinogenic slope factor	Specific
RfD = reference dose for noncarcinogen	Specific
IR = inhalation rate (m/s)	20
EF = adult exposure frequency (days)	90
ED = adult exposure duration (years)	1
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	1
DY = day per year (day/yr)	365
PEF = particulate emission factor (m³/kg)	4.63E+09

Note: Inputs are scenario and site specific

OPC	Concentration Carcinogen (mg/kg)	Particulate Emission Factor (m³/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m³/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (years)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncarc Time (years)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Manganese	4.6E+03	4.6E+09	90	20	1	70	25550	1.0E-03	0.0E+00	0.0E+00	0%	365	7.1E-08	0.0E+00	0.0E+00	0%
Arsenic	1.2E+00	4.6E+09	90	20	1	70	25550	2.6E-13	1.6E+01	4.0E-12	100%	365	1.8E-11	0.0E+00	0.0E+00	0%
Barium	9.1E+00	4.6E+09	90	20	1	70	25550	2.0E-12	0.0E+00	0.0E+00	0%	365	1.4E-10	1.4E-04	0.7E-07	100%
Cadmium (soil)	3.2E-01	4.6E+09	90	20	1	70	25550	6.9E-14	0.0E+00	0.0E+00	0%	365	4.9E-12	0.0E+00	0.0E+00	0%
Chromium	7.0E+00	4.6E+09	90	20	1	70	25550	1.6E-12	0.0E+00	0.0E+00	0%	365	1.1E-10	0.0E+00	0.0E+00	0%
Cobalt	4.7E-01	4.6E+09	90	20	1	70	25550	1.0E-13	0.0E+00	0.0E+00	0%	365	7.1E-12	0.0E+00	0.0E+00	0%
Copper	1.3E+00	4.6E+09	90	20	1	70	25550	2.8E-13	0.0E+00	0.0E+00	0%	365	2.0E-11	0.0E+00	0.0E+00	0%
Lead	6.5E+00	4.6E+09	90	20	1	70	25550	1.2E-12	0.0E+00	0.0E+00	0%	365	8.4E-11	0.0E+00	0.0E+00	0%
Manganese (soil)	8.6E+00	4.6E+09	90	20	1	70	25550	1.8E-12	0.0E+00	0.0E+00	0%	365	1.3E-10	0.0E+00	0.0E+00	0%
Nickel	1.6E+00	4.6E+09	90	20	1	70	25550	3.2E-13	0.0E+00	0.0E+00	0%	365	2.2E-11	0.0E+00	0.0E+00	0%
Vanadium	8.3E+00	4.6E+09	90	20	1	70	25550	1.8E-12	0.0E+00	0.0E+00	0%	365	1.3E-10	0.0E+00	0.0E+00	0%
Zinc	6.4E+00	4.6E+09	90	20	1	70	25550	1.2E-12	0.0E+00	0.0E+00	0%	365	8.3E-11	0.0E+00	0.0E+00	0%
Di(2-ethylhexyl)phthalate	4.4E-01	4.6E+09	90	20	1	70	25550	9.6E-14	0.0E+00	0.0E+00	0%	365	6.6E-12	0.0E+00	0.0E+00	0%
TOTAL								4.0E-12								9.7E-07

**SITE 28**

---

GROUNDWATER INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 28)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL CHILD

Intake from drinking water is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * IRw * EF * ED/BW * AT or ATnc * DY$$

$$\text{Risk} = \text{Intake} * \text{CSF or } / \text{RfD}$$

Where:

	INPUTS
C = contaminant concentration in water (mg/l)	
IRw = child daily water ingestion rate (L/Day)	1
EF = child exposure frequency (days/yr)	350
ED = child exposure duration (yr)	6
BW = child body weight (kg)	15
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	6
DY = days per year (day/year)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RfD = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

SHALLOW AND DEEP GROUNDWATER

CP/C	Concentration Carcinogen (mg/l)	Ingestion Rate (L/day) Child	Exposure Frequency (day/year) Child	Exposure Duration (year) Child	Body Weight (kg) Child	Average Carc Time (days)	Carc Dose (mg/kg-day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc Time (days)	Noncarc Dose (mg/kg-day) Child	Reference Dose (mg/kg-day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child
Arsenic	2.7E-03	1	350	6	15	25550	1.5E-05	1.8E+00	2.8E-05	63%	2190	1.7E-04	3.0E-04	5.8E-01	3%
Barium	7.8E-01	1	350	6	15	25550	4.2E-03	0.0E+00	0.0E+00	0%	2190	4.9E-02	7.0E-02	6.9E-01	3%
Lead	3.7E-02	1	350	6	15	25550	2.0E-04	0.0E+00	0.0E+00	0%	2190	2.4E-03	0.0E+00	0.0E+00	0%
Manganese (water)	1.5E+00	1	350	6	15	25550	7.9E-03	0.0E+00	0.0E+00	0%	2190	9.3E-02	5.0E-03	1.9E+01	92%
Mercury	5.0E-04	1	350	6	15	25550	2.7E-06	0.0E+00	0.0E+00	0%	2190	3.2E-05	3.0E-04	1.1E-01	1%
2,4-Dimethylphenol	4.3E-03	1	350	6	15	25550	2.4E-05	0.0E+00	0.0E+00	0%	2190	2.7E-04	2.0E-02	1.4E-02	0%
Acenaphthene	1.0E-02	1	350	6	15	25550	5.5E-05	0.0E+00	0.0E+00	0%	2190	6.4E-04	6.0E-02	1.1E-02	0%
4,4'-DDE	1.6E-03	1	350	6	15	25550	8.8E-06	3.4E-01	3.0E-06	7%	2190	1.0E-04	0.0E+00	0.0E+00	0%
4,4'-DDD	9.0E-03	1	350	6	15	25550	4.9E-05	2.4E-01	1.2E-05	29%	2190	6.8E-04	0.0E+00	0.0E+00	0%
4,4'-DDT	1.8E-04	1	350	6	15	25550	9.8E-07	3.4E-01	3.4E-07	1%	2190	1.2E-05	5.0E-04	2.3E-02	0%
Chloroform	1.2E-03	1	350	6	15	25550	6.6E-06	6.1E-03	4.0E-06	0%	2190	7.7E-05	1.0E-02	7.7E-03	0%
2-Methylnaphthalene	9.3E-03	1	350	6	15	25550	5.1E-05	0.0E+00	0.0E+00	0%	2190	5.9E-04	4.0E-02	1.5E-02	0%
Phenanthrene	8.6E-03	1	350	6	15	25550	3.6E-05	0.0E+00	0.0E+00	0%	2190	4.2E-04	3.0E-02	1.4E-02	0%
4-Methylphenol	9.3E-03	1	350	6	15	25550	5.1E-05	0.0E+00	0.0E+00	0%	2190	5.9E-04	5.0E-03	1.2E-01	1%
TOTAL								4.1E-05						2.0E+01	

GROUNDWATER DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCBS CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL CHILD

Dermal Contact from groundwater is calculated as follows:

$$\text{Intake (mg/kg-day)} = \text{CW} * \text{SA} * \text{PC} * \text{ET} * \text{EF} * \text{ED} * \text{CF/BW} * \text{ATc or ATnc} * \text{DY}$$

Risk = Intake \* CSF or RfD

Where:

CW = contaminant concentration in water (mg/l)

SA = child skin surface available for contact (cm<sup>2</sup>)

INPUTS

Specific

10000

0.26

350

6

0.001

15

70

6

365

PC = contaminant specific dermal permeability (cm/h)

ET = child exposure time (hours/day)

EF = child exposure frequency (days/yr)

ED = child exposure duration (years)

CF = volumetric conversion factor for water (liter/1000 cm<sup>3</sup>)

BW = child body weight (kg)

ATc = averaging time for carcinogen (yr)

ATnc = averaging time for noncarcinogen (yr)

DY = days per year (days)

Note: Inputs are site and scenario specific

SHALLOW AND DEEP GROUNDWATER

CP/C	Concentration Carcinogen (mg/l)	Surfaces Area (cm <sup>2</sup> ) Child	Dermal Permeability (cm/h)	Exposure Time (hour/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	Volumetric Conversion (L/m <sup>3</sup> )	body Weight (kg) Child	Averaging Carc Time (days)	Carc Dose (mg/kg-day) Child	Dermal Adjust. Slope Factor (mg/kg-day) <sup>-1</sup>	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc Time (days)	Noncarc Dose (mg/kg-day) Child	Dermal Adjust. Reference Dose (mg/kg-day)	Noncarc Risk Child	Percent Noncarcogenic Risk Child	
Asinine	2.7E-03	10000	1.0E-03	0.25	350	6	0.001	15	25500	8.0E-07	3.0E-07	2E-06	2E-06	2190	1.3E-07	6.0E-05	7.0E-03	2%	
Barium	7.0E-01	10000	1.0E-03	0.25	350	6	0.001	15	25500	1.0E-05	0.0E+00	0%	2190	1.2E-04	1.4E-02	8.7E-03	2%		
Lead	3.7E-02	10000	4.0E-06	0.25	350	6	0.001	15	25500	2.0E-09	0.0E+00	0%	2190	2.4E-08	0.0E+00	0.0E+00	0%		
Manganese (water)	1.5E-01	10000	1.0E-03	0.25	350	6	0.001	15	25500	2.0E-05	0.0E+00	0%	2190	2.3E-04	1.0E-03	2.3E-01	67%		
Mercury	5.0E-04	10000	1.0E-03	0.25	350	6	0.001	15	25500	6.0E-09	0.0E+00	0%	2190	8.0E-08	6.0E-05	1.3E-03	0%		
2,4-Dimethylphenol	4.0E-03	10000	1.0E-02	0.25	350	6	0.001	15	25500	8.0E-07	0.0E+00	0%	2190	1.0E-05	1.0E-03	1.0E-03	0%		
Naphthalene	1.0E-02	10000	1.0E-01	0.25	350	6	0.001	15	25500	2.1E-05	0.0E+00	0%	2190	2.4E-04	3.0E-03	8.1E-03	2%		
o-XOe	1.0E-03	10000	2.4E-01	0.25	350	6	0.001	15	25500	6.3E-08	0.0E+01	1%	2190	8.1E-05	0.0E+00	0.0E+00	0%		
p-XOD	9.0E-03	10000	2.0E-01	0.25	350	6	0.001	15	25500	3.6E-05	4.0E-01	1.7E-05	7.0E-04	2190	4.0E-04	0.0E+00	0.0E+00	0%	
p-XOT	1.0E-04	10000	4.3E-01	0.25	350	6	0.001	15	25500	1.1E-06	6.0E-01	7.2E-07	3%	2190	1.2E-05	2.0E-04	4.0E-02	14%	
Chromate	1.2E-03	10000	8.0E-03	0.25	350	6	0.001	15	25500	1.5E-07	7.0E-03	1.1E-09	0%	2190	1.7E-03	8.0E-03	2.1E-04	0%	
p-Methylnaphthalene	9.0E-03	10000	1.4E-01	0.25	350	6	0.001	15	25500	1.8E-05	0.0E+00	0%	2190	2.1E-04	2.0E-02	1.1E-02	3%		
Phenanthrene	8.0E-03	10000	2.7E-01	0.25	350	6	0.001	15	25500	2.4E-05	0.0E+00	0%	2190	2.8E-04	1.6E-02	1.0E-02	5%		
t-Methylphenol	9.0E-03	10000	1.8E-02	0.25	350	6	0.001	15	25500	2.3E-06	0.0E+00	0%	2190	2.7E-05	2.0E-02	1.1E-02	3%		
TOTAL											2.0E-05					3.0E-01			

CTO-231 SITE 28 GROUNDWATER INHALATION  
 13-Jun-95  
 FILE: SHOW.WQ1

GROUNDWATER INHALATION - RME CASE

CHEMICAL	C rme	ED	EF	AT	ATC	CDI	CDIC	RFD	PF	HI	CR	% CONTRIB	% CONTRIB
	mg/kg/shw	y	shwr/y	d	d	mg/kg/d	mg/kg/d	mg/kg/d	(mg/kg/d) <sup>-1</sup>			NC RISK	CARC RISK
Chloroform	7.09E-05	9	350	3285	25550	6.8E-05	8.7E-06	0.0E+00	8.1E-02	---	7.0E-07	0%	88%
Phenanthrene	1.57E-04	9	350	3285	25550	1.5E-04	1.9E-05	0.0E+00	0.0E+00	---	0.0E+00	0%	0%
2, 4-Dimethylphenol	8.71E-07	9	350	3285	25550	8.4E-07	1.1E-07	0.0E+00	0.0E+00	---	0.0E+00	0%	0%
Acenaphthene	1.81E-04	9	350	3285	25550	1.7E-04	2.2E-05	0.0E+00	0.0E+00	---	0.0E+00	0%	0%
4, 4'-DDE	2.06E-05	9	350	3285	25550	2.0E-05	2.5E-06	0.0E+00	0.0E+00	---	0.0E+00	0%	0%
4, 4'-DDT	2.20E-06	9	350	3285	25550	2.1E-06	2.7E-07	0.0E+00	3.4E-01	---	9.2E-08	0%	12%
4-Methylphenol	3.05E-06	9	350	3285	25550	2.9E-06	3.8E-07	0.0E+00	0.0E+00	---	0.0E+00	0%	0%
									TOTAL		0.0E+00	8.0E-07	

site 28 groundwater  
13-Jun-96 file: showmod.wq1

CAS No.	TARGET COMPOUND LIST CONSTITUE	OVERALL MASS TRANSF COEFFICIENT	HENRY'S LAW CONSTAN	GAS CONSTAN	ABSOLUTE TEMP.	GAS CONST.	GAS-FLM MASS TRANSF.	LIQUID-FLM MASS TRANSF.	CARBON DIOXIDE COEFFICIENT	WATER TRANS. COEFF.	MOLECULAR WEIGHT	RECIPROCAL OF (RT)/(H)(kg)	OVERALL MASS TRANSF. COEFF.	WATER VISCOSITY	WATER VISCOSITY	SHOWER ((T1)*(U1))/((T2)*(U2))		
		(KL) (cm/hr)	(H) (atm-m3/mol-K)	(R) (atm-m3/mol-K)	(K)	(RT)	(kg) (cm/mr)	(cm/mr)	(kg) (cm/hr)	(kg-CO2) (cm/hr)	(gram/mole)	(K) (cal)	(AT T1) (Us) (cp)	(AT T1) (U1) (cp)	(T2) (K)			
<b>site 28 RME adult</b>																		
	Chloroform	1.12E+01	2.87E-03	8.20E-05	293.0	2.40E-02	1.16E+03	1.21E+01	20.0	3000.0	119.40	8.24E-02	7.19E-03	1.51E+01	0.596	1.002	318.0	5.48E-01
	Phenanthrene	3.88E+00	1.59E-04	8.20E-05	293.0	2.40E-02	9.53E+02	9.94E+00	20.0	3000.0	178.22	1.01E-01	1.58E-01	6.21E+00	0.596	1.002	318.0	5.48E-01
	2,4-dimethylphenol	3.01E-02	6.30E-07	8.20E-05	293.0	2.40E-02	1.15E+03	1.20E+01	20.0	3000.0	122.16	8.33E-02	3.31E+01	4.07E-02	0.596	1.002	318.0	5.48E-01
	acenaphthene	2.87E+00	9.20E-05	8.20E-05	293.0	2.40E-02	1.03E+03	1.07E+01	20.0	3000.0	154.00	9.35E-02	2.55E-01	3.88E+00	0.596	1.002	318.0	5.48E-01
	4,4'-DDE	2.00E+00	9.20E-05	8.20E-05	293.0	2.40E-02	7.14E+02	7.44E+00	20.0	3000.0	318.00	1.34E-01	3.66E-01	2.70E+00	0.596	1.002	318.0	5.48E-01
	4,4'-DDT	1.89E+00	9.20E-05	8.20E-05	293.0	2.40E-02	6.76E+02	7.05E+00	20.0	3000.0	354.50	1.42E-01	3.86E-01	2.56E+00	0.596	1.002	318.0	5.48E-01
	4-Methylphenol	4.87E-02	9.60E-07	8.20E-05	293.0	2.40E-02	1.22E+03	1.28E+01	20.0	3000.0	108.13	7.84E-02	2.04E+01	6.58E-02	0.596	1.002	318.0	5.48E-01
child	Chloroform	1.12E+01	2.87E-03	8.20E-05	293.0	2.40E-02	1.16E+03	1.21E+01	20.0	3000.0	118.40	8.24E-02	7.19E-03	1.51E+01	0.596	1.002	318.0	5.48E-01
	Phenanthrene	3.88E+00	1.59E-04	8.20E-05	293.0	2.40E-02	9.53E+02	9.94E+00	20.0	3000.0	178.22	1.01E-01	1.58E-01	6.21E+00	0.596	1.002	318.0	5.48E-01
	2,4-dimethylphenol	3.01E-02	6.30E-07	8.20E-05	293.0	2.40E-02	1.15E+03	1.20E+01	20.0	3000.0	122.16	8.33E-02	3.31E+01	4.07E-02	0.596	1.002	318.0	5.48E-01
	acenaphthene	2.87E+00	9.20E-05	8.20E-05	293.0	2.40E-02	1.03E+03	1.07E+01	20.0	3000.0	154.00	9.35E-02	2.55E-01	3.88E+00	0.596	1.002	318.0	5.48E-01
	4,4'-DDE	2.00E+00	9.20E-05	8.20E-05	293.0	2.40E-02	7.14E+02	7.44E+00	20.0	3000.0	318.00	1.34E-01	3.66E-01	2.70E+00	0.596	1.002	318.0	5.48E-01
	4,4'-DDT	1.89E+00	9.20E-05	8.20E-05	293.0	2.40E-02	6.76E+02	7.05E+00	20.0	3000.0	354.50	1.42E-01	3.86E-01	2.56E+00	0.596	1.002	318.0	5.48E-01
	4-Methylphenol	4.87E-02	9.60E-07	8.20E-05	293.0	2.40E-02	1.22E+03	1.28E+01	20.0	3000.0	108.13	7.84E-02	2.04E+01	6.58E-02	0.596	1.002	318.0	5.48E-01

etc 20 groundwater  
12.5-26

the 28 groundwater

SURFACE SOIL INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 26)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT RESIDENTIAL CHILD

Intake from ingestion of soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times EF \times ED \times IR/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RID$$

INPUTS																
C = contaminant concentration in soil (mg/kg)																
CF = conversion for kg to mg	1E-06															
EF = child exposure frequency (days/yr)	360															
ED = child exposure duration (yr)	6															
IR = child soil ingestion rate (mg/day)	200															
BW = child body weight (kg)	15															
ATc = averaging time for carcinogen (yr)	70															
ATnc = averaging time for noncarcinogen (yr)	6															
DY = days per year (days/year)	365															
CSF = cancer slope factor (mg/kg-day)-1																
specific																
RID = reference dose (mg/kg-day)																
specific																

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Exposure Frequency Child	Exposure Duration Child	Conversion Factor (kg/mg)	Ingestion Rate (mg/day) Child	Body Weight (kg) Child	Average Care Time (days)	Care Dose (mg/kg/day)-1 Child	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncare Time (days)	Noncare Dose (mg/kg/day) Child	Reference Dose (mg/kg/day)	Risk Child	Percent Noncarcinogenic Risk Child
Aluminum	1.4E+03	360	6	1E-06	200	15	25550	3.7E-03	0.00E+00	0%	2190	4.4E-02	1.00E+00	4.4E-02	7%	
Antimony	8.7E+00	360	6	1E-06	200	15	25550	6.2E-06	0.00E+00	0%	2190	7.2E-05	4.00E-04	1.8E-01	31%	
Arsenic	1.7E+00	360	6	1E-06	200	15	25550	1.9E-06	1.78E+00	51%	2190	2.2E-05	3.00E-04	7.4E-02	13%	
Barium	3.5E+01	360	6	1E-06	200	15	25550	3.8E-05	0.00E+00	0%	2190	4.4E-04	7.00E-02	6.3E-03	1%	
Cadmium (soil)	1.6E+00	360	6	1E-06	200	15	25550	1.7E-06	0.00E+00	0%	2190	2.0E-05	1.00E-03	2.0E-02	3%	
Chromium	1.0E+01	360	6	1E-06	200	15	25550	1.1E-05	0.00E+00	0%	2190	1.3E-04	1.00E+00	1.3E-04	0%	
Cobalt	9.1E+01	360	6	1E-06	200	15	25550	1.0E-06	0.00E+00	0%	2190	1.2E-05	6.00E-02	1.9E-04	0%	
Copper	1.3E+02	360	6	1E-06	200	15	25550	1.4E-04	0.00E+00	0%	2190	1.7E-03	3.71E-02	4.5E-02	8%	
Lead	1.7E+02	360	6	1E-06	200	15	25550	1.9E-04	0.00E+00	0%	2190	2.2E-03	0.00E+00	0.0E+00	0%	
Manganese (soil)	3.3E+02	360	6	1E-06	200	15	25550	3.6E-04	0.00E+00	0%	2190	4.2E-03	1.40E-01	3.0E-02	5%	
Mercury	2.0E-01	360	6	1E-06	200	15	25550	3.1E-07	0.00E+00	0%	2190	3.0E-06	3.00E-04	1.2E-02	2%	
Nickel	5.6E+00	360	6	1E-06	200	15	25550	6.1E-06	0.00E+00	0%	2190	7.1E-05	2.00E-02	3.6E-03	1%	
Silver	1.1E+00	360	6	1E-06	200	15	25550	1.2E-06	0.00E+00	0%	2190	1.4E-05	5.00E-03	2.0E-03	0%	
Thallium	8.1E-01	360	6	1E-06	200	15	25550	5.6E-07	0.00E+00	0%	2190	6.8E-06	8.00E-05	8.2E-02	14%	
Vanadium	7.9E+00	360	6	1E-06	200	15	25550	8.7E-06	0.00E+00	0%	2190	1.0E-04	7.00E-03	1.5E-02	2%	
Zinc	1.1E+03	360	6	1E-06	200	15	25550	1.2E-03	0.00E+00	0%	2190	1.4E-02	3.00E-01	4.5E-02	8%	
4,4'-DDD	6.4E-02	360	6	1E-06	200	15	25550	7.0E-08	2.40E-01	1.7E-06	0%	2190	8.2E-07	0.00E+00	0.0E+00	0%
4,4'-DDE	3.5E-01	360	6	1E-06	200	15	25550	3.8E-07	3.40E-01	1.3E-07	2%	2190	4.4E-06	0.00E+00	0.0E+00	0%
4,4'-DDT	1.6E-01	360	6	1E-06	200	15	25550	1.6E-07	3.40E-01	6.6E-08	1%	2190	1.9E-06	6.00E-04	3.6E-03	1%
Anthracene	2.1E-01	360	6	1E-06	200	15	25550	2.2E-07	0.00E+00	0%	2190	2.0E-06	3.00E-01	8.7E-08	0%	
Benz(a)anthracene	2.8E-01	360	6	1E-06	200	15	25550	3.0E-07	7.30E-01	2.2E-07	3%	2190	3.4E-06	0.00E+00	0.0E+00	0%
Benz(a)pyrene	2.7E-01	360	6	1E-06	200	15	25550	2.9E-07	7.30E-01	2.1E-06	33%	2190	3.4E-06	0.00E+00	0.0E+00	0%
Benz(b)fluoranthene	2.9E-01	360	6	1E-06	200	15	25550	3.2E-07	7.30E-01	2.4E-07	4%	2190	3.8E-06	0.00E+00	0.0E+00	0%
Benz(g,h)perylene	2.5E-01	360	6	1E-06	200	15	25550	2.8E-07	0.00E+00	0%	2190	3.2E-06	3.00E-02	1.1E-04	0%	
Carbazole	1.7E-01	360	6	1E-06	200	15	25550	1.9E-07	2.00E-02	3.7E-09	0%	2190	2.2E-06	0.00E+00	0.0E+00	0%
Chrysene	2.8E-01	360	6	1E-06	200	15	25550	3.1E-07	7.30E-03	2.2E-09	0%	2190	3.8E-06	0.00E+00	0.0E+00	0%
Heptachlor epoxide	6.8E-03	360	6	1E-06	200	15	25550	7.6E-09	9.10E+00	6.9E-08	1%	2190	8.8E-06	1.30E-05	6.8E-03	1%
Indeno(1,2,3-cd)pyrene	2.5E-01	360	6	1E-06	200	15	25550	2.8E-07	7.30E-01	2.0E-07	3%	2190	3.2E-06	0.00E+00	0.0E+00	0%
Phenanthrene	2.8E-01	360	6	1E-06	200	15	25550	3.1E-07	0.00E+00	0%	2190	3.8E-06	3.00E-02	1.2E-04	0%	
alpha-Chlordane	3.8E-02	360	6	1E-06	200	15	25550	4.2E-08	1.30E+00	8.4E-08	1%	2190	4.8E-07	6.00E-05	8.1E-03	1%
gamma-Chlordane	1.3E-02	360	6	1E-06	200	15	25550	1.4E-08	1.30E+00	1.8E-08	0%	2190	1.7E-07	6.00E-05	2.9E-03	0%
TOTAL									6.5E-08						5.8E-01	

## SURFACE SOIL DERMAL CONTACT EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 2B)

REMEDIAL INVESTIGATION CTO-0231

MCB CAMP LEJEUNE, NORTH CAROLINA

CURRENT RESIDENTIAL CHILD

Dermal contact with soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \times ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RID$$

Where:

C = contaminant concentration in soil (mg/kg)

1E+00

CF = conversion factor (kg/mg)

2300

SA = child exposed skin surface area (cm<sup>2</sup>)

1

AF = soil/skin adherence factor (mg/cm<sup>2</sup>)

Specific

Abs = fraction absorbed (unitless)

350

EF = child exposure frequency (events/yr)

6

ED = child exposure duration (years)

15

BW = child body weight (kg)

70

ATc = averaging time for carcinogen (yr)

6

ATnc = averaging time for noncarcinogen (yr)

365

DY = day per year (day/yr)

specific

CSF = cancer slope factor (mg/kg-day)<sup>-1</sup>

specific

RID = reference dose (mg/kg-day)

specific

Note: Inputs are scenario and site specific

## INPUTS

COPC	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area Child (cm <sup>2</sup> )	Xference Factor (mg/cm <sup>2</sup> )	Fraction Absorbed (%)	Exposure Frequency (events/yr) Child	Exposure Duration (yrs) Child	Body Weight (kg) Child	Average Carc Time (days) Child	Carc Dose (mg/kg/day) Child	Dermal Adjust Slope Factor (mg/kg-day) <sup>-1</sup> Child	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day) Child	Dermal Adjust Reference Dose (mg/kg/day) Child	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child
Nickel	3.4E+00	1E-00	2300	1	0.001	350	6	15	25500	4.3E-05	0.0E+00	0%	2190	0.0E+04	2.0E-01	2.0E-03	7%	
Antimony	5.7E+00	1E-00	2300	1	0.001	350	6	15	25500	7.1E-06	0.0E+00	0%	2190	0.0E+07	8.0E-05	1.0E-02	28%	
Asenic	1.7E+00	1E-00	2300	1	0.001	350	6	15	25500	2.2E-06	0.0E+00	1.9E-07	21%	2190	2.0E-07	6.0E-05	4.3E-03	11%
Boron	3.5E+01	1E-00	2300	1	0.001	350	6	15	25500	4.4E-07	0.0E+00	0%	2190	5.1E-08	1.4E-02	3.7E-04	1%	
Cadmium (soil)	1.8E+00	1E-00	2300	1	0.001	350	6	15	25500	2.0E-06	0.0E+00	0%	2190	2.3E-07	2.0E-04	1.1E-03	3%	
Chromium	1.0E+01	1E-00	2300	1	0.001	350	6	15	25500	1.3E-07	0.0E+00	0%	2190	1.5E-06	2.0E-01	7.0E-08	0%	
Dobalt	9.1E-01	1E-00	2300	1	0.001	350	6	15	25500	1.1E-08	0.0E+00	0%	2190	1.3E-07	1.2E-02	1.1E-05	0%	
Copper	1.3E+02	1E-00	2300	1	0.001	350	6	15	25500	1.6E-06	0.0E+00	0%	2190	1.9E-05	7.4E-03	2.6E-03	7%	
Lead	1.7E+02	1E-00	2300	1	0.001	350	6	15	25500	2.1E-06	0.0E+00	0%	2190	2.5E-06	0.0E+00	0.0E+00	0%	
Manganese (soil)	3.3E+02	1E-00	2300	1	0.001	350	6	15	25500	4.1E-06	0.0E+00	0%	2190	4.6E-05	2.8E-02	1.7E-03	5%	
Mercury	2.8E-01	1E-00	2300	1	0.001	350	6	15	25500	3.6E-09	0.0E+00	0%	2190	4.2E-08	6.0E-05	7.0E-04	2%	
Nickel	5.0E+00	1E-00	2300	1	0.001	350	6	15	25500	7.0E-06	0.0E+00	0%	2190	8.2E-07	4.0E-03	2.0E-04	1%	
Silver	1.1E+00	1E-00	2300	1	0.001	350	6	15	25500	1.4E-08	0.0E+00	0%	2190	1.7E-07	1.0E-03	1.7E-04	0%	
Thallium	6.1E-01	1E-00	2300	1	0.001	350	6	15	25500	6.5E-09	0.0E+00	0%	2190	7.0E-08	1.6E-06	4.7E-03	13%	
Vanadium	7.9E+00	1E-00	2300	1	0.001	350	6	15	25500	1.0E-07	0.0E+00	0%	2190	1.2E-06	1.4E-03	6.3E-04	2%	
Zinc	1.1E+03	1E-00	2300	1	0.001	350	6	15	25500	1.3E-05	0.0E+00	0%	2190	1.0E-04	6.0E-02	2.6E-03	7%	
1,4'-DDD	6.4E-02	1E-00	2300	1	0.01	350	6	15	25500	8.1E-09	4.8E-01	3.9E-09	0%	2190	8.4E-08	0.0E+00	0.0E+00	0%
1,4'-DDE	3.5E-01	1E-00	2300	1	0.01	350	6	15	25500	4.4E-08	6.0E-01	3.0E-08	3%	2190	5.1E-07	0.0E+00	0.0E+00	0%
1,4'-DDT	1.6E-01	1E-00	2300	1	0.01	350	6	15	25500	1.9E-08	6.0E-01	1.3E-08	1%	2190	2.2E-07	2.6E-04	8.7E-04	2%
Anthracene	2.1E-01	1E-00	2300	1	0.01	350	6	15	25500	2.6E-06	0.0E+00	0%	2190	3.0E-07	1.5E-01	2.0E-08	0%	
Benz(a)anthracene	2.8E-01	1E-00	2300	1	0.01	350	6	15	25500	3.5E-08	1.5E+00	5.1E-08	6%	2190	4.1E-07	0.0E+00	0.0E+00	0%
Benz(a)pyrene	2.7E-01	1E-00	2300	1	0.01	350	6	15	25500	3.4E-08	1.5E+01	4.9E-07	54%	2190	3.9E-07	0.0E+00	0.0E+00	0%
Benz(b)fluoranthene	2.9E-01	1E-00	2300	1	0.01	350	6	15	25500	3.7E-08	1.5E+00	5.4E-08	6%	2190	4.3E-07	0.0E+00	0.0E+00	0%
Benz(g,h,i)perylene	2.5E-01	1E-00	2300	1	0.01	350	6	15	25500	3.2E-08	0.0E+00	0%	2190	3.7E-07	1.5E-02	2.5E-05	0%	
Carbazole	1.7E-01	1E-00	2300	1	0.01	350	6	15	25500	2.1E-08	4.0E-02	8.6E-10	0%	2190	2.5E-07	0.0E+00	0.0E+00	0%
Chrysene	2.8E-01	1E-00	2300	1	0.01	350	6	15	25500	3.5E-08	1.5E-02	5.1E-10	0%	2190	4.1E-07	0.0E+00	0.0E+00	0%
Heptachlor epoxide	6.9E-03	1E-00	2300	1	0.01	350	6	15	25500	8.7E-10	1.8E+01	1.0E-08	2%	2190	1.0E-08	6.8E-08	1.8E-03	4%
Indeno(1,2,3-cd)pyrene	2.8E-01	1E-00	2300	1	0.01	350	6	15	25500	3.2E-08	1.8E+00	4.7E-08	5%	2190	3.7E-07	0.0E+00	0.0E+00	0%
Phenanthrene	2.8E-01	1E-00	2300	1	0.01	350	6	15	25500	3.6E-08	0.0E+00	0.0E+00	0%	2190	4.2E-07	1.5E-02	2.8E-05	0%
alpha-Chlordane	3.8E-02	1E-00	2300	1	0.01	350	6	15	25500	4.8E-09	2.0E+00	1.3E-08	1%	2190	6.0E-08	3.0E-05	1.9E-03	5%
gamma-Chlordane	1.3E-02	1E-00	2300	1	0.01	350	6	15	25500	1.0E-08	2.0E+00	4.2E-09	0%	2190	1.8E-08	3.0E-05	6.3E-04	2%
TOTAL											9.2E-07						3.7E-02	

SURFACE SOIL PARTICULATES INHALATION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT RESIDENTIAL CHILD

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (C * EF * ED * IR * 1/PEF) / (BW * ATc \text{ or } ATnc * DY)$$

Risk = Intake \* CSF or /RfD

Where:

C = contaminant concentration in soil (mg/kg)  
 CSF = carcinogenic slope factor  
 RfD = reference dose for noncarcinogen  
 IR = inhalation rate (m³)  
 EF = child exposure frequency (days)  
 ED = child exposure duration (years)  
 BW = child body weight (kg)  
 ATc = averaging time for carcinogen (yr)  
 ATnc = averaging time for noncarcinogen (yr)  
 DY = day per year (day/yr)  
 PEF = particulate emission factor (m³/kg)

INPUTS  
 Calculated  
 Specific  
 Specific  
 10  
 350  
 6  
 15  
 70  
 366  
 4.63E+09

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Particulate Emission Factor (m³/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m³/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Antimony	3.4E+03	4.6E+09	350	10	6	15	25550	4.0E-08	0.0E+00	0%	2190	4.7E-07	0.0E+00	0.0E+00	0%	
Antimony	5.7E+00	4.6E+09	350	10	6	15	25550	6.7E-11	0.0E+00	0%	2190	7.6E-10	0.0E+00	0.0E+00	0%	
Arsenic	1.7E+00	4.6E+09	350	10	6	15	25550	2.1E-11	1.5E+01	92%	2190	2.4E-10	0.0E+00	0.0E+00	0%	
Barium	3.5E+01	4.6E+09	350	10	6	15	25550	4.1E-10	0.0E+00	0%	2190	4.8E-09	1.4E-04	3.4E-05	99%	
Cadmium (soil)	1.6E+00	4.6E+09	350	10	6	15	25550	1.8E-11	0.0E+00	0%	2190	2.1E-10	0.0E+00	0.0E+00	0%	
Chromium	1.0E+01	4.6E+09	350	10	6	15	25550	1.2E-10	0.0E+00	0%	2190	1.4E-09	0.0E+00	0.0E+00	0%	
Cobalt	9.1E-01	4.6E+09	350	10	6	15	25550	1.1E-11	0.0E+00	0%	2190	1.3E-10	0.0E+00	0.0E+00	0%	
Copper	1.3E+02	4.6E+09	350	10	6	15	25550	1.6E-09	0.0E+00	0%	2190	1.8E-08	0.0E+00	0.0E+00	0%	
Lead	1.7E+02	4.6E+09	350	10	6	15	25550	2.0E-09	0.0E+00	0%	2190	2.3E-08	0.0E+00	0.0E+00	0%	
Manganese (soil)	3.3E+02	4.6E+09	350	10	6	15	25550	3.9E-09	0.0E+00	0%	2190	4.5E-08	0.0E+00	0.0E+00	0%	
Mercury	2.8E-01	4.6E+09	350	10	6	15	25550	3.4E-12	0.0E+00	0%	2190	3.8E-11	8.6E-06	4.8E-07	1%	
Nickel	5.0E+00	4.6E+09	350	10	6	15	25550	6.6E-11	0.0E+00	0%	2190	7.7E-10	0.0E+00	0.0E+00	0%	
Silver	1.1E+00	4.6E+09	350	10	6	15	25550	1.3E-11	0.0E+00	0%	2190	1.8E-10	0.0E+00	0.0E+00	0%	
Thallium	5.1E-01	4.6E+09	350	10	6	15	25550	6.1E-12	0.0E+00	0%	2190	7.1E-11	0.0E+00	0.0E+00	0%	
Titanium	7.9E+00	4.6E+09	350	10	6	15	25550	9.4E-11	0.0E+00	0%	2190	1.1E-09	0.0E+00	0.0E+00	0%	
Zinc	1.1E+03	4.6E+09	350	10	6	15	25550	1.3E-08	0.0E+00	0%	2190	1.5E-07	0.0E+00	0.0E+00	0%	
1,4'-DDO	0.4E-02	4.6E+09	350	10	6	15	25550	7.6E-13	0.0E+00	0%	2190	6.8E-12	0.0E+00	0.0E+00	0%	
1,4'-DDE	3.6E-01	4.6E+09	350	10	6	15	25550	4.1E-12	0.0E+00	0%	2190	4.8E-11	0.0E+00	0.0E+00	0%	
1,4'-DDT	1.5E-01	4.6E+09	350	10	6	15	25550	1.8E-12	3.4E-01	6.0E-13	0%	2190	2.1E-11	0.0E+00	0.0E+00	0%
Anthracene	2.1E-01	4.6E+09	350	10	6	15	25550	2.4E-12	0.0E+00	0%	2190	2.8E-11	0.0E+00	0.0E+00	0%	
Benzo(a)anthracene	2.8E-01	4.6E+09	350	10	6	15	25550	3.3E-12	6.1E-01	2.0E-12	1%	2190	3.6E-11	0.0E+00	0.0E+00	0%
Benzo(a)pyrene	2.7E-01	4.6E+09	350	10	6	15	25550	3.2E-12	6.1E+00	1.9E-11	6%	2190	3.7E-11	0.0E+00	0.0E+00	0%
Benzo(b)fluoranthene	2.9E-01	4.6E+09	350	10	6	15	25550	3.5E-12	6.1E-01	2.1E-12	1%	2190	4.1E-11	0.0E+00	0.0E+00	0%
Benzo(g,h,i)perylene	2.5E-01	4.6E+09	350	10	6	15	25550	3.0E-12	0.0E+00	0.0E+00	0%	2190	3.8E-11	0.0E+00	0.0E+00	0%
Carbazole	1.7E-01	4.6E+09	350	10	6	15	25550	2.0E-12	0.0E+00	0.0E+00	0%	2190	2.3E-11	0.0E+00	0.0E+00	0%
Chrysene	2.8E-01	4.6E+09	350	10	6	15	25550	3.3E-12	6.1E-03	2.0E-14	0%	2190	3.8E-11	0.0E+00	0.0E+00	0%
Heptachlor epoxide	6.9E-03	4.6E+09	350	10	6	15	25550	8.2E-14	9.1E+00	7.4E-13	0%	2190	9.5E-13	0.0E+00	0.0E+00	0%
Indeno(1,2,3-cd)pyrene	2.5E-01	4.6E+09	350	10	6	15	25550	3.0E-12	6.1E-01	1.8E-12	1%	2190	3.8E-11	0.0E+00	0.0E+00	0%
Phenanthrene	2.8E-01	4.6E+09	350	10	6	15	25550	3.4E-12	0.0E+00	0.0E+00	0%	2190	3.9E-11	0.0E+00	0.0E+00	0%
alpha-Chlordane	3.8E-02	4.6E+09	350	10	6	15	25550	4.5E-13	1.3E+00	5.8E-13	0%	2190	6.3E-12	0.0E+00	0.0E+00	0%
gamma-Chlordane	1.3E-02	4.6E+09	350	10	6	15	25550	1.5E-13	1.3E+00	2.0E-13	0%	2190	1.8E-12	0.0E+00	0.0E+00	0%
<b>TOTAL</b>									3.4E-10					3.4E-06		

SUBSURFACE SOIL INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL CHILD

Intake from ingestion of soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * CF * EF * ED * IR/BW * ATc \text{ or } ATnc * DY$$

$$\text{Risk} = \text{Intake} * CSF \text{ or } RID$$

Where:	INPUTS
C = contaminant concentration in soil (mg/kg)	
CF = conversion for kg to mg	1E-06
EF = child exposure frequency (days/yr)	350
ED = child exposure duration (yr)	6
IR = child soil ingestion rate (mg/day)	200
BW = child body weight (kg)	15
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	6
DY = days per year (days/year)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RID = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Exposure Frequency (days/yr) Child	Exposure Duration (yr) Child	Conversion Factor (kg/mg)	Ingestion Rate (mg/day) Child	Body Weight (kg) Child	Average Carc Time (days)	Carc Dose (mg/kg/day) Child	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day) Child	Reference Dose (mg/kg/day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child
Nickel	7.8E+03	350	6	1E-06	200	15	25550	9.5E-03	0.00E+00	0.0E+00	0%	2190	1.0E-01	1.0E+00	1.0E-01	6%
Antimony	1.2E+01	350	6	1E-06	200	15	25550	1.3E-05	0.00E+00	0.0E+00	0%	2190	1.5E-04	4.00E-04	3.8E-01	24%
Arsenic	7.9E+00	350	6	1E-06	200	15	25550	8.7E-06	1.75E+00	1.5E-05	61%	2190	1.0E-04	3.00E-04	3.4E-01	22%
Barium	7.8E+01	350	6	1E-06	200	15	25550	8.8E-05	0.00E+00	0.0E+00	0%	2190	1.0E-03	7.00E-02	1.4E-02	1%
Beryllium	1.4E+01	350	6	1E-06	200	15	25550	1.8E-07	4.30E+00	6.8E-07	3%	2190	1.8E-06	5.00E-03	3.7E-04	0%
Cadmium (soil)	3.2E+00	350	6	1E-06	200	15	25550	3.5E-06	0.00E+00	0.0E+00	0%	2190	4.1E-05	1.00E-03	4.1E-02	3%
Chromium	3.4E+01	350	6	1E-06	200	15	25550	3.7E-05	0.00E+00	0.0E+00	0%	2190	4.4E-04	1.00E+00	4.4E-04	0%
Cobalt	2.7E+00	350	6	1E-06	200	15	25550	2.9E-06	0.00E+00	0.0E+00	0%	2190	3.4E-05	6.00E-02	5.7E-04	0%
Copper	8.9E+02	350	6	1E-06	200	15	25550	9.7E-04	0.00E+00	0.0E+00	0%	2190	1.1E-02	3.71E-02	3.1E-01	20%
Lead	1.7E+03	350	6	1E-06	200	15	25550	1.0E-03	0.00E+00	0.0E+00	0%	2190	2.2E-02	0.00E+00	0.0E+00	0%
Manganese (soil)	1.2E+03	350	6	1E-06	200	15	25550	1.3E-03	0.00E+00	0.0E+00	0%	2190	1.5E-02	1.40E-01	1.1E-01	7%
Mercury	2.3E+01	350	6	1E-06	200	15	25550	2.5E-07	0.00E+00	0.0E+00	0%	2190	2.9E-06	3.00E-04	9.7E-03	1%
Nickel	2.3E+01	350	6	1E-06	200	15	25550	2.6E-05	0.00E+00	0.0E+00	0%	2190	3.0E-04	2.00E-02	1.5E-02	1%
Silver	2.5E+00	350	6	1E-06	200	15	25550	2.7E-06	0.00E+00	0.0E+00	0%	2190	3.2E-05	5.00E-03	6.4E-03	0%
Vanadium	2.4E+01	350	6	1E-06	200	15	25550	2.7E-05	0.00E+00	0.0E+00	0%	2190	3.1E-04	7.00E-03	4.5E-02	3%
Zinc	4.3E+03	350	6	1E-06	200	15	25550	4.7E-03	0.00E+00	0.0E+00	0%	2190	5.5E-02	3.00E-01	1.8E-01	12%
2-Methylnaphthalene	8.0E-02	350	6	1E-06	200	15	25550	9.8E-08	0.00E+00	0.0E+00	0%	2190	1.1E-06	4.00E-02	2.8E-05	0%
4,4'-DDD	8.0E-01	350	6	1E-06	200	15	25550	9.6E-07	2.40E-01	2.3E-07	1%	2190	1.1E-06	0.00E+00	0.0E+00	0%
4,4'-DDE	1.6E+00	350	6	1E-06	200	15	25550	1.6E-05	3.40E-01	6.0E-07	2%	2190	2.0E-05	0.00E+00	0.0E+00	0%
4,4'-DDT	1.2E+01	350	6	1E-06	200	15	25550	1.3E-07	3.40E-01	4.4E-08	0%	2190	1.5E-06	5.00E-04	3.0E-03	0%
Benz(a)anthracene	6.8E-01	350	6	1E-06	200	15	25550	6.4E-07	7.30E-01	4.6E-07	2%	2190	7.4E-06	0.00E+00	0.0E+00	0%
Benz(a)pyrene	6.0E-01	350	6	1E-06	200	15	25550	6.2E-07	7.30E+00	4.6E-06	18%	2190	7.2E-06	0.00E+00	0.0E+00	0%
Benz(0)fluoranthene	6.2E-01	350	6	1E-06	200	15	25550	6.7E-07	7.30E-01	4.9E-07	2%	2190	7.9E-06	0.00E+00	0.0E+00	0%
Benz(g,h)perylene	4.4E-01	350	6	1E-06	200	15	25550	4.9E-07	0.00E+00	0.0E+00	0%	2190	6.7E-06	3.00E-02	1.9E-04	0%
Benz(i,j)fluoranthene	6.4E-01	350	6	1E-06	200	15	25550	5.9E-07	7.30E-02	4.3E-08	0%	2190	6.8E-06	0.00E+00	0.0E+00	0%
Bis(2-ethylhexyl)phthalate	4.8E-01	350	6	1E-06	200	15	25550	6.2E-07	1.40E-02	7.3E-09	0%	2190	6.1E-06	2.00E-02	3.1E-04	0%
Chrysene	8.7E-01	350	6	1E-06	200	15	25550	6.5E-07	7.30E-03	4.6E-09	0%	2190	7.3E-06	0.00E+00	0.0E+00	0%
Dibenz(a,h)anthracene	2.9E-01	350	6	1E-06	200	15	25550	3.1E-07	7.30E+00	2.3E-06	9%	2190	3.6E-06	0.00E+00	0.0E+00	0%
Fluorene	2.8E-01	350	6	1E-06	200	15	25550	3.1E-07	0.00E+00	0.0E+00	0%	2190	3.6E-06	4.00E-02	8.9E-05	0%
Indeno(1,2,3-cd)pyrene	4.3E-01	350	6	1E-06	200	15	25550	4.6E-07	7.30E-01	3.8E-07	1%	2190	5.0E-06	0.00E+00	0.0E+00	0%
Naphthalene	3.4E-01	350	6	1E-06	200	15	25550	3.8E-07	0.00E+00	0.0E+00	0%	2190	4.4E-06	4.00E-02	1.1E-04	0%
Phenanthrene	6.4E-01	350	6	1E-06	200	15	25550	7.0E-07	0.00E+00	0.0E+00	0%	2190	8.2E-06	3.00E-02	2.7E-04	0%
alpha-Chlordane	1.2E-02	350	6	1E-06	200	15	25550	1.3E-08	1.30E+00	1.6E-08	0%	2190	1.5E-07	8.00E-05	2.5E-03	0%
gamma-Chlordane	6.7E-03	350	6	1E-06	200	15	25550	7.3E-09	1.30E+00	9.8E-09	0%	2190	8.5E-06	8.00E-05	1.4E-03	0%
TOTAL							2.6E-05							1.5E+00		

## SUBSURFACE SOIL DERMAL CONTACT EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 26)  
REMEDIAl INVESTIGATION CTO-0231  
MCB CAMP LEJEUNE, NORTH CAROLINA  
FUTURE RESIDENTIAL CHILD

Dental contact with soft is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \cdot CF \cdot SA \cdot AF \cdot Abs \cdot EF \cdot ED/BW \cdot ATc \text{ or } ATnc \cdot DTF$$

$$P_{\text{link}} = \text{Intrinsic} \cdot \text{CSF or FID}$$

Where:

- C = contaminant concentration in soil (mg/kg)
- CF = conversion factor (kg/mg)
- SA = child exposed skin surface area (cm<sup>2</sup>)
- AF = adult to skin adherence factor (mg/cm<sup>2</sup>)
- Abs = fraction absorbed (unitless)
- EF = child exposure frequency (events/yr)
- ED = child exposure duration (years)
- BW = child body weight (kg)
- ATC = averaging time for carcinogen (yr)
- ATHC = averaging time for noncarcinogen (yr)
- DY = day per year (day/yr)
- CSF = cancer slope factor (mg/kg-day)<sup>-1</sup>
- RDF = reference dose (mg/kg-day)

## INPUTS

1E-08
2300
1
Specific
350
6
15
70
6
385
specific
specific

Note: Inputs are scenario and site specific

COPC	Concentration Carcinogen (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> )	Adherence Factor (mg/cm <sup>2</sup> )	Fraction Absorbed (%)	Exposure Frequency (event/day)	Exposure Duration (yr)	Body Weight (kg)	Average Carc Time (day)	Carc Dose (mg/kg/day)	Dermal Adjust Slope Factor	Carcinogenic Risk (mg/kg-day) <sup>-1</sup>	Percent Carcinogenic	Average Noncarc. Time (days)	Noncarc. Dose (mg/kg/day)	Dermal Adjust. Reference Dose	Noncarcinogenic Risk Child	Percent Noncarcinogenic
Naphthalene	2.4E+03	1E-05	2300	1	0.001	360	8	15	25500	0.3E-05	0.0E+00	0%	2150	1.1E-03	2.0E-01	5.7E-03	6%	
Aromatic Hydrocarbons	1.2E+01	1E-05	2300	1	0.001	360	8	15	25500	1.6E-07	0.0E+00	0%	2150	8.0E-05	8.0E-02	2.2E-02	24%	
Asbestos	7.0E+02	1E-05	2300	1	0.001	360	8	15	25500	1.0E-07	8.8E+00	8.7E-07	2%	2150	1.2E-08	6.0E-05	1.9E-02	21%
Benzene	7.0E+01	1E-05	2300	1	0.001	360	8	15	25500	9.0E-07	0.0E+00	0.0E+00	0%	2150	1.2E-05	1.4E-02	8.2E-04	1%
Beryllium	1.4E+01	1E-05	2300	1	0.001	360	8	15	25500	1.0E-08	2.1E+01	3.9E-08	1%	2150	2.1E-08	1.0E-03	2.1E-05	0%
Cadmium (sol)	3.2E+00	1E-05	2300	1	0.001	360	8	15	25500	4.0E-08	0.0E+00	0.0E+00	0%	2150	4.7E-07	2.0E-04	2.4E-03	3%
Chromium	3.4E+01	1E-05	2300	1	0.001	360	8	15	25500	3.4E-08	0.0E+00	0.0E+00	0%	2150	6.0E-08	2.0E-01	2.6E-05	0%
Cobalt	2.7E+00	1E-05	2300	1	0.001	360	8	15	25500	4.3E-07	0.0E+00	0.0E+00	0%	2150	4.0E-07	1.2E-02	3.3E-05	0%
Copper	8.9E+02	1E-05	2300	1	0.001	360	8	15	25500	1.1E-05	0.0E+00	0.0E+00	0%	2150	1.3E-04	7.4E-03	1.8E-02	19%
Lead	1.7E+03	1E-05	2300	1	0.001	360	8	15	25500	2.2E-05	0.0E+00	0.0E+00	0%	2150	2.9E-04	0.0E+00	0.0E+00	0%
Manganese (sol)	1.2E+03	1E-05	2300	1	0.001	360	8	15	25500	1.0E-05	0.0E+00	0.0E+00	0%	2150	1.7E-04	2.9E-02	8.2E-03	7%
Mercury	2.3E-01	1E-05	2300	1	0.001	360	8	15	25500	2.0E-08	0.0E+00	0.0E+00	0%	2150	3.4E-08	6.0E-05	6.0E-04	1%
Nickel	2.3E+01	1E-05	2300	1	0.001	360	8	15	25500	2.9E-07	0.0E+00	0.0E+00	0%	2150	3.4E-03	8.0E-04	1%	1%
Silver	2.9E+00	1E-05	2300	1	0.001	360	8	15	25500	3.2E-08	0.0E+00	0.0E+00	0%	2150	3.7E-07	1.0E-03	3.7E-04	0%
Titanium	2.4E+01	1E-05	2300	1	0.001	360	8	15	25500	3.1E-07	0.0E+00	0.0E+00	0%	2150	3.6E-08	1.4E-03	2.6E-03	5%
Zinc	4.3E+03	1E-05	2300	1	0.001	360	8	15	25500	6.0E-05	0.0E+00	0.0E+00	0%	2150	8.0E-04	8.0E-02	1.1E-02	12%
2-Methylnaphthalene	8.9E-02	1E-05	2300	1	0.01	360	8	15	25500	1.1E-08	0.0E+00	0.0E+00	0%	2150	1.3E-07	2.0E-02	8.0E-05	0%
1,4-DOD	8.6E-01	1E-05	2300	1	0.01	360	8	15	25500	1.1E-07	4.0E-01	5.0E-08	2%	2150	1.3E-08	0.0E+00	0.0E+00	0%
1,4-DDT	1.0E+00	1E-05	2300	1	0.01	360	8	15	25500	2.0E-07	0.0E+00	1.4E-07	0%	2150	2.4E-08	0.0E+00	0.0E+00	0%
Benz(a)anthracene	5.6E-01	1E-05	2300	1	0.01	360	8	15	25500	1.0E-08	0.0E+00	1.0E-08	0%	2150	1.7E-07	2.6E-04	8.0E-04	1%
Benz(a)pyrene	6.6E-01	1E-05	2300	1	0.01	360	8	15	25500	7.0E-08	1.0E-00	1.1E-07	4%	2150	8.0E-07	0.0E+00	0.0E+00	0%
Benz(b)anthracene	6.2E-01	1E-05	2300	1	0.01	360	8	15	25500	7.0E-08	1.0E-01	1.0E-08	3%	2150	8.0E-07	0.0E+00	0.0E+00	0%
Benz(b)phenanthrene	4.4E-01	1E-05	2300	1	0.01	360	8	15	25500	8.0E-08	0.0E+00	0.0E+00	0%	2150	8.0E-07	1.0E-02	4.4E-05	0%
Benz(g,h,i)perylene	8.4E-01	1E-05	2300	1	0.01	360	8	15	25500	6.7E-08	1.0E-01	8.0E-08	0%	2150	7.0E-07	0.0E+00	0.0E+00	0%
Bis(2-Butylnaphthalene)	8.8E-01	1E-05	2300	1	0.01	360	8	15	25500	6.0E-08	2.0E-02	1.7E-08	0%	2150	7.0E-07	1.0E-02	7.0E-05	0%
Drysdale	6.7E-01	1E-05	2300	1	0.01	360	8	15	25500	7.2E-08	1.0E-02	1.1E-08	0%	2150	8.0E-07	0.0E+00	0.0E+00	0%
Oxidized(a)anthracene	2.6E-01	1E-05	2300	1	0.01	360	8	15	25500	3.0E-08	1.0E-01	8.0E-07	17%	2150	4.2E-07	0.0E+00	0.0E+00	0%
Fluorene	2.8E-01	1E-05	2300	1	0.01	360	8	15	25500	3.0E-08	0.0E+00	0.0E+00	0%	2150	4.1E-07	2.0E-02	2.1E-05	0%
Indeno(1,2,3-cd)pyrene	4.3E-01	1E-05	2300	1	0.01	360	8	15	25500	8.0E-08	1.0E-02	8.0E-08	3%	2150	6.0E-07	0.0E+00	0.0E+00	0%
Naphthalene	3.4E-01	1E-05	2300	1	0.01	360	8	15	25500	4.3E-08	0.0E+00	0.0E+00	0%	2150	8.0E-07	2.0E-02	2.0E-05	0%
Phenanthrene	8.4E-01	1E-05	2300	1	0.01	360	8	15	25500	8.1E-08	0.0E+00	0.0E+00	0%	2150	9.4E-07	1.0E-02	8.0E-05	0%
alpha-Chlordane	1.2E-02	1E-05	2300	1	0.01	360	8	15	25500	1.6E-09	2.0E-04	3.0E-08	0%	2150	1.7E-08	3.0E-05	8.7E-04	1%
gamma-Chlordane	6.7E-03	1E-05	2300	1	0.01	360	8	15	25500	8.4E-10	2.0E-04	2.2E-08	0%	2150	9.0E-08	3.0E-05	3.3E-04	0%
TOTAL														3.0E-08		8.1E-02		

SUBSURFACE SOIL PARTICLES INHALATION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL CHILD

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (\text{C} * \text{EF} * \text{ED} * \text{IR} * 1/\text{PEF}) / (\text{BW} * \text{ATc or ATnc} * \text{DY})$$

$$\text{Risk} = \text{Intake} * \text{CSF or RID}$$

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	Calculated
CSF = carcinogenic slope factor	Specific
RID = reference dose for noncarcinogen	Specific
IR = inhalation rate (m3)	10
EF = child exposure frequency (days)	360
ED = child exposure duration (years)	6
BW = child body weight (kg)	15
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	6
DY = day per year (day/yr)	365
PEF = particulate emission factor (m3/kg)	4.6E+09

Note: Inputs are scenario and site specific.

COPC	Concentration Carcinogen (mg/kg)	Particulate Emission Factor (m3/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m3/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Antimony	7.8E+03	4.6E+09	360	10	6	15	25550	9.2E-08	0.0E+00	0.0E+00	0%	2190	1.1E-05	0.0E+00	0.0E+00	0%
Antimony	1.2E+01	4.6E+09	360	10	6	15	25550	1.4E-10	0.0E+00	0.0E+00	0%	2190	1.6E-09	0.0E+00	0.0E+00	0%
Antimony	7.9E+00	4.6E+09	360	10	6	15	25550	9.4E-11	1.5E+01	1.4E-09	94%	2190	1.1E-09	0.0E+00	0.0E+00	0%
Barium	7.8E+01	4.6E+09	360	10	6	15	25550	9.3E-10	0.0E+00	0.0E+00	0%	2190	1.1E-08	1.4E-04	7.6E-05	100%
Beryllium	1.4E-01	4.6E+09	360	10	6	15	25550	1.7E-12	8.4E+00	1.4E-11	1%	2190	2.0E-11	0.0E+00	0.0E+00	0%
Cadmium (soil)	3.2E+00	4.6E+09	360	10	6	15	25550	3.6E-11	0.0E+00	0.0E+00	0%	2190	4.4E-10	0.0E+00	0.0E+00	0%
Chromium	3.4E+01	4.6E+09	360	10	6	15	25550	4.0E-10	0.0E+00	0.0E+00	0%	2190	4.7E-09	0.0E+00	0.0E+00	0%
Cobalt	2.7E+00	4.6E+09	360	10	6	15	25550	3.2E-11	0.0E+00	0.0E+00	0%	2190	3.7E-10	0.0E+00	0.0E+00	0%
Copper	8.9E+02	4.6E+09	360	10	6	15	25550	1.0E-08	0.0E+00	0.0E+00	0%	2190	1.2E-07	0.0E+00	0.0E+00	0%
Lead	1.7E+03	4.6E+09	360	10	6	15	25550	2.0E-08	0.0E+00	0.0E+00	0%	2190	2.4E-07	0.0E+00	0.0E+00	0%
Manganese (soil)	1.2E+03	4.6E+09	360	10	6	15	25550	1.4E-08	0.0E+00	0.0E+00	0%	2190	1.6E-07	0.0E+00	0.0E+00	0%
Mercury	2.3E-01	4.6E+09	360	10	6	15	25550	2.7E-12	0.0E+00	0.0E+00	0%	2190	3.1E-11	8.6E-05	3.7E-07	0%
Nickel	2.3E+01	4.6E+09	360	10	6	15	25550	2.7E-10	0.0E+00	0.0E+00	0%	2190	3.2E-09	0.0E+00	0.0E+00	0%
Silver	2.8E+00	4.6E+09	360	10	6	15	25550	3.0E-11	0.0E+00	0.0E+00	0%	2190	3.5E-10	0.0E+00	0.0E+00	0%
Titanium	2.4E+01	4.6E+09	360	10	6	15	25550	2.9E-10	0.0E+00	0.0E+00	0%	2190	3.4E-09	0.0E+00	0.0E+00	0%
Zinc	4.3E+03	4.6E+09	360	10	6	15	25550	6.1E-08	0.0E+00	0.0E+00	0%	2190	5.0E-07	0.0E+00	0.0E+00	0%
2-Methylnaphthalene	8.9E-02	4.6E+09	360	10	6	15	25550	1.1E-12	0.0E+00	0.0E+00	0%	2190	1.2E-11	0.0E+00	0.0E+00	0%
4,4'-DDD	8.8E-01	4.6E+09	360	10	6	15	25550	1.0E-11	0.0E+00	0.0E+00	0%	2190	1.2E-10	0.0E+00	0.0E+00	0%
4,4'-DDT	1.6E+00	4.6E+09	360	10	6	15	25550	1.9E-11	0.0E+00	0.0E+00	0%	2190	2.2E-10	0.0E+00	0.0E+00	0%
4,4'-DDT	1.2E-01	4.6E+09	360	10	6	15	25550	1.4E-12	3.4E-01	4.7E-13	0%	2190	1.6E-11	0.0E+00	0.0E+00	0%
Benz(a)anthracene	8.8E-01	4.6E+09	360	10	6	15	25550	6.9E-12	6.1E-01	4.2E-12	0%	2190	8.0E-11	0.0E+00	0.0E+00	0%
Benz(a)pyrene	5.6E-01	4.6E+09	360	10	6	15	25550	6.7E-12	6.1E+00	4.1E-11	3%	2190	7.8E-11	0.0E+00	0.0E+00	0%
Benz(b)fluoranthene	6.2E-01	4.6E+09	360	10	6	15	25550	7.3E-12	6.1E-01	4.4E-12	0%	2190	8.5E-11	0.0E+00	0.0E+00	0%
Benz(a,h)perylene	4.4E-01	4.6E+09	360	10	6	15	25550	5.3E-12	0.0E+00	0.0E+00	0%	2190	6.1E-11	0.0E+00	0.0E+00	0%
Benz(k)fluoranthene	6.4E-01	4.6E+09	360	10	6	15	25550	6.3E-12	6.1E-02	3.9E-13	0%	2190	7.4E-11	0.0E+00	0.0E+00	0%
Bis(2-ethylhexyl)phthalate	4.8E-01	4.6E+09	360	10	6	15	25550	5.7E-12	0.0E+00	0.0E+00	0%	2190	6.6E-11	0.0E+00	0.0E+00	0%
Chrysene	8.7E-01	4.6E+09	360	10	6	15	25550	6.8E-12	6.1E-03	4.1E-14	0%	2190	7.9E-11	0.0E+00	0.0E+00	0%
Dibenz(a,h)anthracene	2.9E-01	4.6E+09	360	10	6	15	25550	3.3E-12	6.1E+00	2.0E-11	1%	2190	3.9E-11	0.0E+00	0.0E+00	0%
Fluorene	2.8E-01	4.6E+09	360	10	6	15	25550	3.3E-12	0.0E+00	0.0E+00	0%	2190	3.9E-11	0.0E+00	0.0E+00	0%
Indeno(1,2,3-cd)pyrene	4.5E-01	4.6E+09	360	10	6	15	25550	6.1E-12	6.1E-01	3.1E-12	0%	2190	6.0E-11	0.0E+00	0.0E+00	0%
Naphthalene	3.4E-01	4.6E+09	360	10	6	15	25550	4.1E-12	0.0E+00	0.0E+00	0%	2190	4.7E-11	0.0E+00	0.0E+00	0%
Phenanthrene	6.4E-02	4.6E+09	360	10	6	15	25550	7.0E-12	0.0E+00	0.0E+00	0%	2190	8.8E-11	0.0E+00	0.0E+00	0%
alpha-Chlordane	1.2E-02	4.6E+09	360	10	6	15	25550	1.4E-13	1.3E+00	1.8E-13	0%	2190	1.6E-12	0.0E+00	0.0E+00	0%
gamma-Chlordane	6.7E-03	4.6E+09	360	10	6	15	25550	7.9E-14	1.3E+00	1.0E-13	0%	2190	9.2E-13	0.0E+00	0.0E+00	0%
<b>TOTAL</b>										1.8E-09					7.6E-05	

SEDIMENT INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B - NEW RIVER AND COODLES CREEK)  
 REMEDIAL INVESTIGATION CTO-0201  
 MC8 CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT AND FUTURE RESIDENTIAL CHILD

Intake from ingestion of sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times IR \times CF \times EF \times ED / BW \times ATC \text{ or } ATnc \times DY$$

Risk = Intake \* CSF or RfD

Where:

C = contaminant concentration in sediment (mg/kg)

INPUTS

CF = conversion for kg to mg

1E-03

EF = exposure frequency for child (days/yr)

45

ED = exposure duration for child (yr)

6

IR = soil ingestion rate for child (mg/day)

200

BW = body weight for child (kg)

15

ATC = averaging time for carcinogen (yr)

70

ATnc = averaging time for noncarcinogen (yr)

8

DY = days per year (days/year)

365

CSF = cancer slope factor (mg/kg-day)-1

Specific

RfD = reference dose (mg/kg-day)

Specific

Note: Inputs are scenario and site specific

NEW RIVER

Contaminant	Concentration (mg/kg)	Exposure Frequency (days/yr) Child	Exposure Duration (yr) Child	Ingestion Rate (mg/day) Child	Conversion Factor (kg/mg)	Body Weight (kg) Child	Average Carc Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child
Antimony	2.0E+00	45	6	200	1E-03	15	25500	3.7E-05	0.0E+00	0%	2100	4.3E-04	4.3E-04	1.1E+00	91%	
Asenic	9.0E-01	45	6	200	1E-03	15	25500	1.3E-05	1.8E+00	2.2E-08	72%	2100	1.8E-05	3.0E-04	4.9E-02	4%
Boron	1.0E+01	45	6	200	1E-03	15	25500	2.2E-05	0.0E+00	0%	2100	2.2E-05	7.0E-02	3.0E-04	0%	
Copper	1.3E+03	45	6	200	1E-03	15	25500	1.0E-04	0.0E+00	0.0E+00	0%	2100	2.2E-03	3.7E-02	5.8E-02	5%
Lead	3.0E+04	45	6	200	1E-03	15	25500	5.6E-03	0.0E+00	0.0E+00	0%	2100	8.4E-02	0.0E+00	0.0E+00	0%
Silver	2.1E+00	45	6	200	1E-03	15	25500	3.0E-07	0.0E+00	0.0E+00	0%	2100	3.0E-09	6.0E-03	7.0E-04	0%
Zinc	8.0E+01	45	6	200	1E-03	15	25500	1.0E-05	0.0E+00	0.0E+00	0%	2100	1.8E-04	3.0E-01	4.9E-04	0%
Alkyl-ODD	1.0E-02	45	6	200	1E-03	15	25500	2.1E-09	2.4E-01	8.1E-10	0%	2100	2.6E-08	0.0E+00	0.0E+00	0%
Alkyl-ODE	8.0E-03	45	6	200	1E-03	15	25500	1.2E-08	3.4E-01	4.1E-10	0%	2100	1.4E-08	0.0E+00	0.0E+00	0%
Alkyl-OOT	3.0E-01	45	6	200	1E-03	15	25500	4.2E-08	3.4E-01	1.4E-08	0%	2100	4.9E-07	6.0E-04	9.9E-04	0%
Arinonane	2.0E-01	45	6	200	1E-03	15	25500	3.0E-08	0.0E+00	0.0E+00	0%	2100	4.6E-07	3.0E-01	1.5E-08	0%
Benz(a)anthracene	1.0E-00	45	6	200	1E-03	15	25500	1.7E-07	7.3E-01	1.2E-07	4%	2100	1.9E-08	0.0E+00	0.0E+00	0%
Benz(a)pyrene	6.0E-01	45	6	200	1E-03	15	25500	7.0E-08	7.3E+00	6.0E-07	1%	2100	9.3E-07	0.0E+00	0.0E+00	0%
Benz(b)fluoranthene	1.1E-03	45	6	200	1E-03	15	25500	1.0E-07	7.3E-01	1.1E-07	4%	2100	1.8E-08	0.0E+00	0.0E+00	0%
Benz(g,h)perylene	3.0E-01	45	6	200	1E-03	15	25500	4.0E-08	0.0E+00	0.0E+00	0%	2100	4.9E-07	3.0E-02	1.0E-02	0%
Benz(k)fluoranthene	4.0E-01	45	6	200	1E-03	15	25500	6.0E-08	7.3E-02	5.0E-08	0%	2100	8.0E-07	0.0E+00	0.0E+00	0%
Bis(2-ethylhexyl)phthalate	1.0E+00	45	6	200	1E-03	15	25500	1.0E-07	1.4E-02	2.5E-08	0%	2100	2.1E-08	2.0E-02	1.0E-04	0%
Carbazole	1.0E-01	45	6	200	1E-03	15	25500	2.0E-08	4.5E-10	4.5E-10	0%	2100	2.9E-07	0.0E+00	0.0E+00	0%
Chrysene	1.0E-00	45	6	200	1E-03	15	25500	2.0E-07	7.3E-03	1.0E-08	0%	2100	2.0E-08	0.0E+00	0.0E+00	0%
Dibenzofuran	6.0E-02	45	6	200	1E-03	15	25500	8.0E-08	0.0E+00	0.0E+00	0%	2100	9.0E-08	0.0E+00	0.0E+00	0%
Fluoranthene	1.0E-00	45	6	200	1E-03	15	25500	1.0E-07	0.0E+00	0.0E+00	0%	2100	2.1E-08	4.0E-02	5.1E-05	0%
Indeno(1,2,3-cd)pyrene	3.0E-01	45	6	200	1E-03	15	25500	4.0E-08	7.3E-01	3.1E-08	1%	2100	4.9E-07	0.0E+00	0.0E+00	0%
Phenanthrene	7.7E-01	45	6	200	1E-03	15	25500	1.1E-07	0.0E+00	0.0E+00	0%	2100	1.3E-08	3.0E-02	4.2E-05	0%
Tyrene	1.1E+00	45	6	200	1E-03	15	25500	1.0E-07	0.0E+00	0.0E+00	0%	2100	1.1E-08	3.0E-02	8.0E-05	0%
Alpha-Chordane	6.0E-03	45	6	200	1E-03	15	25500	9.0E-10	1.3E-010	1.2E-09	0%	2100	1.1E-08	8.0E-05	1.8E-04	0%
gamma-Chordane	4.0E-03	45	6	200	1E-03	15	25500	8.0E-10	1.3E-010	8.4E-10	0%	2100	7.0E-09	8.0E-05	1.2E-04	0%
<b>TOTAL</b>								<b>3.1E-05</b>						<b>1.2E+00</b>		

SEDIMENT DERMAL CONTACT Dermal ASSESSMENT  
 OPERABLE UNIT NO.7 (SITE 2B - NEW RIVER AND COODLES CREEK)  
 REMEDIAL INVESTIGATION CTD-0231  
 MC8 CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT AND FUTURE RESIDENTIAL CHILD

The intake from dermal contact to sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

Risk = Intake  $\times$  CSF or RD

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	1.0E-05
CF = conversion factor (kg/mg)	2300
SA = child exposed skin surface area (cm <sup>2</sup> )	1
AF = sediment to skin absorption factor (mg/cm <sup>2</sup> )	Specific
Abs = fraction absorbed (unitless) (contaminant specific)	45
EF = child exposure frequency (events/yr)	6
ED = child exposure duration (years)	15
BW = child body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	6
DY = day per year (days/yr)	365
CSF = cancer slope factor (mg/kg-day) <sup>-1</sup>	Specific
RD = reference dose (mg/kg-day)	Specific

Note: Inputs are scenario and site specific.

NEW RIVER

Contaminant	Concentration Factor (mg/mg)	Surface Area (cm <sup>2</sup> )	Adherence Factor (mg/cm <sup>2</sup> )	AF <sub>CS</sub> Factor (%)	Exposure Frequency (events/yr)	Duration (yrs)	Body Weight (kg)	Average Caco Time (days)	Carc. Dose (mg/kg-day)	Dermal Absorb. Slope Factor (mg/kg-day) <sup>-1</sup>	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncaco Time (days)	Noncaco Dose (mg/kg-day)	Dermat. Absorb. Reference Dose (mg/kg-day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child	
Acetone	2.0E+04	1E-05	2300	1	0.001	45	6	15	22500	1.0E-07	0.0E+00	0%	2100	0.0E+00	0.0E+00	0.0E+00	0%	
Acetic acid	9.0E+00	1E-05	2300	1	0.001	45	6	15	22500	1.0E-08	8.0E+00	1.3E-07	44%	2100	1.7E-07	6.0E-05	2.0E-03	4%
Barium	1.0E+01	1E-05	2300	1	0.001	45	6	15	22500	2.0E-08	0.0E+00	0%	2100	2.0E-07	1.4E-02	2.1E-05	0%	
Copper	1.0E+03	1E-05	2300	1	0.001	45	6	15	22500	2.0E-08	0.0E+00	0%	2100	2.0E-05	7.4E-03	3.4E-03	5%	
Lead	3.0E+04	1E-05	2300	1	0.001	45	6	15	22500	0.0E+00	0.0E+00	0%	2100	0.0E+00	0.0E+00	0.0E+00	0%	
Silver	2.0E+00	1E-05	2300	1	0.001	45	6	15	22500	3.0E-08	0.0E+00	0%	2100	3.0E-04	1.0E+00	4.0E-05	0%	
Zinc	0.0E+01	1E-05	2300	1	0.001	45	6	15	22500	1.0E-07	0.0E+00	0%	2100	1.0E-08	8.0E-05	2.0E-05	0%	
1,4-DCD	1.0E-02	1E-05	2300	1	0.01	45	6	15	22500	2.4E-10	4.0E-01	1.2E-10	0%	2100	2.0E-09	0.0E+00	0.0E+00	0%
1,4-DDE	0.0E+03	1E-05	2300	1	0.01	45	6	15	22500	1.0E-10	0.0E+00	0%	2100	1.0E-09	0.0E+00	0.0E+00	0%	
1,4-DDT	3.0E-01	1E-05	2300	1	0.01	45	6	15	22500	4.0E-09	6.0E-01	9.4E-11	0%	2100	1.0E-09	0.0E+00	0.0E+00	0%
Anthracene	2.0E-01	1E-05	2300	1	0.01	45	6	15	22500	4.0E-09	3.0E-01	3.3E-09	1%	2100	6.7E-08	2.0E-04	2.3E-04	0%
benzo(a)anthracene	1.0E+00	1E-05	2300	1	0.01	45	6	15	22500	1.0E-08	0.0E+00	0%	2100	8.0E-09	1.0E-01	3.0E-07	0%	
benzo(a)pyrene	5.0E-01	1E-05	2300	1	0.01	45	6	15	22500	9.0E-09	1.0E-01	1.3E-07	4%	2100	1.1E-07	0.0E+00	0.0E+00	0%
benzo(b)fluoranthene	1.1E+00	1E-05	2300	1	0.01	45	6	15	22500	1.0E-08	1.0E+00	2.0E-08	9%	2100	2.1E-07	0.0E+00	0.0E+00	0%
benzo(k)fluoranthene	3.0E-01	1E-05	2300	1	0.01	45	6	15	22500	4.0E-09	0.0E+00	0%	2100	6.7E-08	1.0E-02	3.0E-06	0%	
benzo(a,h,i)perylene	4.0E-01	1E-05	2300	1	0.01	45	6	15	22500	7.0E-09	1.0E-01	1.1E-09	0%	2100	9.0E-08	0.0E+00	0.0E+00	0%
benzo(a,h)fluoranthene	4.0E-01	1E-05	2300	1	0.01	45	6	15	22500	2.1E-08	2.0E-02	6.0E-10	0%	2100	2.4E-07	0.0E+00	2.4E-06	0%
2-(2-ethylhexyl)phthalate	1.0E-01	1E-05	2300	1	0.01	45	6	15	22500	2.0E-09	4.0E-03	1.0E-10	0%	2100	3.0E-08	0.0E+00	0.0E+00	0%
Chloroacetaldehyde	1.0E-01	1E-05	2300	1	0.01	45	6	15	22500	2.0E-09	0.0E+00	0%	2100	3.0E-07	0.0E+00	0.0E+00	0%	
Chloroform	1.0E-01	1E-05	2300	1	0.01	45	6	15	22500	2.0E-09	0.0E+00	0%	2100	3.0E-07	0.0E+00	0.0E+00	0%	
Dibenzofuran	0.0E+00	1E-05	2300	1	0.01	45	6	15	22500	9.7E-10	0.0E+00	0%	2100	1.1E-08	0.0E+00	0.0E+00	0%	
Fluorene	1.0E+00	1E-05	2300	1	0.01	45	6	15	22500	2.0E-08	0.0E+00	0%	2100	2.4E-07	2.0E-02	1.2E-05	0%	
Indeno(1,2,3- <i>cd</i> )pyrene	3.0E-01	1E-05	2300	1	0.01	45	6	15	22500	4.0E-09	1.0E+00	7.1E-09	2%	2100	6.7E-08	0.0E+00	0.0E+00	0%
Phenanthrene	7.7E-01	1E-05	2300	1	0.01	45	6	15	22500	1.0E-08	0.0E+00	0%	2100	1.0E-07	1.0E-02	9.7E-06	0%	
Pyrene	1.1E+00	1E-05	2300	1	0.01	45	6	15	22500	1.0E-08	0.0E+00	0%	2100	2.1E-07	1.0E-02	1.4E-05	0%	
alpha-Chlordene	6.0E-03	1E-05	2300	1	0.01	45	6	15	22500	1.1E-10	2.0E-00	2.0E-10	0%	2100	1.0E-08	3.0E-05	4.2E-05	0%
gamma-Chlordene	0.0E+00	1E-05	2300	1	0.01	45	6	15	22500	7.0E-11	2.0E-00	1.0E-10	0%	2100	8.7E-10	3.0E-05	2.0E-05	0%
<b>TOTAL</b>														2.0E-07				

SEDIMENT INGESTION EXPOSURE ASSESSMENT  
OPERABLE UNIT NO. 7 (SITE 28 - NEW RIVER AND COGDELLS CREEK)  
REMEDIAl INVESTIGATION CTD-0231  
MCB CAMP LEJEUNE, NORTH CAROLINA  
CURRENT AND FUTURE RESIDENTIAL CHILD

Intake from ingestion of sediment is calculated as follows:

$$\text{intake (mg/kg-day)} = C \cdot \eta \cdot CF \cdot EF \cdot ED / BW \cdot ATC \text{ or } ATne \cdot CY$$

Risk = threats • CSF or FFD

Where:  
 C = contaminant concentration in sediment (mg/kg)  
 CF = conversion factor kg to mg  
 EF = exposure frequency for child (days/yr)  
 ED = exposure duration for child (yr)  
 R = soil ingestion rate for child (mg/day)  
 BW = body weight for child (kg)  
 ATc = averaging time for carcinogen (yr)  
 ATnc = averaging time for noncarcinogen (yr)  
 DY = days per year (days/year)  
 CSF = cancer slope factor (mg/kg-day)  
 RD = reference dose (mg/kg-day)

## INPUTS

1E-08  
4E-08  
6E-08  
2E-07  
1E-07  
7E-08  
5E-08  
3E-08

Note: Inputs are scenario and site specific

COGGELS CREEK

COPC	Concentration (mg/kg)	Exposure Frequency (days/yr)	Exposure Duration (yr)	Ingestion Rate (mg/day)	Conversion Factor	Body Weight (kg)	Average Carc Time (day)	Carc Date	Slope Factor (mg/kg/day) <sup>-1</sup>	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc. Time (days)	Noncarc. Date (mg/kg/day)	Reference Child	Noncarcinogenic Date (mg/kg/day)	Percent Noncarcinogenic Risk Child
aluminum	2.5E+04	45	6	200	1E-08	15	25500	3.8E-03	1.0E+00	0%	0%	2100	4.1E-02	1.0E+00	4.1E-02	25%
arsenic	7.4E+00	45	6	200	1E-08	15	25500	1.0E-08	1.0E+00	0%	0%	2100	1.2E-05	3.0E-04	4.1E-02	28%
barium	6.8E+01	45	6	200	1E-08	15	25500	8.4E-03	0.0E+00	0%	0%	2100	9.0E-05	2.0E-02	1.4E-03	1%
chromium	2.6E+01	45	6	200	1E-08	15	25500	3.8E-03	0.0E+00	0%	0%	2100	4.5E-05	1.0E+00	4.5E-05	0%
copper	6.1E+01	45	6	200	1E-08	15	25500	7.2E-09	0.0E+00	0%	0%	2100	8.4E-05	3.7E-02	2.3E-03	1%
lead	1.6E+02	45	6	200	1E-08	15	25500	2.3E-05	0.0E+00	0%	0%	2100	2.0E-04	0.0E+00	0.0E+00	0%
manganese (not)	2.5E+02	45	6	200	1E-08	15	25500	3.2E-05	0.0E+00	0%	0%	2100	3.7E-04	1.4E-05	2.7E-03	2%
mercury	3.8E+01	45	6	200	1E-08	15	25500	6.3E-08	0.0E+00	0%	0%	2100	6.2E-07	3.0E-04	2.1E-03	1%
tin	2.9E+00	45	6	200	1E-08	15	25500	4.0E-07	0.0E+00	0%	0%	2100	4.7E-05	8.0E-05	5.6E-02	37%
zirconium	3.8E+01	45	6	200	1E-08	15	25500	5.1E-08	0.0E+00	0%	0%	2100	8.0E-05	7.0E-03	8.4E-03	5%
zinc	1.7E+02	45	6	200	1E-08	15	25500	2.6E-05	0.0E+00	0%	0%	2100	2.0E-04	3.0E-01	9.0E-04	1%
1,3-Dichlorobenzene	4.1E+01	45	6	200	1E-08	15	25500	5.8E-08	4.0E-01	2.0E-08	1%	2100	8.7E-07	0.0E+00	0.0E+00	0%
1,4-DCD	4.8E+01	45	6	200	1E-08	15	25500	6.3E-08	2.4E-01	1.5E-08	1%	2100	7.4E-07	0.0E+00	0.0E+00	0%
1,4-DDE	1.0E+01	45	6	200	1E-08	15	25500	2.2E-08	3.4E-01	7.0E-09	0%	2100	2.0E-07	0.0E+00	0.0E+00	0%
benzo(a)anthracene	1.4E+01	45	6	200	1E-08	15	25500	2.0E-08	7.3E-01	1.4E-08	0%	2100	2.3E-07	0.0E+00	0.0E+00	0%
benzo(a)pyrene	1.1E+00	45	6	200	1E-08	15	25500	1.6E-07	7.3E+00	1.1E-08	37%	2100	1.8E-06	0.0E+00	0.0E+00	0%
benzo(b)fluoranthene	8.3E-02	45	6	200	1E-08	15	25500	8.0E-09	7.3E-01	8.0E-09	0%	2100	1.0E-07	0.0E+00	0.0E+00	0%
benzo(k)fluoranthene	4.8E-02	45	6	200	1E-08	16	26600	6.8E-09	7.3E-02	4.3E-10	0%	2100	8.0E-08	0.0E+00	0.0E+00	0%
2a(2-methylphenyl)phenoxazin	7.0E-01	45	6	200	1E-08	15	25500	9.0E-08	1.4E-02	1.4E-09	0%	2100	1.2E-03	2.0E-02	8.8E-05	0%
Carbon Dioxide	1.3E-02	45	6	200	1E-08	15	25500	1.8E-08	0.0E+00	0%	0%	2100	2.1E-08	1.0E-01	2.1E-07	0%
Chrysene	1.6E-01	45	6	200	1E-08	15	25500	2.3E-08	7.3E-03	1.6E-10	0%	2100	2.8E-07	0.0E+00	0.0E+00	0%
Fluoranthene	3.4E-01	45	6	200	1E-08	15	25500	4.0E-08	0.0E+00	0%	0%	2100	5.0E-07	4.0E-02	1.4E-05	0%
Phenanthrene	2.6E-01	45	6	200	1E-08	15	25500	3.7E-08	0.0E+00	0%	0%	2100	4.3E-07	3.0E-02	1.4E-05	0%
pyrene	2.9E-01	45	6	200	1E-08	15	25500	3.8E-08	0.0E+00	0%	0%	2100	4.1E-07	3.0E-02	1.4E-05	0%
alpha-Chlordane	6.5E-03	45	6	200	1E-08	15	25500	8.8E-10	1.8E+00	1.1E-08	0%	2100	9.7E-08	6.0E-05	1.6E-04	0%
gamma-Chlordane	8.4E-03	45	6	200	1E-08	15	25500	1.2E-08	1.8E+00	1.5E-08	0%	2100	1.4E-08	6.0E-05	2.3E-04	0%
<b>TOTAL</b>												<b>SUM</b>				

SEDIMENT DERMAL CONTACT ASSESSMENT  
OPERABLE UNIT NO.7 (SITE 2B - N.C. RIVER AND COODELS CREEK)  
REMEDIAl INVESTIGATION CTD-0201  
MCB CAMP LEELINE, NORTH CAROLINA  
CURRENT AND FUTURE RESIDENTIAL CHILD

The intake from dermal contact to sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

Risk = Intake  $\times$  CSF or RD

Where:

C = contaminant concentration in soil (mg/kg)  
CF = conversion factor (mg/mg)  
SA = child exposed skin surface area (cm<sup>2</sup>)  
AF = sediment to skin adherence factor (mg/cm<sup>2</sup>)  
Abs = fraction absorbed (unitless) (contaminant specific)  
EF = child exposure frequency (events/yr)  
ED = child exposure duration (year)  
BW = child body weight (kg)  
ATc = averaging time for carcinogen (yr)  
ATnc = averaging time for noncarcinogen (yr)  
DY = day per year (day/yr)  
CSF = cancer slope factor (mg/kg-day)  
RD = reference dose (mg/kg-day)

INPUTS

1.00E-08  
2300  
1  
Specific  
45  
6  
15  
70  
8  
355  
Specific  
Specific

Note: Inputs are scenario and site specific

COODELS CREEK

POPC	Concentration Carcogen (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> ) Child	Adherence Factor (mg/cm <sup>2</sup> )	Abs Factor (%)	Exposure Frequency (events/yr) Child	Exposure Duration (yr) Child	Body Weight (kg) Child	Average Case Time (days) Child	Dose (mg/kg/day) Child	Dermal Risks Slope Factor (mg/kg-day) Child	Carcinogenic Risk Child	Average Noncarc. Noncarcogen Risk Child	Reference Dose (mg/kg/day) Child	Reference Dose (mg/kg-day) Noncarcogen Child	Average Noncarcogen Risk Child
Antimony	2.0E-04	1E-08	2300	1	0.001	45	6	15	23000	1.1E-03	0.0E+00	0%	2100	4.7E-04	2.2E-03	20%
Asenic	7.4E+00	1E-08	2300	1	0.001	45	6	15	23000	1.2E-03	1.0E-07	20%	2100	6.0E-05	2.3E-03	25%
Boron	6.0E+01	1E-08	2300	1	0.001	45	6	15	23000	8.6E-04	0.0E+00	0%	2100	1.1E-03	6.0E-05	1%
Chromium	2.0E+01	1E-08	2300	1	0.001	45	6	15	23000	4.6E-04	0.0E+00	0%	2100	8.2E-07	2.0E-01	2.0E-06
Copper	5.1E+01	1E-08	2300	1	0.001	45	6	15	23000	9.3E-04	0.0E+00	0%	2100	8.6E-07	7.4E-03	1%
Lead	1.0E+02	1E-08	2300	1	0.001	45	6	15	23000	2.0E-07	0.0E+00	0%	2100	3.0E-07	0.0E+00	0%
Manganese (edt)	2.3E+02	1E-08	2300	1	0.001	45	6	15	23000	3.7E-07	0.0E+00	0%	2100	4.3E-06	2.0E-02	1.5E-04
Mercury	3.0E-01	1E-08	2300	1	0.001	45	6	15	23000	6.1E-10	0.0E+00	0%	2100	7.1E-09	0.0E+00	1%
Thallium	2.0E+00	1E-08	2300	1	0.001	45	6	15	23000	4.6E-05	0.0E+00	0%	2100	6.4E-08	1.0E-05	37%
Vanadium	3.0E+01	1E-08	2300	1	0.001	45	6	15	23000	6.0E-04	0.0E+00	0%	2100	6.0E-07	4.0E-04	0%
Zinc	1.7E+02	1E-08	2300	1	0.001	45	6	15	23000	2.8E-07	0.0E+00	0%	2100	3.3E-06	6.0E-05	1%
1,3-Dichlorobenzidine	4.1E-01	1E-08	2300	1	0.01	45	6	15	23000	6.0E-06	0.0E-01	0%	2100	7.6E-06	0.0E+00	0%
1,4-DDO	4.0E-01	1E-08	2300	1	0.01	45	6	15	23000	7.3E-06	4.0E-01	0%	2100	8.5E-06	0.0E+00	0%
1,4-DOE	1.0E-01	1E-08	2300	1	0.01	45	6	15	23000	2.8E-06	6.0E-01	0%	2100	3.0E-06	0.0E+00	0%
3-Benz(a)anthracene	1.4E-01	1E-08	2300	1	0.01	45	6	15	23000	8.5E-06	0.0E+00	0%	2100	2.0E-06	0.0E+00	0%
3-Benz(a)pyrene	1.1E+00	1E-08	2300	1	0.01	45	6	15	23000	1.7E-06	1.0E-01	0%	2100	2.0E-07	0.0E+00	0%
3-Benz(b)fluoranthene	6.3E-02	1E-08	2300	1	0.01	45	6	15	23000	1.0E-06	1.0E-01	0%	2100	2.0E-06	0.0E+00	0%
3-Benz(k)fluoranthene	4.0E-02	1E-08	2300	1	0.01	45	6	15	23000	6.0E-10	1.0E-01	0%	2100	8.5E-11	0.0E+00	0%
3a(2-ethyl)phenolphenoate	7.0E-01	1E-08	2300	1	0.01	45	6	15	23000	1.1E-06	2.0E-02	0%	2100	7.0E-09	0.0E+00	0%
Carbon Disulfide	1.3E-02	1E-08	2300	1	0.01	45	6	15	23000	2.1E-10	0.0E+00	0%	2100	2.0E-09	3.1E-08	0%
Chrysene	1.0E-01	1E-08	2300	1	0.01	45	6	15	23000	2.0E-06	1.0E-01	0%	2100	8.0E-11	0.0E+00	0%
Durene	3.4E-01	1E-08	2300	1	0.01	45	6	15	23000	8.0E-06	0.0E+00	0%	2100	8.0E-08	2.0E-02	3.0E-06
hexamethane	2.0E-01	1E-08	2300	1	0.01	45	6	15	23000	4.0E-06	0.0E+00	0%	2100	4.0E-08	1.0E-03	3.0E-06
Irene	2.0E-01	1E-08	2300	1	0.01	45	6	15	23000	4.1E-06	0.0E+00	0%	2100	4.7E-08	1.0E-03	3.0E-06
Ions-Chlorides	5.0E-03	1E-08	2300	1	0.01	45	6	15	23000	0.0E+11	2.0E-10	0%	2100	1.1E-09	3.0E-03	3.0E-08
gamma-Chlorides	8.4E-03	1E-08	2300	1	0.01	45	6	15	23000	1.4E-10	2.0E+00	0%	2100	1.0E-09	3.0E-05	1%
UTNL										3.0E-07					3.0E-03	

SURFACE WATER INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NOL (SITE 2B - NEW RIVER AND COODLES CREEK)  
 REMEDIAL INVESTIGATION CTD-0221  
 MCAMP CAMP LEADLINE, NORTH CAROLINA  
 CURRENT AND FUTURE CHILD RESIDENT

The intake from the ingestion of surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = Cr * CR * ET * EF * ED/BW * ATc \text{ or } ATnc * DY$$

$$\text{Risk} = \text{Intake} * CSF \text{ or } RD$$

Where:

Cr = contaminant concentration in surface water (mg/l)

CR = contact rate (Lhr/hour)

INPUT

	0.005													
Cr		0.005												
CR			2.6											
ET				45										
EF					45									
ED						5								
BW							15							
ATc								70						
ATnc									6					
DY									365					
CSF										specific				
RD										specific				

Note: Inputs are scenario and site specific

NEW RIVER

COPC	Concentration Carcinogen (mg/l)	Contact Rate (hour) Child	Exposure Time (hrs/week) Child	Exposure Frequency (events/yr) Child	Exposure Duration (years) Child	Body Weight (kg) Child	Averaging Carc. Time (days)	Carc. Dose (mg/kg-day)-1 Child	Cancer Slope Factor (mg/kg-day)-1 Child	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Averaging Time Noncarc. (days)	Noncarc. Dose (mg/kg-day) Child	Reference Dose (mg/kg-day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child	
Nickel	1.7E+00	0.005	2.6	45	6	15	22550	1.3E-05	1.0E+00	0%	2190	1.3E-14	1.0E+00	1.0E-04	3%	3%	
Asenic	4.5E-03	0.005	2.6	45	6	15	22550	3.9E-08	1.8E+00	100%	2190	4.5E-07	3.0E-04	1.0E-03	30%	30%	
Chromium (water)	4.5E-03	0.005	2.6	45	6	15	22550	3.9E-08	0.0E+00	0%	2190	4.5E-07	5.0E-04	9.0E-04	23%	23%	
Copper	1.0E-02	0.005	2.6	45	6	15	22550	1.7E-07	0.0E+00	0%	2190	1.0E-09	3.7E-02	6.2E-05	1%	1%	
Lead	2.5E-02	0.005	2.6	45	6	15	22550	2.1E-07	0.0E+00	0%	2190	2.5E-09	0.0E+00	0.0E+00	0%	0%	
Manganese (water)	6.0E-02	0.005	2.6	45	6	15	22550	4.6E-07	0.0E+00	0%	2190	6.0E-09	6.0E-03	1.1E-03	27%	27%	
Vanadium	6.1E-03	0.005	2.6	45	6	15	22550	8.0E-08	0.0E+00	0%	2190	6.0E-07	7.0E-03	9.0E-05	2%	2%	
Zinc	3.0E-01	0.005	2.6	45	6	15	22550	3.2E-08	0.0E+00	0%	2190	3.0E-05	3.0E-01	1.0E-04	3%	3%	
Al-ODE	4.0E-05	0.005	2.6	45	6	15	22550	3.7E-10	3.4E-01	1.0E-10	0%	2190	4.0E-09	0.0E+00	0.0E+00	0%	0%
Al-ODDE	8.0E-05	0.005	2.6	45	6	15	22550	4.6E-10	2.4E-01	1.0E-10	0%	2190	8.0E-09	0.0E+00	0.0E+00	0%	0%
TOTAL										8.9E-08					3.1E-03		

COODLES CREEK

COPC	Concentration Carcinogen (mg/l)	Contact Rate (hour) Child	Exposure Time (hrs/week) Child	Exposure Frequency (events/yr) Child	Exposure Duration (years) Child	Body Weight (kg) Child	Averaging Carc. Time (days)	Carc. Dose (mg/kg-day)-1 Child	Cancer Slope Factor (mg/kg-day)-1 Child	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Averaging Time Noncarc. (days)	Noncarc. Dose (mg/kg-day) Child	Reference Dose (mg/kg-day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child	
Nickel	1.4E-01	0.005	2.6	45	6	15	22550	1.0E-08	1.0E+00	0%	2190	1.0E-04	1.0E+00	1.0E-04	5%	5%	
Asenic	2.0E-03	0.005	2.6	45	6	15	22550	2.4E-08	1.8E+00	100%	2190	2.7E-07	3.0E-04	9.0E-04	44%	44%	
Lead	3.7E-03	0.005	2.6	45	6	15	22550	3.4E-08	0.0E+00	0%	2190	3.0E-07	0.0E+00	0.0E+00	0%	0%	
Manganese (water)	4.5E-02	0.005	2.6	45	6	15	22550	4.6E-07	0.0E+00	0%	2190	5.0E-09	6.0E-03	1.0E-03	50%	50%	
Vanadium	2.4E-03	0.005	2.6	45	6	15	22550	2.0E-08	0.0E+00	0%	2190	2.0E-07	7.0E-03	3.7E-05	2%	2%	
Zinc	1.2E-02	0.005	2.6	45	6	15	22550	1.1E-07	0.0E+00	0%	2190	1.2E-09	3.0E-01	4.0E-08	2%	2%	
TOTAL										4.1E-08					2.1E-03		

SURFACE WATER DERMAL CO. EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO.7 (SITE 28 - NEW RIVER AND COODLES CREEK)  
 REMEDIAL INVESTIGATION - CTD-0231  
 MCB CAMP LEESBURG, NORTH CAROLINA  
 CURRENT AND FUTURE CHILD RESIDENT

The intake from dermal contact with surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = \text{Cw} * \text{SA} * \text{PC} * \text{ET} * \text{EF} * \text{ED} * \text{CF/BW} * \text{ATc or ATnc} * \text{DY}$$

Risk = Intake \* CSF or RID

Where:

	INPUTS
CW = contaminant concentration in water (mg/l)	
SA = child skin surface available for contact (cm <sup>2</sup> )	2300
PC = contaminant specific dermal permeability (cm/h)	Specific
ET = child exposure time (hours/day)	2.6
EF = child exposure frequency (days/yr)	45
ED = child exposure duration (years)	8
CF = volumetric conversion factor for water (1lta/1000 cm <sup>3</sup> )	0.001
BW = child body weight (kg)	15
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	6
DY = days per year (days)	365
CSF = cancer slope factor (mg/kg-day) <sup>-1</sup>	Specific
RID = reference dose (mg/kg-day)	Specific

Note: Inputs are site and element specific

NEW RIVER

Element	Concentration Carcinogen (mg/l)	Surface Area (cm <sup>2</sup> )	Dermal Permeability (cm/hr)	Exposure Time (hours/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	Volumetric Conversion (l/m <sup>3</sup> )	Body Weight (kg) Child	Averaging Carc Time (days)	Carc. Dose (mg/kg-day)	Dermat. Adjust. Slope Factor (mg/kg-day) <sup>-1</sup>	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc. Nonzero Time (days)	Noncarc. Dose (mg/kg-day)	Dermat. Adjust. Reference Dose (mg/kg-day)	Noncarc. Risk Child	Percent Noncarcinogenic Risk Child
Arsenic	1.7E+00	2300	1.0E-03	2.6	45	5	0.001	15	25500	7.0E-05	0.0E+00	0.0E+00	0%	2100	2.0E-03	0.0E+00	3.1E-04	3%
Arsenic	4.3E-03	2300	1.0E-03	2.6	45	5	0.001	15	25500	1.8E-08	0.0E+00	1.0E-07	74%	2100	2.1E-07	0.0E+00	3.6E-03	39%
Boron (water)	4.2E-03	2300	1.0E-03	2.6	45	5	0.001	15	25500	1.8E-08	0.0E+00	0.0E+00	0%	2100	2.1E-07	1.0E-04	2.1E-03	20%
Copper	1.8E-02	2300	1.0E-03	2.6	45	5	0.001	15	25500	7.0E-08	0.0E+00	0.0E+00	0%	2100	3.1E-07	7.4E-03	1.2E-04	1%
Lead	2.3E-02	2300	4.0E-06	2.6	45	5	0.001	15	25500	3.0E-10	0.0E+00	0.0E+00	0%	2100	4.0E-09	0.0E+00	0.0E+00	0%
Manganese (water)	6.0E-02	2300	1.0E-03	2.6	45	5	0.001	15	25500	2.1E-07	0.0E+00	0.0E+00	0%	2100	2.4E-08	1.0E-03	2.4E-03	27%
Mercury	6.1E-03	2300	1.0E-03	2.6	45	5	0.001	15	25500	2.0E-08	0.0E+00	0.0E+00	0%	2100	3.0E-07	1.4E-03	2.1E-04	2%
Zinc	3.8E-01	2300	6.0E-04	2.6	45	5	0.001	15	25500	9.0E-07	0.0E+00	0.0E+00	0%	2100	1.1E-06	0.0E+00	1.8E-04	2%
K-CDE	4.0E-05	2300	2.4E-01	2.6	45	5	0.001	15	25500	4.0E-08	0.0E+00	2.0E-08	13%	2100	4.7E-07	0.0E+00	0.0E+00	0%
K-DOE	5.0E-05	2300	2.8E-01	2.6	45	5	0.001	15	25500	5.0E-08	4.0E-01	2.0E-08	13%	2100	5.0E-07	0.0E+00	0.0E+00	0%
<b>TOTAL</b>															2.1E-07			

SURFACE WATER DERIVATIVE ASSESSMENT  
 OPERABLE UNIT NO.7 (SITE 2B - COGEELS CREEK AND COGEELS CREEK)  
 REMEDIAL INVESTIGATION - CTO-0291  
 MC8 CAMP LINE, NORTH CAROLINA  
 CURRENT AND FUTURE CHILD RESIDENT

The intake from dermal contact with surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = \text{Cw} * \text{SA} * \text{PC} * \text{ET} * \text{EF} * \text{ED} * \text{CF/BW} * \text{ATc or ATnc} * \text{DY}$$

Risk = Intake \* CSF or RfD

Where:

CW = contaminant concentration in water (mg/l)  
 SA = child skin surface available for contact (cm<sup>2</sup>)  
 PC = contaminant specific dermal permeability (cm/h)  
 ET = child exposure time (hours/day)  
 EF = child exposure frequency (days/year)  
 ED = child exposure duration (years)  
 CF = volumetric conversion factor for water (1 liter/1000 cm<sup>3</sup>)  
 BW = child body weight (kg)  
 ATc = averaging time for carcinogen (yr)  
 ATnc = averaging time for noncarcinogen (yr)  
 DY = days per year (days)  
 CSF = cancer slope factor (mg/kg-day)  
 RfD = reference dose (mg/kg-day)

INPUTS

Specific  
 Specific  
 Specific  
 Specific  
 Specific  
 Specific  
 Specific  
 Specific

Note: Inputs are site and scenario specific.

COGEELS CREEK

COPC	Concentration Carcinogen (mg/l)	Contact Area (cm <sup>2</sup> )	Dermal Permeability (cm/h)	Exposure Time (hours/day)	Exposure Frequency (days/year)	Duration (years)	Volumetric Conversion (L/m <sup>3</sup> )	Body Weight (kg)	Averaging Carc Time (days)	Carc Dose (mg/kg-day)	Dermal Adjust- ment Factor (mg/kg-day) <sup>-1</sup>	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Averaging Noncans Time (days)	Noncancer Dose (mg/kg-day)	Dermal Adjust- ment Factor (mg/kg-day) <sup>-1</sup>	Noncancer Risk Child	Percent Noncancerog- enic Risk Child
chromate	3.4E-01	2300	1.0E-03	2.5	45	5	0.001	15	25500	3.0E-05	0.0E+00	0.0E+00	0%	2190	4.0E-05	2.0E-01	2.0E-04	0%
Asentic	2.8E-03	2300	1.0E-03	2.5	45	6	0.001	15	25500	1.1E-08	0.0E+00	0.0E-08	100%	2190	1.3E-07	0.0E-05	2.1E-03	44%
Lead	3.7E-03	2300	4.0E-06	2.5	45	6	0.001	15	25500	6.2E-11	0.0E+00	0.0E+00	0%	2190	7.3E-10	0.0E+00	0.0E+00	0%
Manganese (water)	4.0E-02	2300	1.0E-03	2.5	45	6	0.001	15	25500	2.0E-07	0.0E+00	0.0E+00	0%	2190	2.4E-09	1.0E-03	2.4E-03	50%
Zincum	2.4E-03	2300	1.0E-03	2.5	45	6	0.001	15	25500	1.0E-08	0.0E+00	0.0E+00	0%	2190	1.2E-07	1.0E-03	8.8E-05	2%
Zinc	1.2E-02	2300	6.0E-04	2.5	45	6	0.001	15	25500	3.0E-08	0.0E+00	0.0E+00	0%	2190	3.6E-07	6.0E-02	5.9E-05	0%
<b>TOTAL</b>																		

GROUNDWATER INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 29)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL ADULT

Intake from drinking water is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * IRw * EF * ED/BW * AT \text{ or } ATnc * DY$$

$$\text{Risk} = \text{Intake} * CSF \text{ or } RID$$

Where:	INPUTS
C = contaminant concentration in water (mg/l)	
IRw = adult daily water ingestion rate (L/Day)	2
EF = adult exposure frequency (days/yr)	350
ED = adult exposure duration (yr)	30
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	30
DY = days per year (day/year)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RID = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

Shallow and Deep Groundwater

COPC	Concentration (mg/l)	Ingestion Rate (L/day) Adult	Exposure Frequency (day/year) Adult	Exposure Duration (year) Adult	Body Weight (kg) Adult	Average Carc Time (days)	Carc Dose (mg/kg-day)-1 Adult	Slope Factor (mg/kg-day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (days)	Noncarc Dose (mg/kg-day) Adult	Reference Dose (mg/kg-day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Arsenic	2.7E-03	2	350	30	70	25550	3.2E-05	1.8E+00	5.8E-05	63%	10950	7.4E-05	3.0E-04	2.5E-01	3%
Barium	7.6E-01	2	350	30	70	25550	8.9E-03	0.0E+00	0%	0%	10950	2.1E-02	7.0E-02	3.0E-01	3%
Lead	3.7E-02	2	350	30	70	25550	4.4E-04	0.0E+00	0%	0%	10950	1.0E-03	0.0E+00	0.0E+00	0%
Manganese (water)	1.6E+00	2	350	30	70	25550	1.7E-02	0.0E+00	0.0E+00	0%	10950	4.0E-02	5.0E-03	7.9E+00	92%
Mercury	5.0E-04	2	350	30	70	25550	5.9E-06	0.0E+00	0.0E+00	0%	10950	1.4E-05	3.0E-04	4.6E-02	1%
2,4-Dimethylphenol	4.3E-03	2	350	30	70	25550	5.0E-05	0.0E+00	0.0E+00	0%	10950	1.2E-04	2.0E-02	5.9E-03	0%
Aceanaphthalene	1.0E-02	2	350	30	70	25550	1.2E-04	0.0E+00	0.0E+00	0%	10950	2.7E-04	6.0E-02	4.6E-03	0%
4,4'-DDE	1.6E-03	2	350	30	70	25550	1.9E-05	3.4E-01	6.4E-06	7%	10950	4.4E-05	0.0E+00	0.0E+00	0%
4,4'-DDD	9.0E-03	2	350	30	70	25550	1.1E-04	2.4E-01	2.6E-05	29%	10950	2.5E-04	0.0E+00	0.0E+00	0%
4,4'-DDT	1.8E-04	2	350	30	70	25550	2.1E-06	3.4E-01	7.2E-07	1%	10950	4.9E-06	5.0E-04	9.9E-03	0%
Chloroform	1.2E-03	2	350	30	70	25550	1.4E-05	6.1E-03	8.8E-08	0%	10950	3.3E-05	1.0E-02	3.3E-03	0%
2-Methylnaphthalene	9.3E-03	2	350	30	70	25550	1.1E-04	0.0E+00	0.0E+00	0%	10950	2.5E-04	4.0E-02	6.4E-03	0%
Phenanthrene	6.6E-03	2	350	30	70	25550	7.7E-05	0.0E+00	0.0E+00	0%	10950	1.8E-04	3.0E-02	6.0E-03	0%
4-Methylphenol	9.3E-03	2	350	30	70	25550	1.1E-04	0.0E+00	0.0E+00	0%	10950	2.5E-04	5.0E-03	5.1E-02	1%
TOTAL							8.8E-05						8.8E+00		

## GROUNDWATER DERMAL CONTAMINANT EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 26)

MEDICAL INVESTIGATION (D-020)

MCB CAMP LEJEUNE, NORTH CAROLINA

FUTURE RESIDENTIAL ADULT

Dermal Contact from groundwater is calculated as follows:

$$\text{Intake (mg/kg-day)} = \text{CW} * \text{SA} * \text{PC} * \text{ET} * \text{EF} * \text{ED} * \text{CF/BW} * \text{ATc or ATnc} * \text{DF}$$

Risk = Intake \* CSF or RID

Where:

CW = contaminant concentration in water (mg/l)

## INPUTS

SA = adult skin surface available for contact (cm<sup>2</sup>)

23000

PC = contaminant specific dermal permeability (cm/h)

Specific

ET = adult exposure time (hours/day)

0.25

EF = adult exposure frequency (days/yr)

350

ED = adult exposure duration (years)

30

CF = volumetric conversion factor for water (liter/1000 cm<sup>3</sup>)

0.001

BW = adult body weight (kg)

70

ATc = averaging time for carcinogen (yr)

70

ATnc = averaging time for noncarcinogen (yr)

30

DF = days per year (day)

365

Note: Inputs are site and scenario specific

## SHALLOW AND DEEP GROUNDWATER

CP/C	Concentration Carcinogen (mg/l)	Surface Area (cm <sup>2</sup> )	Dermal Permeability (cm/hr)	Exposure Time (hours/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	Volumetric Conversion (L/m <sup>3</sup> )	Body Weight (kg)	Averaging Carc Time (years)	Carc Dose (mg/kg-day)	Dermal Adj. Slope Factor (mg/kg-day) <sup>-1</sup>	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (years)	Noncarc Dose (mg/kg-day)	Dermal Adjust. Reference Dose (mg/kg-day)	Noncarc Risk Adult	Percent Noncarcinogenic Risk Adult
Asinic	2.7E-03	23000	1.0E+03	0.25	350	30	0.001	70	25500	9.2E-08	8.0E+00	2%	10960	2.1E-07	0.0E+00	3.0E-13	2%	
Barium	7.0E-01	23000	1.0E+03	0.25	350	30	0.001	70	25500	2.0E-05	0.0E+00	0%	10960	8.0E-05	1.4E-02	4.3E-03	2%	
Lead	3.7E-02	23000	4.0E-06	0.25	350	30	0.001	70	25500	6.0E-08	0.0E+00	0%	10960	1.2E-08	0.0E+00	0.0E+00	0%	
Manganese (water)	1.0E+00	23000	1.0E+03	0.25	350	30	0.001	70	25500	4.9E-05	0.0E+00	0%	10960	1.1E-04	1.0E-03	1.1E-01	67%	
Mercury	6.0E-04	23000	1.0E+03	0.25	350	30	0.001	70	25500	1.7E-08	0.0E+00	0%	10960	3.9E-08	6.0E-05	9.0E-04	0%	
p-Dimethylphenol	4.3E-03	23000	1.6E-02	0.25	350	30	0.001	70	25500	2.2E-08	0.0E+00	0%	10960	5.1E-08	1.0E-02	5.1E-04	0%	
Acenaphthene	1.0E-02	23000	1.5E-01	0.25	350	30	0.001	70	25500	5.1E-05	0.0E+00	0%	10960	1.2E-04	3.0E-02	4.0E-03	2%	
Acenaphthene	1.0E-02	23000	2.4E-01	0.25	350	30	0.001	70	25500	1.3E-05	8.8E-01	17%	10960	3.0E-05	0.0E+00	0.0E+00	0%	
Acenaphthene	1.0E-02	23000	2.8E-01	0.25	350	30	0.001	70	25500	8.5E-05	4.8E-01	4.1E-05	78%	1.0E-04	2.0E-04	0.0E+00	0.0E+00	0%
Acenaphthene	1.0E-02	23000	4.3E-01	0.25	350	30	0.001	70	25500	2.0E-08	8.8E-01	1.1E-05	3%	10960	6.1E-08	2.5E-04	2.4E-02	14%
Chloroform	1.2E-03	23000	8.0E-03	0.25	350	30	0.001	70	25500	3.0E-07	7.8E-03	2.7E-08	0%	10960	8.4E-07	8.0E-03	1.1E-04	0%
p-Methylnaphthalene	9.3E-03	23000	1.4E-01	0.25	350	30	0.001	70	25500	4.5E-05	0.0E+00	0%	10960	1.0E-04	2.0E-02	6.0E-03	3%	
p-Nitroaniline	6.0E-03	23000	2.7E-01	0.25	350	30	0.001	70	25500	6.0E-05	0.0E+00	0%	10960	1.4E-04	1.0E-02	9.0E-03	5%	
p-Nitroaniline	6.0E-03	23000	1.6E-02	0.25	350	30	0.001	70	25500	5.7E-08	0.0E+00	0%	10960	1.3E-05	2.5E-03	5.5E-03	3%	
TOTAL																		

CTO-231 SITE 28 GROUNDWATER INHALATION  
 13-Jun-95  
 FILE: SHOW.WQ1

GROUNDWATER INHALATION - RME CASE

CHEMICAL	ADULT	C rme	ED	EF	AT	ATC	CDI	CDIC	RFD	PF	HI	CR	% CONTRIB	% CONTRIB
		mg/kg/shw	y	shwr/y	d	d	mg/kg/d	mg/kg/d	mg/kg/d	(mg/kg/d) <sup>-1</sup>		NC RISK	CARC RISK	
Chloroform		1.2E-05	30	350	10950	25550	1.1E-05	4.8E-06	0.0E+00	8.1E-02	---	3.9E-07	0%	88%
Phenanthrene		2.6E-05	30	350	10950	25550	2.5E-05	1.1E-05	0.0E+00	0.0E+00	---	0.0E+00	0%	0%
2, 4-Dimethylphenol		1.4E-07	30	350	10950	25550	1.4E-07	5.9E-08	0.0E+00	0.0E+00	---	0.0E+00	0%	0%
Acenaphthene		3.0E-05	30	350	10950	25550	2.9E-05	1.2E-05	0.0E+00	0.0E+00	---	0.0E+00	0%	0%
4, 4'-DDE		3.4E-06	30	350	10950	25550	3.3E-06	1.4E-06	0.0E+00	0.0E+00	---	0.0E+00	0%	0%
4, 4'-DDT		3.6E-07	30	350	10950	25550	3.5E-07	1.5E-07	0.0E+00	3.4E-01	---	5.1E-08	0%	12%
4-Methylphenol		5.0E-07	30	350	10950	25550	4.8E-07	2.1E-07	0.0E+00	0.0E+00	---	0.0E+00	0%	0%
										TOTAL	0.0E+00	4.4E-07		

site 28 groundwater  
13-Jun-95 file: showmod.wq1

CAS No.	TARGET COMPOUND LIST CONSTITUTE	OVERALL MASS TRANSF COEFFICIENT	HENRY'S LAW CONSTAN	GAS CONSTAN	ABSOLUTE TEMP.	GAS CONST.	GAS-FLM MASS TRANSF. COEFFICIENT	Liquid-film COEFFICIENT	CARBON DIOXIDE TRANS. COEFF.	WATER TRANS. COEFF.	MOLECULAR WEIGHT	RECIPROCAL OF K	(RT)/(H)*KG)	OVERALL MASS TRANSF. COEFF.	WATER VISCOSITY AT T1	WATER VISCOSITY AT T1	SHOWER WATER TEMP. ((T1)*(U1))/((T2)*(U1))	
		(KL/cm/hr)	(atm-m3/mol-K)	(atm-m3/mol-K)	(K)	(RT)	(kg/cm/hr)	(kg/cm/hr)	(kg/cm/hr)	(cm/hr)	(gram/mole)	(K)	(ATTs)	(Us)	(U1)	(Ts)		
<b>site 28 PME adult</b>																		
	Chloroform	1.12E+01	2.87E-03	8.20E-05	293.0	2.40E-02	1.16E+03	1.21E+01	20.0	3000.0	119.40	8.24E-02	7.19E-03	1.51E+01	0.596	1.002	318.0	5.48E-01
	Phenanthrene	3.98E+00	1.59E-04	8.20E-05	293.0	2.40E-02	9.63E+02	9.94E+00	20.0	3000.0	178.22	1.01E-01	1.58E-01	5.21E+00	0.596	1.002	318.0	5.48E-01
	2,4-dimethylphenol	3.01E-02	6.30E-07	8.20E-05	293.0	2.40E-02	1.15E+03	1.20E+01	20.0	3000.0	122.16	8.33E-02	3.31E+01	4.07E-02	0.596	1.002	318.0	5.48E-01
	acenaphthene	2.97E+00	9.20E-05	8.20E-05	293.0	2.40E-02	1.03E+03	1.07E+01	20.0	3000.0	154.00	9.35E-02	2.55E-01	3.88E+00	0.596	1.002	318.0	5.48E-01
	4,4'-DDE	2.00E+00	9.20E-05	8.20E-05	293.0	2.40E-02	7.14E+02	7.44E+00	20.0	3000.0	318.00	1.34E-01	3.66E-01	2.70E+00	0.596	1.002	318.0	5.48E-01
	4,4'-DDT	1.89E+00	9.20E-05	8.20E-05	293.0	2.40E-02	6.76E+02	7.05E+00	20.0	3000.0	354.50	1.42E-01	3.86E-01	2.56E+00	0.596	1.002	318.0	5.48E-01
	4-Methylphenol	4.87E-02	9.60E-07	8.20E-05	293.0	2.40E-02	1.22E+03	1.28E+01	20.0	3000.0	108.13	7.84E-02	2.04E+01	6.58E-02	0.596	1.002	318.0	5.48E-01
<b>child</b>																		
	Chloroform	1.12E+01	2.87E-03	8.20E-05	293.0	2.40E-02	1.16E+03	1.21E+01	20.0	3000.0	119.40	8.24E-02	7.19E-03	1.51E+01	0.596	1.002	318.0	5.48E-01
	Phenanthrene	3.98E+00	1.59E-04	8.20E-05	293.0	2.40E-02	9.63E+02	9.94E+00	20.0	3000.0	178.22	1.01E-01	1.58E-01	5.21E+00	0.596	1.002	318.0	5.48E-01
	2,4-dimethylphenol	3.01E-02	6.30E-07	8.20E-05	293.0	2.40E-02	1.15E+03	1.20E+01	20.0	3000.0	122.16	8.33E-02	3.31E+01	4.07E-02	0.596	1.002	318.0	5.48E-01
	acenaphthene	2.97E+00	9.20E-05	8.20E-05	293.0	2.40E-02	1.03E+03	1.07E+01	20.0	3000.0	154.00	9.35E-02	2.55E-01	3.88E+00	0.596	1.002	318.0	5.48E-01
	4,4'-DDE	2.00E+00	9.20E-05	8.20E-05	293.0	2.40E-02	7.14E+02	7.44E+00	20.0	3000.0	318.00	1.34E-01	3.66E-01	2.70E+00	0.596	1.002	318.0	5.48E-01
	4,4'-DDT	1.89E+00	9.20E-05	8.20E-05	293.0	2.40E-02	6.76E+02	7.05E+00	20.0	3000.0	354.50	1.42E-01	3.86E-01	2.56E+00	0.596	1.002	318.0	5.48E-01
	4-Methylphenol	4.87E-02	9.60E-07	8.20E-05	293.0	2.40E-02	1.22E+03	1.28E+01	20.0	3000.0	108.13	7.84E-02	2.04E+01	6.58E-02	0.596	1.002	318.0	5.48E-01

site 20 groundwater  
13-June-05

CHS No.	TARGET COMPOUND LIST CONSTITUENT	CONCENTRATION				SHOWER		SHOWER		Kd's		Droplet Size		INDOOR VOC		SHOWER		SHOWER		INDOOR AIR		AIR EXCHANGE		SHOWER		TOTAL		exp(0.0)		exp(0.1)		INHALATION		VENTILATION		BODY		exp(-0.0)		exp(0.0)		exp(0.1)	
		SHOWER DROPLET CONCENTRATION		WATER DROPLET DIAETER		DROPOLET DROPOUT		INDOOR VOC CONCENTRATION		SHOWER WATER FLOW RATE		SHOWER WATER FLOW RATE		INDOOR AIR VOC CONC. ATTIME		SHOWER WATER FLOW RATE		INDOOR AIR VOC CONC. ATTIME		SHOWER WATER FLOW RATE		TOTAL TIME		exp(0.0)		exp(0.1)		PER SHOWER		RATE		BODY WEIGHT		exp(-0.0)		exp(0.0)		exp(0.1)					
		Droplet Conc Mg/m <sup>3</sup>	Droplet Conc Mg/m <sup>3</sup>	Dia mm	Dia mm	Droplet Droponut Sec	Droplet Droponut Sec	Indoor Voc Conc Mg/m <sup>3</sup>	Indoor Voc Conc Mg/m <sup>3</sup>	Shower Flow Rate L/min	Shower Flow Rate L/min	Shower Flow Rate L/min	Shower Flow Rate L/min	Indoor Air Voc Conc Mg/m <sup>3</sup>	Indoor Air Voc Conc Mg/m <sup>3</sup>	Shower Flow Rate L/min	Shower Flow Rate L/min	Shower Flow Rate L/min	Shower Flow Rate L/min	Total Time Sec	exp(0.0)	exp(0.1)	Per Shower Time Sec	Rate Sec	Body Weight kg	exp(-0.0)	exp(0.0)	exp(0.1)	Exposure Time Sec	Exposure Rate Sec	Body Weight kg	exp(-0.0)	exp(0.0)	exp(0.1)									
<b>Me-26 PME</b>																																											
adult	Chlordimefuran	4.74E-01	1.2	1.00	2.00	3.02E+01	6.00E+01	-5.00E-01	6.048E-01	7.80E-01	10	0	3.42E+01	0.0088	12	15	4.08E-01	8.89E-01	1.17E-05	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	1.20E-05																
	Phenanthrene	1.02E+03	6.6	1.00	2.00	1.04E+01	6.00E+01	-1.74E-01	6.404E-01	1.78E+02	10	0	7.98E+01	0.0088	12	15	4.08E-01	8.89E-01	2.88E-05	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	2.02E-05																
	2,4-dimethylphenol	6.93E-03	4.3	1.00	2.00	8.14E-02	6.00E+01	-1.30E-03	9.888E-01	9.71E-03	10	0	4.20E-01	0.0088	12	15	4.08E-01	8.89E-01	1.44E-07	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	1.67E-07																
	acenaphthene	1.21E+00	100	1.00	2.00	7.79E+00	6.00E+01	-1.20E-01	8.789E-01	2.02E+00	10	0	8.74E+01	0.0088	12	15	4.08E-01	8.89E-01	2.88E-05	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	3.48E-05																
	4,4'-DEE	1.39E-01	1.8	1.00	2.00	8.40E+00	6.00E+01	-4.02E-02	8.810E-01	2.30E-01	10	0	6.62E+01	0.0088	12	15	4.08E-01	8.89E-01	2.88E-05	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	3.90E-05																
	4,4'-DDT	1.47E-02	0.2	1.00	2.00	6.11E+00	6.00E+01	-4.82E-02	8.820E-01	2.48E-02	10	0	1.08E+02	0.0088	12	15	4.08E-01	8.89E-01	2.88E-07	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	4.22E-07																
	4,4'-DHT	0.22E-02	0.8	1.00	2.00	1.02E-01	6.00E+01	-0.12E-02	8.829E-01	0.40E-02	10	0	1.42E+02	0.0088	12	15	4.08E-01	8.89E-01	2.88E-07	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	5.82E-07																
child	Chlordimefuran	4.74E-01	1.2	1.00	2.00	3.02E+01	6.00E+01	-4.02E-01	6.048E-01	7.80E-01	10	0	8.42E+01	0.0088	12	15	4.08E-01	8.89E-01	7.70E-05	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	8.20E-05																
	Phenanthrene	1.02E+03	6.6	1.00	2.00	1.04E+01	6.00E+01	-1.74E-01	8.404E-01	1.78E+00	10	0	7.98E+01	0.0088	12	15	4.08E-01	8.89E-01	1.87E-04	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	1.80E-04																
	2,4-dimethylphenol	6.93E-03	4.3	1.00	2.00	8.14E-02	6.00E+01	-4.30E-03	9.888E-01	9.71E-02	10	0	4.20E-01	0.0088	12	15	4.08E-01	8.89E-01	8.71E-07	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	1.91E-06																
	acenaphthene	1.21E+00	100	1.00	2.00	7.79E+00	6.00E+01	-1.20E-01	8.789E-01	2.02E+00	10	0	8.74E+01	0.0088	12	15	4.08E-01	8.89E-01	1.87E-04	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	2.11E-04																
	4,4'-DEE	1.39E-01	1.8	1.00	2.00	8.40E+00	6.00E+01	-4.02E-02	8.810E-01	2.30E-01	10	0	6.62E+01	0.0088	12	15	4.08E-01	8.89E-01	2.88E-05	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	2.40E-05																
	4,4'-DDT	1.47E-02	0.2	1.00	2.00	6.11E+00	6.00E+01	-4.82E-02	8.820E-01	2.48E-02	10	0	1.08E+02	0.0088	12	15	4.08E-01	8.89E-01	2.88E-08	10	70	8.89E-01	15	1.18E+02	1.17E+02	-3	5.82E-08																

SURFACE SOIL INGESTION EXPOSURE ASSESSMENT  
 OPERATIONAL UNIT NO. 7 (SITE 26)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT RESIDENTIAL ADULT

Intake from ingestion of soil is calculated as follows:

$$\text{Intake}^* (\text{mg/kg-day}) = C \times CF \times EF \times ED \times IR/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RfD$$

Where:

C = contaminant concentration in soil (mg/kg)	INPUTS
CF = conversion for kg to mg	1E-06
EF = adult exposure frequency (days/yr)	350
ED = adult exposure duration (yr)	30
IR = adult soil ingestion rate (mg/day)	100
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	30
DY = days per year (days/year)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RfD = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Exposure Frequency (days/yr)	Exposure Duration (yr)	Conversion Factor (kg/mg)	Ingestion Rate (mg/day)	Body Weight (kg)	Average Care Time (days)	Care Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncare Time (days)	Noncare Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	3.4E+03	350	30	1E-06	100	70	25550	2.0E-03	0.0E+00	0%	10950	4.7E-03	1.0E+00	4.7E-03	7%	
Antimony	8.7E+00	350	30	1E-06	100	70	25550	3.3E-06	0.0E+00	0%	10950	7.8E-06	4.0E-04	1.9E-02	31%	
Arsenic	1.7E+00	350	30	1E-06	100	70	25550	1.0E-06	1.8E+00	51%	10950	2.4E-06	3.0E-04	8.0E-03	13%	
Barium	3.5E+01	350	30	1E-06	100	70	25550	2.0E-06	0.0E+00	0%	10950	4.8E-05	7.0E-02	8.8E-04	1%	
Cadmium (soil)	1.0E+00	350	30	1E-06	100	70	25550	9.1E-07	0.0E+00	0%	10950	2.1E-06	1.0E-03	2.1E-03	3%	
Chromium	1.0E+01	350	30	1E-06	100	70	25550	8.1E-06	0.0E+00	0%	10950	1.4E-05	1.0E+00	1.4E-05	0%	
Cobalt	9.1E-01	350	30	1E-06	100	70	25550	5.3E-07	0.0E+00	0%	10950	1.2E-06	6.0E-02	2.1E-05	0%	
Copper	1.0E+02	350	30	1E-06	100	70	25550	7.6E-06	0.0E+00	0%	10950	1.8E-04	3.7E-02	4.8E-03	8%	
Lead	1.7E+02	350	30	1E-06	100	70	25550	1.0E-04	0.0E+00	0%	10950	2.3E-04	0.0E+00	0.0E+00	0%	
Manganese (soil)	3.5E+02	350	30	1E-06	100	70	25550	1.9E-04	0.0E+00	0%	10950	4.5E-04	1.4E-01	3.2E-03	5%	
Mercury	2.8E-01	350	30	1E-06	100	70	25550	1.7E-07	0.0E+00	0%	10950	3.9E-07	3.0E-04	1.3E-03	2%	
Nickel	6.0E+00	350	30	1E-06	100	70	25550	3.3E-06	0.0E+00	0%	10950	7.6E-06	2.0E-02	3.8E-04	1%	
Silver	1.1E+00	350	30	1E-06	100	70	25550	8.6E-07	0.0E+00	0%	10950	1.5E-06	5.0E-03	3.1E-04	0%	
Thallium	5.1E-01	350	30	1E-06	100	70	25550	3.0E-07	0.0E+00	0%	10950	7.0E-07	8.0E-05	8.8E-03	14%	
Vanadium	7.9E+00	350	30	1E-06	100	70	25550	4.7E-06	0.0E+00	0%	10950	1.1E-05	7.0E-03	1.6E-03	2%	
Zinc	1.1E+03	350	30	1E-06	100	70	25550	8.3E-04	0.0E+00	0%	10950	1.5E-03	3.0E-01	4.9E-03	8%	
4,4'-DDD	6.4E-02	350	30	1E-06	100	70	25550	3.8E-06	2.4E-01	9.0E-09	0%	10950	8.8E-08	0.0E+00	0.0E+00	0%
4,4'-DDE	3.5E-01	350	30	1E-06	100	70	25550	2.0E-07	3.4E-01	6.9E-08	2%	10950	4.8E-07	0.0E+00	0.0E+00	0%
4,4'-DDT	1.5E-01	350	30	1E-06	100	70	25550	9.7E-06	3.4E-01	3.0E-08	1%	10950	2.0E-07	6.0E-04	4.1E-04	1%
Anthracene	2.1E-01	350	30	1E-06	100	70	25550	1.2E-07	0.0E+00	0.0E+00	0%	10950	2.8E-07	3.0E-01	9.4E-07	0%
Benzo(a)anthracene	2.8E-01	350	30	1E-06	100	70	25550	1.8E-07	7.3E-01	1.2E-07	3%	10950	3.8E-07	0.0E+00	0.0E+00	0%
Benzo(a)pyrene	2.7E-01	350	30	1E-06	100	70	25550	1.8E-07	7.3E+00	1.1E-06	33%	10950	3.7E-07	0.0E+00	0.0E+00	0%
Benzo(b)fluoranthene	2.9E-01	350	30	1E-06	100	70	25550	1.7E-07	7.3E-01	1.3E-07	4%	10950	4.0E-07	0.0E+00	0.0E+00	0%
Benzo(g,h,i)perylene	2.5E-01	350	30	1E-06	100	70	25550	1.6E-07	0.0E+00	0.0E+00	0%	10950	3.5E-07	3.0E-02	1.2E-05	0%
Carbazole	1.7E-01	350	30	1E-06	100	70	25550	1.0E-07	2.0E-02	2.0E-09	0%	10950	2.3E-07	0.0E+00	0.0E+00	0%
Chrysene	2.8E-01	350	30	1E-06	100	70	25550	1.6E-07	7.3E-03	1.2E-09	0%	10950	3.8E-07	0.0E+00	0.0E+00	0%
Heptachlor epoxide	6.9E-03	350	30	1E-06	100	70	25550	4.1E-09	9.1E+00	3.7E-08	1%	10950	9.5E-09	1.3E-05	7.3E-04	1%
Indeno(1,2,3-cd)pyrene	2.5E-01	350	30	1E-06	100	70	25550	1.6E-07	7.3E-01	1.1E-07	3%	10950	3.5E-07	0.0E+00	0.0E+00	0%
Phenanthrene	2.8E-01	350	30	1E-06	100	70	25550	1.7E-07	0.0E+00	0.0E+00	0%	10950	3.9E-07	3.0E-02	1.3E-05	0%
alpha-Chlordane	3.6E-02	350	30	1E-06	100	70	25550	2.2E-08	1.3E+00	2.9E-08	1%	10950	6.2E-08	6.0E-05	8.7E-04	1%
gamma-Chlordane	1.3E-02	350	30	1E-06	100	70	25550	7.8E-09	1.3E+00	9.9E-09	0%	10950	1.9E-08	6.0E-05	3.0E-04	0%
TOTAL								3.5E-06						6.2E-02		

SURFACE SOIL DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT RESIDENTIAL ADULT

Dermal contact with soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RID$$

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	
CF = conversion factor (kg/mg)	1E-08
SA = adult exposed skin surface area (cm <sup>2</sup> )	5800
AF = soil to skin adherence factor (mg/cm <sup>2</sup> )	1
Abs = fraction absorbed (unitless)	Specific
EF = adult exposure frequency (events/yr)	350
ED = adult exposure duration (years)	30
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	30
DY = day per year (day/yr)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RID = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> ) Adult	Adherence Factor (mg/cm <sup>2</sup> )	Fraction Absorbed (%)	Exposure Frequency (events/yr)	Exposure Duration (yrs) Adult	body Weight (kg) Adult	Average Carc Time (days)	Carc Dose (mg/kg/day) Adult	Dermal Adjust. Slope Factor (mg/kg-day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (days)	Noncarc/c Dose (mg/kg/day) Adult	Dermal Adjust. Reference Dose (mg/kg/day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	3.4E+03	1E-08	5800	1	0.001	350	30	70	25550	1.2E-08	0.0E+00	0.0E+00	0%	10950	2.7E-14	2.0E-01	1.4E-03	7%
Antimony	5.7E+00	1E-08	5800	1	0.001	350	30	70	25550	1.9E-07	0.0E+00	0.0E+00	0%	10950	4.5E-07	8.0E-05	5.0E-03	28%
Arsenic	1.7E+00	1E-08	5800	1	0.001	350	30	70	25550	6.9E-08	8.8E+00	6.2E-07	21%	10950	1.4E-07	6.0E-05	2.3E-03	11%
Barium	3.5E+01	1E-08	5800	1	0.001	350	30	70	25550	1.2E-08	0.0E+00	0.0E+00	0%	10950	2.8E-08	1.4E-02	2.0E-04	1%
Cadmium (sol)	1.0E+00	1E-08	5800	1	0.001	350	30	70	25550	5.3E-08	0.0E+00	0.0E+00	0%	10950	1.2E-07	2.0E-04	6.2E-04	3%
Chromium	1.0E+01	1E-08	5800	1	0.001	350	30	70	25550	3.5E-07	0.0E+00	0.0E+00	0%	10950	8.2E-07	2.0E-01	4.1E-08	0%
Cobalt	9.1E-01	1E-08	5800	1	0.001	350	30	70	25550	3.1E-08	0.0E+00	0.0E+00	0%	10950	7.2E-08	1.2E-02	6.0E-08	0%
Copper	1.3E+02	1E-08	5800	1	0.001	350	30	70	25550	4.4E-08	0.0E+00	0.0E+00	0%	10950	1.0E-05	7.4E-03	1.4E-03	7%
Lead	1.7E+02	1E-08	5800	1	0.001	350	30	70	25550	8.8E-08	0.0E+00	0.0E+00	0%	10950	1.3E-05	0.0E+00	0.0E+00	0%
Manganese (sol)	3.3E+02	1E-08	5800	1	0.001	350	30	70	25550	1.1E-08	0.0E+00	0.0E+00	0%	10950	2.6E-05	2.8E-02	9.5E-04	5%
Mercury	2.8E-01	1E-08	5800	1	0.001	350	30	70	25550	9.7E-09	0.0E+00	0.0E+00	0%	10950	2.3E-08	6.0E-05	3.8E-04	2%
Nickel	5.6E+00	1E-08	5800	1	0.001	350	30	70	25550	1.0E-07	0.0E+00	0.0E+00	0%	10950	4.4E-07	4.0E-03	1.1E-04	1%
Silver	1.1E+00	1E-08	5800	1	0.001	350	30	70	25550	3.8E-08	0.0E+00	0.0E+00	0%	10950	9.0E-08	1.0E-03	9.0E-05	0%
Thallium	5.1E-01	1E-08	5800	1	0.001	350	30	70	25550	1.7E-08	0.0E+00	0.0E+00	0%	10950	4.1E-08	1.6E-05	2.8E-03	13%
Vanadium	7.9E-01	1E-08	5800	1	0.001	350	30	70	25550	2.7E-07	0.0E+00	0.0E+00	0%	10950	6.3E-07	1.4E-03	4.5E-04	2%
Zinc	1.1E+03	1E-08	5800	1	0.001	350	30	70	25550	3.6E-08	0.0E+00	0.0E+00	0%	10950	8.8E-05	8.0E-02	1.4E-03	7%
4,4'-DDD	6.4E-02	1E-08	5800	1	0.01	350	30	70	25550	2.2E-08	4.8E-01	1.0E-08	0%	10950	5.1E-08	0.0E+00	0.0E+00	0%
4,4'-DDE	3.5E-01	1E-08	5800	1	0.01	350	30	70	25550	1.2E-07	6.8E-01	8.1E-08	3%	10950	2.8E-07	0.0E+00	0.0E+00	0%
4,4'-DDT	1.5E-01	1E-08	5800	1	0.01	350	30	70	25550	5.1E-08	6.8E-01	3.4E-08	1%	10950	1.2E-07	2.5E-04	4.7E-04	2%
Anthracene	2.1E-01	1E-08	5800	1	0.01	350	30	70	25550	7.0E-08	0.0E+00	0.0E+00	0%	10950	1.6E-07	1.8E-01	1.1E-05	0%
Benzo(a)anthracene	2.8E-01	1E-08	5800	1	0.01	350	30	70	25550	9.5E-08	1.5E+00	1.4E-07	6%	10950	2.2E-07	0.0E+00	0.0E+00	0%
Benzo(a)pyrene	2.7E-01	1E-08	5800	1	0.01	350	30	70	25550	9.1E-08	1.5E+01	1.3E-06	54%	10950	2.1E-07	0.0E+00	0.0E+00	0%
Benzo(b)fluoranthene	2.9E-01	1E-08	5800	1	0.01	350	30	70	25550	1.0E-07	1.5E+00	1.5E-07	6%	10950	2.3E-07	0.0E+00	0.0E+00	0%
Benzo(g,h,i)perylene	2.8E-01	1E-08	5800	1	0.01	350	30	70	25550	8.6E-08	0.0E+00	0.0E+00	0%	10950	2.0E-07	1.5E-02	1.3E-05	0%
Carbazole	1.7E-01	1E-08	5800	1	0.01	350	30	70	25550	6.9E-08	4.0E-02	2.3E-09	0%	10950	1.4E-07	0.0E+00	0.0E+00	0%
Chrysene	2.8E-01	1E-08	5800	1	0.01	350	30	70	25550	9.5E-08	1.5E-02	1.4E-09	0%	10950	2.2E-07	0.0E+00	0.0E+00	0%
Heptachlor epoxide	6.9E-03	1E-08	5800	1	0.01	350	30	70	25550	2.4E-09	1.8E+01	4.3E-08	2%	10950	6.5E-09	6.5E-06	8.4E-04	4%
Indeno(1,2,3-cd)pyrene	2.8E-01	1E-08	5800	1	0.01	350	30	70	25550	9.7E-08	0.0E+00	0.0E+00	0%	10950	2.0E-07	0.0E+00	0.0E+00	0%
Phenanthrene	2.8E-01	1E-08	5800	1	0.01	350	30	70	25550	9.7E-08	0.0E+00	0.0E+00	0%	10950	2.3E-07	1.5E-02	1.5E-05	0%
alpha-Chlordane	3.8E-02	1E-08	5800	1	0.01	350	30	70	25550	1.3E-08	2.6E+00	3.4E-08	1%	10950	3.0E-08	3.0E-05	1.0E-03	6%
gamma-Chlordane	1.3E-02	1E-08	5800	1	0.01	350	30	70	25550	4.4E-09	2.6E+00	1.1E-08	0%	10950	1.0E-08	3.4E-04	2%	
TOTAL											2.6E-08					2.0E-02		

SURFACE SOIL PARTICULATE INHALATION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT RESIDENTIAL ADULT

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (C * EF * ED * IR * 1/PEF) / (BW * ATc or ATnc * DY)$$

$$\text{Risk} = \text{Intake} * \text{CSF or}^* / \text{RfD}$$

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	Calculated
CSF = carcinogenic slope factor	Specific
RfD = reference dose for noncarcinogen	Specific
IR = Inhalation rate (m3)	20
EF = adult exposure frequency (days)	350
ED = adult exposure duration (years)	30
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	30
DY = day per year (day/yr)	365
PEF = particulate emission factor (m3/kg)	4.63E+09

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Particulate Emission Factor (m3/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m3/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Car Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncarc Time (days)	NONCarc Dose (mg/kg/day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Aluminum	3.4E+03	4.6E+09	350	20	30	70	25550	8.7E-08	0.0E+00	0.0E+00	0%	10950	2.0E-07	0.0E+00	0.0E+00	0%
Antimony	6.7E+00	4.6E+09	350	20	30	70	25550	1.4E-10	0.0E+00	0.0E+00	0%	10950	3.4E-10	0.0E+00	0.0E+00	0%
Arsenic	1.7E+00	4.6E+09	350	20	30	70	25550	4.4E-11	1.8E+01	0.7E-10	92%	10950	1.0E-10	0.0E+00	0.0E+00	0%
Barium	3.5E-01	4.6E+09	350	20	30	70	25550	8.8E-10	0.0E+00	0.0E+00	0%	10950	2.1E-09	1.4E-04	1.4E-05	99%
Cadmium (soil)	1.6E+00	4.6E+09	350	20	30	70	25550	3.9E-11	0.0E+00	0.0E+00	0%	10950	9.2E-11	0.0E+00	0.0E+00	0%
Chromium	1.0E+01	4.6E+09	350	20	30	70	25550	2.8E-10	0.0E+00	0.0E+00	0%	10950	6.1E-10	0.0E+00	0.0E+00	0%
Cobalt	9.1E-01	4.6E+09	350	20	30	70	25550	2.3E-11	0.0E+00	0.0E+00	0%	10950	5.4E-11	0.0E+00	0.0E+00	0%
Copper	1.3E+02	4.6E+09	350	20	30	70	25550	3.3E-09	0.0E+00	0.0E+00	0%	10950	7.7E-09	0.0E+00	0.0E+00	0%
Lead	1.7E+02	4.6E+09	350	20	30	70	25550	4.3E-09	0.0E+00	0.0E+00	0%	10950	1.0E-08	0.0E+00	0.0E+00	0%
Manganese (soil)	3.3E+02	4.6E+09	350	20	30	70	25550	8.3E-09	0.0E+00	0.0E+00	0%	10950	1.9E-08	0.0E+00	0.0E+00	0%
Mercury	2.9E-01	4.6E+09	350	20	30	70	25550	7.2E-12	0.0E+00	0.0E+00	0%	10950	1.7E-11	8.8E-05	2.0E-07	1%
Nickel	6.6E+00	4.6E+09	350	20	30	70	25550	1.4E-10	0.0E+00	0.0E+00	0%	10950	3.3E-10	0.0E+00	0.0E+00	0%
Silver	1.1E+00	4.6E+09	350	20	30	70	25550	2.9E-11	0.0E+00	0.0E+00	0%	10950	6.7E-11	0.0E+00	0.0E+00	0%
Thallium	6.1E-01	4.6E+09	350	20	30	70	25550	1.3E-11	0.0E+00	0.0E+00	0%	10950	3.0E-11	0.0E+00	0.0E+00	0%
Zinc	7.9E+00	4.6E+09	350	20	30	70	25550	2.0E-10	0.0E+00	0.0E+00	0%	10950	4.7E-10	0.0E+00	0.0E+00	0%
4,4'-DDD	1.1E+03	4.6E+09	350	20	30	70	25550	2.7E-08	0.0E+00	0.0E+00	0%	10950	6.3E-08	0.0E+00	0.0E+00	0%
4,4'-DDE	6.4E-02	4.6E+09	350	20	30	70	25550	1.8E-12	0.0E+00	0.0E+00	0%	10950	3.8E-12	0.0E+00	0.0E+00	0%
4,4'-DDT	3.6E-01	4.6E+09	350	20	30	70	25550	8.8E-12	0.0E+00	0.0E+00	0%	10950	2.1E-11	0.0E+00	0.0E+00	0%
Anthracene	1.5E-01	4.6E+09	350	20	30	70	25550	3.8E-12	3.4E-01	1.3E-12	0%	10950	8.8E-12	0.0E+00	0.0E+00	0%
Benz(a)anthracene	2.9E-01	4.6E+09	350	20	30	70	25550	7.1E-12	6.1E-01	4.3E-12	1%	10950	1.2E-11	0.0E+00	0.0E+00	0%
Benz(a)pyrene	2.7E-01	4.6E+09	350	20	30	70	25550	8.8E-12	6.1E+00	4.1E-11	6%	10950	1.6E-11	0.0E+00	0.0E+00	0%
Benz(b)fluoranthene	2.9E-01	4.6E+09	350	20	30	70	25550	7.5E-12	6.1E-01	4.6E-12	1%	10950	1.7E-11	0.0E+00	0.0E+00	0%
Benz(g,h,i)perylene	2.9E-01	4.6E+09	350	20	30	70	25550	6.4E-12	0.0E+00	0.0E+00	0%	10950	1.5E-11	0.0E+00	0.0E+00	0%
Carbazole	1.7E-01	4.6E+09	350	20	30	70	25550	4.3E-12	0.0E+00	0.0E+00	0%	10950	1.0E-11	0.0E+00	0.0E+00	0%
Chrysene	2.9E-01	4.6E+09	350	20	30	70	25550	7.1E-12	6.1E-03	4.3E-14	0%	10950	1.7E-11	0.0E+00	0.0E+00	0%
Heptachlor epoxide	6.9E-03	4.6E+09	350	20	30	70	25550	1.8E-13	9.1E+00	1.6E-12	0%	10950	4.1E-13	0.0E+00	0.0E+00	0%
Indeno(1,2,3-cd)pyrene	2.8E-01	4.6E+09	350	20	30	70	25550	8.4E-12	6.1E-01	3.9E-12	1%	10950	1.5E-11	0.0E+00	0.0E+00	0%
Phenanthrene	2.9E-01	4.6E+09	350	20	30	70	25550	7.2E-12	0.0E+00	0.0E+00	0%	10950	1.7E-11	0.0E+00	0.0E+00	0%
alpha-Chlordane	3.8E-02	4.6E+09	350	20	30	70	25550	9.7E-13	1.3E+00	1.2E-12	0%	10950	2.3E-12	0.0E+00	0.0E+00	0%
gamma-Chlordane	1.3E-02	4.6E+09	350	20	30	70	25550	3.3E-13	1.3E+00	4.2E-13	0%	10950	7.7E-13	0.0E+00	0.0E+00	0%
TOTAL								7.3E-10						1.6E-05		

SUBSURFACE SOIL INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL ADULT

Intake from ingestion of soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * CF * EF * ED * IR/BW * ATc \text{ or } ATnc * DY$$

$$\text{Risk} = \text{Intake} * \text{CSF} \text{ or } \text{RfD}$$

Where:

	INPUTS	SPECIFIC
C = contaminant concentration in soil (mg/kg)	1E-06	
CF = conversion for kg to mg	1E-06	
EF = adult exposure frequency (days/yr)	360	
ED = adult exposure duration (yr)	30	
IR = adult soil ingestion rate (mg/day)	100	
BW = adult body weight (kg)	70	
ATc = averaging time for carcinogen (yr)	70	
ATnc = averaging time for noncarcinogen (yr)	30	
DY = days per year (days/year)	365	
CSF = cancer slope factor (mg/kg-day)-1	specific	
RfD = reference dose (mg/kg-day)	specific	

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Exposure Frequency (days/yr)	Exposure Duration (yr)	Conversion Factor (kg/mg)	Ingestion Rate (mg/day)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	7.8E+03	360	30	1E-06	100	70	25550	4.8E-03	0.0E+00	0%	10950	1.1E-02	1.0E+00	1.1E-02	5%	
Antimony	1.2E+01	360	30	1E-06	100	70	25550	6.9E-06	0.0E+00	0%	10950	1.6E-05	4.0E-04	4.0E-02	24%	
Arsenic	7.9E+00	360	30	1E-06	100	70	25550	4.6E-06	1.8E+00	81%	10950	1.1E-05	3.0E-04	3.6E-02	22%	
Barium	7.8E+01	360	30	1E-06	100	70	25550	4.8E-05	0.0E+00	0%	10950	1.1E-04	7.0E-02	1.5E-03	1%	
Beryllium	1.4E-01	360	30	1E-06	100	70	25550	8.4E-06	4.3E+00	3%	10950	2.0E-07	6.0E-03	3.9E-05	0%	
Cadmium (soil)	3.2E+00	360	30	1E-06	100	70	25550	1.9E-06	0.0E+00	0%	10950	4.4E-06	1.0E-03	4.4E-03	3%	
Chromium	3.4E+01	360	30	1E-06	100	70	25550	2.0E-05	0.0E+00	0%	10950	4.7E-06	1.0E+00	4.7E-05	0%	
Cobalt	2.7E+00	360	30	1E-06	100	70	25550	1.6E-06	0.0E+00	0%	10950	3.7E-06	6.0E-02	6.1E-05	0%	
Copper	8.9E+02	360	30	1E-06	100	70	25550	5.2E-04	0.0E+00	0%	10950	1.2E-03	3.7E-02	3.3E-02	20%	
Lead	1.7E+03	360	30	1E-06	100	70	25550	1.0E-03	0.0E+00	0%	10950	2.4E-03	0.0E+00	0.0E+00	0%	
Manganese (soil)	1.2E+03	360	30	1E-06	100	70	25550	7.0E-04	0.0E+00	0%	10950	1.6E-03	1.4E-01	1.2E-02	7%	
Mercury	2.3E-01	360	30	1E-06	100	70	25550	1.3E-07	0.0E+00	0%	10950	3.1E-07	3.0E-04	1.0E-03	1%	
Nickel	2.3E+01	360	30	1E-06	100	70	25550	1.4E-05	0.0E+00	0%	10950	3.2E-05	2.0E-02	1.6E-03	1%	
Silver	2.5E+00	360	30	1E-06	100	70	25550	1.5E-06	0.0E+00	0%	10950	3.4E-06	5.0E-03	6.9E-04	0%	
Vanadium	2.4E+01	360	30	1E-06	100	70	25550	1.4E-05	0.0E+00	0%	10950	3.4E-05	7.0E-03	4.8E-03	3%	
Zinc	4.3E+03	360	30	1E-06	100	70	25550	2.5E-03	0.0E+00	0%	10950	5.9E-03	3.0E-01	2.0E-02	12%	
2-Methylnaphthalene	8.9E-02	360	30	1E-06	100	70	25550	5.2E-08	0.0E+00	0%	10950	1.2E-07	4.0E-02	3.0E-03	0%	
4,4'-DDD	8.8E-01	360	30	1E-06	100	70	25550	5.2E-07	2.4E-01	1%	10950	1.2E-06	0.0E+00	0.0E+00	0%	
4,4'-DDE	1.0E+00	360	30	1E-06	100	70	25550	9.4E-07	3.4E-01	2%	10950	2.2E-06	0.0E+00	0.0E+00	0%	
4,4'-DDT	1.2E-01	360	30	1E-06	100	70	25550	6.9E-08	3.4E-01	2.3E-08	0%	10950	1.8E-07	8.0E-04	3.2E-04	0%
Benzo(a)anthracene	5.8E-01	360	30	1E-06	100	70	25550	3.4E-07	7.3E-01	2.5E-07	2%	10950	8.0E-07	0.0E+00	0.0E+00	0%
Benzo(a)pyrene	5.6E-01	360	30	1E-06	100	70	25550	3.3E-07	7.3E+00	2.4E-06	18%	10950	7.7E-07	0.0E+00	0.0E+00	0%
Benzo(b)fluoranthene	6.2E-01	360	30	1E-06	100	70	25550	3.6E-07	7.3E-01	2.6E-07	2%	10950	8.4E-07	0.0E+00	0.0E+00	0%
Benzo(g,h,i)perylene	4.4E-01	360	30	1E-06	100	70	25550	2.6E-07	0.0E+00	0%	10950	6.1E-07	3.0E-02	2.0E-05	0%	
Benzo(k)fluoranthene	5.4E-01	360	30	1E-06	100	70	25550	3.1E-07	7.3E-02	2.3E-08	0%	10950	7.3E-07	0.0E+00	0.0E+00	0%
Bis(2-ethylhexyl)phthalate	4.8E-01	360	30	1E-06	100	70	25550	2.8E-07	1.4E-02	3.9E-08	0%	10950	8.5E-07	2.0E-02	3.3E-05	0%
Chrysene	8.7E-01	360	30	1E-06	100	70	25550	3.4E-07	7.3E-03	2.5E-09	0%	10950	7.9E-07	0.0E+00	0.0E+00	0%
Dibenz(a,h)anthracene	2.8E-01	360	30	1E-06	100	70	25550	1.7E-07	7.3E+00	1.2E-06	9%	10950	3.9E-07	0.0E+00	0.0E+00	0%
Fluorene	2.8E-01	360	30	1E-06	100	70	25550	1.6E-07	0.0E+00	0.0E+00	0%	10950	3.8E-07	4.0E-02	9.8E-06	0%
Indeno(1,2,3-cd)pyrene	4.3E-01	360	30	1E-06	100	70	25550	2.6E-07	7.3E-01	1.9E-07	1%	10950	6.0E-07	0.0E+00	0.0E+00	0%
Naphthalene	3.4E-01	360	30	1E-06	100	70	25550	2.0E-07	0.0E+00	0.0E+00	0%	10950	4.7E-07	4.0E-02	1.2E-05	0%
Phenanthrene	8.4E-01	360	30	1E-06	100	70	25550	3.8E-07	0.0E+00	0.0E+00	0%	10950	8.8E-07	3.0E-02	2.9E-05	0%
alpha-Chlordane	1.2E-02	360	30	1E-06	100	70	25550	5.8E-09	1.3E+00	8.8E-09	0%	10950	1.6E-08	0.0E-05	2.6E-04	0%
gamma-Chlordane	8.7E-03	360	30	1E-06	100	70	25550	3.9E-09	1.3E+00	5.1E-09	0%	10950	9.2E-09	0.0E-05	1.6E-04	0%
TOTAL									1.3E-05					1.7E-01		

SUBSURFACE SOIL DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL ADULT

Dermal contact with soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * CF * SA * AF * Abs * EF * ED/BW * ATc or ATnc * DY$$

$$\text{Risk} = \text{Intake} * \text{CSF or } / \text{RfD}$$

Where:

C = contaminant concentration in soil (mg/kg)

CF = conversion factor (kg/mg)

1E-06

SA = adult exposed skin surface area (cm<sup>2</sup>)

5800

AF = soil to skin adherence factor (mg/cm<sup>2</sup>)

1

Abs = fraction absorbed (unitless)

Specific

EF = adult exposure frequency (events/yr)

350

ED = adult exposure duration (years)

30

BW = adult body weight (kg)

70

ATc = averaging time for carcinogen (yr)

70

ATnc = averaging time for noncarcinogen (yr)

30

DY = day per year (day/yr)

365

CSF = cancer slope factor (mg/kg-day)<sup>-1</sup>

specific

RfD = reference dose (mg/kg-day)

specific

INPUTS

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> )	Adherence Factor (mg/cm <sup>2</sup> )	Fraction Absorbed (%)	Exposure Frequency (events/yr)	Exposure Duration (yrs)	Body Weight (kg)	Average Care Time (days)	C&C Dose (mg/kg/day)	Dermal Adjust Slope Factor	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncare Time (days)	Noncarc Dose (mg/kg/day)	Dermal Adjust Reference Dose (mg/kg-day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	7.8E+03	1E-06	5800	1	0.001	350	30	70	25550	2.7E-04	0.0E+00	0%	10950	6.2E-04	2.0E-01	3.1E-03	5%	
Antimony	1.2E+01	1E-06	5800	1	0.001	350	30	70	25550	4.0E-05	0.0E+00	0%	10950	9.3E-07	8.0E-05	1.2E-02	24%	
Arsenic	7.9E+00	1E-06	5800	1	0.001	350	30	70	25550	2.7E-07	8.8E+00	2.4E-06	28%	10950	6.3E-07	8.0E-05	1.0E-02	21%
Barium	7.8E+01	1E-06	5800	1	0.001	350	30	70	25550	2.7E-06	0.0E+00	0%	10950	6.2E-06	1.4E-02	4.4E-04	1%	
Beryllium	1.4E-01	1E-06	5800	1	0.001	350	30	70	25550	4.9E-09	2.1E+01	1.1E-07	1%	10950	1.1E-06	1.0E-03	1.1E-06	0%
Cadmium (soil)	3.2E+00	1E-06	5800	1	0.001	350	30	70	25550	1.1E-07	0.0E+00	0%	10950	2.8E-07	2.0E-04	1.3E-03	3%	
Chromium	3.4E+01	1E-06	5800	1	0.001	350	30	70	25550	1.2E-06	0.0E+00	0%	10950	2.7E-06	2.0E-01	1.4E-05	0%	
Cobalt	2.7E+00	1E-06	5800	1	0.001	350	30	70	25550	0.2E-08	0.0E+00	0%	10950	2.1E-07	1.2E-02	1.8E-05	0%	
Copper	8.9E+02	1E-06	5800	1	0.001	350	30	70	25550	3.0E-05	0.0E+00	0%	10950	7.0E-05	7.4E-03	8.8E-03	19%	
Lead	1.7E+03	1E-06	5800	1	0.001	350	30	70	25550	5.9E-05	0.0E+00	0%	10950	1.4E-04	0.0E+00	0.0E+00	0%	
Manganese (soil)	1.2E+03	1E-06	5800	1	0.001	350	30	70	25550	4.0E-05	0.0E+00	0%	10950	9.4E-05	2.8E-02	3.4E-03	7%	
Mercury	2.3E-01	1E-06	5800	1	0.001	350	30	70	25550	7.8E-09	0.0E+00	0%	10950	1.8E-06	8.0E-06	3.0E-04	1%	
Nickel	2.3E+01	1E-06	5800	1	0.001	350	30	70	25550	7.9E-07	0.0E+00	0%	10950	1.8E-06	4.0E-03	4.8E-04	1%	
Silver	2.8E+00	1E-06	5800	1	0.001	350	30	70	25550	8.8E-08	0.0E+00	0%	10950	2.0E-07	1.0E-03	2.0E-04	0%	
Titanium	2.4E+01	1E-06	5800	1	0.001	350	30	70	25550	8.9E-07	0.0E+00	0%	10950	1.9E-06	1.4E-03	1.4E-03	3%	
Zinc	4.3E+03	1E-06	5800	1	0.001	350	30	70	25550	1.5E-04	0.0E+00	0%	10950	3.4E-04	6.0E-02	5.7E-03	12%	
2-Methylnaphthalene	8.9E-02	1E-06	5800	1	0.01	350	30	70	25550	3.0E-08	0.0E+00	0%	10950	7.1E-05	2.0E-02	3.8E-06	0%	
1,4-DDD	8.6E-01	1E-06	5800	1	0.01	350	30	70	25550	3.0E-07	4.8E-01	1.4E-07	2%	10950	7.0E-07	0.0E+00	0.0E+00	0%
1,4-DDE	1.6E+00	1E-06	5800	1	0.01	350	30	70	25550	5.4E-07	6.8E-01	3.7E-07	5%	10950	1.3E-06	0.0E+00	0.0E+00	0%
1,4-DDT	1.2E-01	1E-06	5800	1	0.01	350	30	70	25550	4.0E-08	6.8E-01	2.7E-08	0%	10950	9.3E-08	2.8E-04	3.7E-04	1%
3-nitro(4-ethoxy)anthracene	5.8E-01	1E-06	5800	1	0.01	350	30	70	25550	2.0E-07	1.5E+00	2.9E-07	4%	10950	4.6E-07	0.0E+00	0.0E+00	0%
Benzo(a)pyrene	8.6E-01	1E-06	5800	1	0.01	350	30	70	25550	1.9E-07	1.6E+01	2.8E-06	35%	10950	4.5E-07	0.0E+00	0.0E+00	0%
Benzo(b)fluoranthene	0.2E-01	1E-06	5800	1	0.01	350	30	70	25550	2.1E-07	1.5E+00	3.1E-07	4%	10950	4.9E-07	0.0E+00	0.0E+00	0%
Benzo(a,h)perylene	4.4E-01	1E-06	5800	1	0.01	350	30	70	25550	1.5E-07	0.0E+00	0%	10950	3.5E-07	1.5E-02	2.4E-05	0%	
Benzo(f)fluoranthene	5.4E-01	1E-06	5800	1	0.01	350	30	70	25550	1.6E-07	1.5E-01	2.7E-08	0%	10950	4.3E-07	0.0E+00	0.0E+00	0%
3is(2-ethylhexyl)phthalate	4.8E-01	1E-06	5800	1	0.01	350	30	70	25550	1.0E-07	2.8E-02	4.6E-09	0%	10950	3.8E-07	1.0E-02	3.8E-05	0%
Chrysene	8.7E-01	1E-06	5800	1	0.01	350	30	70	25550	2.0E-07	1.5E-02	2.9E-09	0%	10950	4.6E-07	0.0E+00	0.0E+00	0%
Dibenz(a,h)anthracene	2.8E-01	1E-06	5800	1	0.01	350	30	70	25550	9.6E-08	1.5E+01	1.4E-08	17%	10950	2.2E-07	0.0E+00	0.0E+00	0%
Fluorene	2.8E-01	1E-06	5800	1	0.01	350	30	70	25550	9.5E-08	0.0E+00	0%	10950	2.2E-07	2.0E-02	1.1E-06	0%	
Indeno(1,2,3-cd)pyrene	4.3E-01	1E-06	5800	1	0.01	350	30	70	25550	1.5E-07	1.5E+00	2.2E-07	3%	10950	3.5E-07	0.0E+00	0.0E+00	0%
Naphthalene	3.4E-01	1E-06	5800	1	0.01	350	30	70	25550	1.2E-07	0.0E+00	0%	10950	2.7E-07	2.0E-02	1.4E-05	0%	
Phenanthrene	6.4E-01	1E-06	5800	1	0.01	350	30	70	25550	2.2E-07	0.0E+00	0%	10950	5.1E-07	1.5E-02	3.4E-05	0%	
Upha-Chlordane	1.2E-02	1E-06	5800	1	0.01	350	30	70	25550	3.9E-09	2.6E+00	1.0E-08	0%	10950	9.2E-09	3.0E-05	3.1E-04	1%
gamma-Chlordane	6.7E-03	1E-06	5800	1	0.01	350	30	70	25550	2.3E-09	2.6E+00	5.9E-09	0%	10950	5.3E-09	3.0E-05	1.8E-04	0%
TOTAL														4.1E-06				
																		4.3E-02

SUBSURFACE SOIL PARTICLES & INHALATION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL ADULT

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (C * EF * ED * IR * 1/PEF) / (BW * ATc or ATnc * DY)$$

$$\text{Risk} = \text{Intake} * \text{CSF or } / \text{RD}$$

Where:

C = contaminant concentration in soil (mg/kg)  
 Specific  
 CSF = carcinogenic slope factor  
 Specific  
 RD = reference dose for noncarcinogen  
 Specific  
 IR = inhalation rate (m³)  
 20  
 EF = adult exposure frequency (days)  
 350  
 ED = adult exposure duration (years)  
 30  
 BW = adult body weight (kg)  
 70  
 ATc = averaging time for carcinogen (yr)  
 70  
 ATnc = averaging time for noncarcinogen (yr)  
 30  
 DY = day per year (day/yr)  
 365  
 PEF = particulate emission factor (m³/kg)  
 4.63E+09

Note: Inputs are scenario and site specific

COPC	Concentration Carcinogen (mg/kg)	Particulate Emission Factor (m³/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m³/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (days)	CARC Dose (mg/kg/day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Aluminum	7.8E+03	4.6E+09	350	20	30	70	25550	2.0E-07	0.0E+00	0%	10950	4.6E-07	0.0E+00	0.0E+00	0%	
Antimony	1.2E+01	4.6E+09	350	20	30	70	25550	3.0E-10	0.0E+00	0%	10950	7.0E-10	0.0E+00	0.0E+00	0%	
Arsenic	7.9E+00	4.6E+09	350	20	30	70	25550	2.0E-10	1.5E+01	3.0E-09	94%	10950	4.7E-10	0.0E+00	0.0E+00	0%
Barium	7.8E+01	4.6E+09	350	20	30	70	25550	2.0E-09	0.0E+00	0%	10950	4.6E-09	1.4E-04	3.2E-06	100%	
Beryllium	1.4E+01	4.6E+09	350	20	30	70	25550	3.6E-12	8.4E+00	3.1E-11	1%	10950	8.5E-12	0.0E+00	0.0E+00	0%
Cadmium (soil)	3.2E+00	4.6E+09	350	20	30	70	25550	8.1E-11	0.0E+00	0%	10950	1.9E-10	0.0E+00	0.0E+00	0%	
Chromium	3.4E+01	4.6E+09	350	20	30	70	25550	8.7E-10	0.0E+00	0%	10950	2.0E-09	0.0E+00	0.0E+00	0%	
Cobalt	2.7E+00	4.6E+09	350	20	30	70	25550	6.8E-11	0.0E+00	0%	10950	1.6E-10	0.0E+00	0.0E+00	0%	
Copper	8.9E+02	4.6E+09	350	20	30	70	25550	2.2E-08	0.0E+00	0%	10950	5.2E-08	0.0E+00	0.0E+00	0%	
Lead	1.7E+03	4.6E+09	350	20	30	70	25550	4.4E-08	0.0E+00	0%	10950	1.0E-07	0.0E+00	0.0E+00	0%	
Manganese (soil)	1.2E+03	4.6E+09	350	20	30	70	25550	3.0E-08	0.0E+00	0%	10950	7.0E-08	0.0E+00	0.0E+00	0%	
Mercury	2.3E-01	4.6E+09	350	20	30	70	25550	5.8E-12	0.0E+00	0%	10950	1.3E-11	0.0E-05	1.6E-07	0%	
Nickel	2.3E+01	4.6E+09	350	20	30	70	25550	5.9E-10	0.0E+00	0%	10950	1.4E-09	0.0E+00	0.0E+00	0%	
Silver	2.5E+00	4.6E+09	350	20	30	70	25550	0.3E-11	0.0E+00	0%	10950	1.5E-10	0.0E+00	0.0E+00	0%	
Vanadium	2.4E+01	4.6E+09	350	20	30	70	25550	6.2E-10	0.0E+00	0%	10950	1.4E-09	0.0E+00	0.0E+00	0%	
Zinc	4.3E+03	4.6E+09	350	20	30	70	25550	1.1E-07	0.0E+00	0%	10950	2.6E-07	0.0E+00	0.0E+00	0%	
2-Methylnaphthalene	8.9E-02	4.6E+09	350	20	30	70	25550	2.3E-12	0.0E+00	0%	10950	5.3E-12	0.0E+00	0.0E+00	0%	
4,4'-DDD	8.8E-01	4.6E+09	350	20	30	70	25550	2.2E-11	0.0E+00	0%	10950	5.2E-11	0.0E+00	0.0E+00	0%	
4,4'-DDE	1.6E+00	4.6E+09	350	20	30	70	25550	4.1E-11	0.0E+00	0%	10950	9.5E-11	0.0E+00	0.0E+00	0%	
4,4'-DDT	1.2E-01	4.6E+09	350	20	30	70	25550	3.0E-12	3.4E-01	1.0E-12	0%	10950	6.9E-12	0.0E+00	0.0E+00	0%
Benzo(a)anthracene	5.8E-01	4.6E+09	350	20	30	70	25550	1.5E-11	6.1E-01	9.0E-12	0%	10950	3.4E-11	0.0E+00	0.0E+00	0%
Benzo(a)pyrene	5.6E-01	4.6E+09	350	20	30	70	25550	1.4E-11	6.1E+00	8.7E-11	3%	10950	3.3E-11	0.0E+00	0.0E+00	0%
Benzo(b)fluoranthene	6.2E-01	4.6E+09	360	20	30	70	25550	1.6E-11	6.1E-01	9.5E-12	0%	10950	3.6E-11	0.0E+00	0.0E+00	0%
Benzo(g,h,i)perylene	4.4E-01	4.6E+09	350	20	30	70	25550	1.1E-11	0.0E+00	0%	10950	2.6E-11	0.0E+00	0.0E+00	0%	
Benzo(k)fluoranthene	5.4E-01	4.6E+09	350	20	30	70	25550	1.4E-11	6.1E-02	8.3E-13	0%	10950	3.2E-11	0.0E+00	0.0E+00	0%
Bis(2-ethylhexyl)phthalate	4.8E-01	4.6E+09	350	20	30	70	25550	1.2E-11	0.0E+00	0%	10950	2.9E-11	0.0E+00	0.0E+00	0%	
Chrysene	5.7E-01	4.6E+09	350	20	30	70	25550	1.5E-11	6.1E-03	8.9E-14	0%	10950	3.4E-11	0.0E+00	0.0E+00	0%
Dibenz(a,h)anthracene	2.8E-01	4.6E+09	350	20	30	70	25550	7.2E-12	6.1E+00	4.4E-11	1%	10950	1.7E-11	0.0E+00	0.0E+00	0%
Fluorene	2.8E-01	4.6E+09	350	20	30	70	25550	7.1E-12	0.0E+00	0%	10950	1.7E-11	0.0E+00	0.0E+00	0%	
Indeno(1,2,3-cd)pyrene	4.3E-01	4.6E+09	350	20	30	70	25550	1.1E-11	6.1E-01	6.7E-12	0%	10950	2.6E-11	0.0E+00	0.0E+00	0%
Naphthalene	3.4E-01	4.6E+09	350	20	30	70	25550	8.7E-12	0.0E+00	0%	10950	2.0E-11	0.0E+00	0.0E+00	0%	
Phenanthrene	6.4E-01	4.6E+09	350	20	30	70	25550	1.6E-11	0.0E+00	0.0E+00	0%	10950	3.8E-11	0.0E+00	0.0E+00	0%
alpha-Chlordane	1.2E-02	4.6E+09	350	20	30	70	25550	2.9E-13	1.3E+00	3.8E-13	0%	10950	6.8E-13	0.0E+00	0.0E+00	0%
gamma-Chlordane	6.7E-03	4.6E+09	350	20	30	70	25550	1.7E-13	1.3E+00	2.2E-13	0%	10950	4.0E-13	0.0E+00	0.0E+00	0%
<b>TOTAL</b>												3.2E-09				3.3E-05

**ASSESSMENT**  
**OPERABLE UNIT NO. 7 (SITE 2B - NEW RIVER AND COODLEB CREEK)**  
**REMEDIATION INVESTIGATION CTO-0201**  
**MCB CAMP LEJEUNE, NORTH CAROLINA**  
**CURRENT AND FUTURE RESIDENTIAL ADULT**

Intake from ingestion of sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times R \times CF \times EF \times ED \times BW \times ATC \text{ or } ATnc \times DY$$

CHANGE

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RD$$

Where:

**INPUTS**

C = contaminant concentration in sediment (mg/kg)	
CF = conversion for kg to mg	1E-06
EF = exposure frequency (days/yr)	45
ED = exposure duration (yr)	.30
R = soil ingestion rate (mg/day)	100
BW = body weight (kg)	70
ATC = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	50
DY = days per year (days/year)	365
CSF = cancer slope factor (mg/kg-day)-1	Specific
RD = reference dose (mg/kg-day)	Specific

Note: Inputs are scenario and site specific

**NEW RIVER**

POPC	Concentration (mg/kg)	Exposure Frequency (days/yr)	Exposure Duration (yr)	Ingestion Rate (mg/day)	Conversion Factor (kg/mg)	Body Weight (kg)	Average Cnic Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Acetone	2.0E+02	45	.30	100	1E-06	70	25500	2.0E-05	1.0E+00	0%	10500	4.0E-05	4.0E-04	1.0E-01	0.1%	
Asenic	9.0E+00	45	.30	100	1E-06	70	25500	9.0E-07	1.0E+00	1.0E-08	10500	1.0E-08	3.0E-04	6.0E-03	4%	
Benzen	1.0E+01	45	.30	100	1E-06	70	25500	1.0E-06	0.0E+00	0%	10500	2.7E-08	7.0E-02	3.0E-05	0%	
Copper	1.0E+03	45	.30	100	1E-06	70	25500	1.0E-04	0.0E+00	0%	10500	2.4E-04	3.7E-02	8.0E-03	0%	
Lead	3.0E+04	45	.30	100	1E-06	70	25500	2.0E-03	0.0E+00	0%	10500	6.0E-03	0.0E+00	0.0E+00	0%	
Zinc	2.1E+03	45	.30	100	1E-06	70	25500	1.0E-07	0.0E+00	0%	10500	3.0E-07	8.0E-03	7.0E-05	0%	
1,4-DOD	1.0E-02	45	.30	100	1E-06	70	25500	1.1E-09	2.4E-01	2.7E-10	0%	10500	2.0E-09	0.0E+00	0.0E+00	0%
1,4-DDC	0.0E+00	45	.30	100	1E-06	70	25500	0.0E+00	3.0E-10	3.0E-10	0%	10500	1.0E-08	0.0E+00	0.0E+00	0%
1,4-DDT	3.0E-01	45	.30	100	1E-06	70	25500	2.0E-08	3.0E-01	7.7E-08	0%	10500	6.0E-08	6.0E-04	1.1E-04	0%
1,4-Trich	2.0E-01	45	.30	100	1E-06	70	25500	2.1E-08	0.0E+00	0%	10500	4.0E-08	3.0E-01	1.0E-07	0%	
1,4-Di(4-ethoxy)benzene	1.0E+00	45	.30	100	1E-06	70	25500	0.0E+00	7.0E-01	6.0E-08	4%	10500	2.1E-07	0.0E+00	0.0E+00	0%
1,4-Di(4-pyridyl)pyrene	0.0E+00	45	.30	100	1E-06	70	25500	4.0E-08	7.0E+00	3.1E-07	10%	10500	9.0E-08	0.0E+00	0.0E+00	0%
1,4-Di(4-fluorophenyl)anthracene	1.0E+00	45	.30	100	1E-06	70	25500	0.0E+00	7.0E-01	8.0E-08	4%	10500	1.0E-07	0.0E+00	0.0E+00	0%
1,4-Di(4-phenyl)benzene	3.0E-01	45	.30	100	1E-06	70	25500	2.0E-08	0.0E+00	0%	10500	5.0E-08	3.0E-02	1.0E-08	0%	
1,4-Di(4-fluorophenyl)benzene	4.0E-01	45	.30	100	1E-06	70	25500	3.7E-08	7.0E-02	2.7E-08	0%	10500	8.0E-08	0.0E+00	0.0E+00	0%
2-Ethylhexylphthalate	1.0E+00	45	.30	100	1E-06	70	25500	0.0E+00	1.0E-02	1.0E-08	0%	10500	2.0E-07	2.0E-02	1.1E-05	0%
Carbazole	1.0E-01	45	.30	100	1E-06	70	25500	1.0E-08	2.0E-02	2.4E-10	0%	10500	2.0E-08	0.0E+00	0.0E+00	0%
Chrysene	1.0E+00	45	.30	100	1E-06	70	25500	1.0E-07	7.0E-03	8.0E-10	0%	10500	2.7E-07	0.0E+00	0.0E+00	0%
Obenzotriphen	0.0E+00	45	.30	100	1E-06	70	25500	4.0E-09	0.0E+00	0%	10500	1.0E-08	0.0E+00	0.0E+00	0%	
Toluene	1.0E+00	45	.30	100	1E-06	70	25500	0.0E+00	0.0E+00	0%	10500	2.0E-07	4.0E-02	6.0E-08	0%	
nitro(1,2-diphenyl)pyrene	3.0E-01	45	.30	100	1E-06	70	25500	2.0E-08	7.0E-01	1.7E-08	1%	10500	5.0E-08	0.0E+00	0.0E+00	0%
Thiophene	7.7E-01	45	.30	100	1E-06	70	25500	8.0E-08	0.0E+00	0%	10500	1.4E-07	3.0E-02	4.5E-05	0%	
Yrene	1.0E+00	45	.30	100	1E-06	70	25500	0.0E+00	0.0E+00	0%	10500	1.0E-07	3.0E-02	8.0E-05	0%	
alpha-Chordane	0.0E+00	45	.30	100	1E-06	70	25500	0.0E+00	1.0E-00	0.0E-10	0%	10500	1.0E-08	0.0E-05	1.0E-05	0%
gamma-Chordane	4.0E-03	45	.30	100	1E-06	70	25500	3.0E-10	1.0E+00	4.0E-10	0%	10500	8.0E-10	0.0E-05	1.4E-05	0%
UATL									1.7E-05					1.0E-01		

SEDIMENT DERMAL CONTACT ASSESSMENT  
OPERABLE UNIT NO. 7 SITE 2B: NEW RIVER AND COODLES CREEK  
REMEDIAL INVESTIGATION CTI-0291  
MCB CAMP LEJEUNE, NORTH CAROLINA  
CURRENT AND FUTURE RESIDENTIAL ADULT

The intake from dermal contact to sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \text{ or ATnc} \times DY$$

Risk = Intake / CSF or RfD

Where:

C = contaminant concentration in soil (mg/kg)	
CF = conversion factor (kg/mg)	1.0E-08
SA = exposed skin surface area (cm <sup>2</sup> )	6800
AF = sediment to skin adherence factor (mg/cm <sup>2</sup> )	1
Abs = fraction absorbed (unitsless) (contaminant specific)	
EF = exposure frequency (events/y)	45
ED = exposure duration (years)	.33
BW = body weight (kg)	70
ATc = averaging time for carcinogen (y)	.33
ATnc = averaging time for noncarcinogen (y)	.33
DY = day per year (day/yr)	365
CSF = cancer slope factor (mg/kg-day) <sup>-1</sup>	Specific
RfD = reference dose (mg/kg-day)	Specific

Note: Inputs are scenario and site specific

NEW RIVER

COPC	Concentration (mg/kg)	Conversion Factor (kg/mg)	Sediment Area (cm <sup>2</sup> )	Adherence Factor (mg/cm <sup>2</sup> )	Abs Factor (%)	Exposure Frequency (events/y)	Exposure Duration (yrs)	Body Weight (kg)	Average Carb Time	Carc. Dose (mg/kg/day)	Dermed Adjust Slope Factor (mg/kg-day) <sup>-1</sup>	Carcinogenic Risk	Inherent Carcinogenic Risk	Averaging Noncarb Time (days)	Noncarb Dose (mg/kg/day)	Dermed Adjust Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Noncarcinogenic Risk
anthracene	2.0E+02	1E-08	6800	1	0.01	45	.33	70	25500	1.2E-06	3.4E-10	0%	0.000	2.1E-06	0.0E+00	2.1E-02	0%	
Asentic	1.0E+03	1E-08	6800	1	0.01	45	.33	70	25500	3.6E-06	8.6E-07	30%	0.050	6.2E-06	1.6E-03	1.6E-03	4%	
Barium	1.0E+03	1E-08	6800	1	0.01	45	.33	70	25500	6.0E-06	0.0E+00	0%	0.050	1.6E-07	1.1E-05	0%		
Copper	1.0E+03	1E-08	6800	1	0.01	45	.33	70	25500	8.0E-06	0.0E+00	0%	0.050	1.4E-06	1.6E-03	1.6E-03	5%	
Lead	3.0E+04	1E-08	6800	1	0.01	45	.33	70	25500	1.7E-04	0.0E+00	0%	0.050	4.0E-04	0.0E+00	0.0E+00	0%	
Mercury	2.1E+03	1E-08	6800	1	0.01	45	.33	70	25500	8.4E-06	0.0E+00	0%	0.050	2.2E-06	1.0E-03	2.2E-05	0%	
Phos	6.0E+02	1E-08	6800	1	0.01	45	.33	70	25500	3.6E-07	0.0E+00	0%	0.050	0.0E+00	6.0E-09	1.6E-06	0%	
1,4-EDD	1.0E+02	1E-08	6800	1	0.01	45	.33	70	25500	6.0E-10	4.6E-01	0%	0.050	1.6E-09	0.0E+00	0.0E+00	0%	
1,4-EDE	6.0E+02	1E-08	6800	1	0.01	45	.33	70	25500	3.7E-10	4.6E-01	0%	0.050	8.7E-10	0.0E+00	0.0E+00	0%	
1,4-EDO	3.0E+03	1E-08	6800	1	0.01	45	.33	70	25500	1.2E-08	8.6E-01	1%	0.050	3.1E-08	2.6E-04	1.2E-04	0%	
1,4-EDP	3.0E+01	1E-08	6800	1	0.01	45	.33	70	25500	0.0E+00	0.0E+00	0%	0.050	2.6E-08	1.2E-01	1.6E-07	0%	
1,4-EDT	2.0E+01	1E-08	6800	1	0.01	45	.33	70	25500	6.1E-08	7.6E-09	0%	0.050	1.2E-07	0.0E+00	0.0E+00	0%	
benzene	1.0E+00	1E-08	6800	1	0.01	45	.33	70	25500	2.6E-08	1.6E-01	41%	0.050	5.6E-08	0.0E+00	0.0E+00	0%	
benzo(a)anthracene	6.0E-01	1E-08	6800	1	0.01	45	.33	70	25500	4.6E-08	7.0E-08	0%	0.050	1.1E-07	0.0E+00	0.0E+00	0%	
benzo(a)anthracene	1.1E+00	1E-08	6800	1	0.01	45	.33	70	25500	1.2E-08	0.0E+00	0%	0.050	3.1E-08	1.6E-02	2.0E-08	0%	
benzo(a)anthracene	1.1E+00	1E-08	6800	1	0.01	45	.33	70	25500	1.2E-08	0.0E+00	0%	0.050	3.1E-08	1.6E-02	2.0E-08	0%	
benzo(a)anthracene	3.0E-01	1E-08	6800	1	0.01	45	.33	70	25500	2.1E-08	1.9E-01	0%	0.050	5.1E-08	0.0E+00	0.0E+00	0%	
benzo(a)anthracene	4.0E-01	1E-08	6800	1	0.01	45	.33	70	25500	2.1E-08	3.1E-09	0%	0.050	6.0E-08	0.0E+00	0.0E+00	0%	
benzo(a)anthracene	4.0E-01	1E-08	6800	1	0.01	45	.33	70	25500	2.1E-08	3.1E-09	0%	0.050	6.0E-08	0.0E+00	0.0E+00	0%	
3-(4-chlorophenyl)propanoate	1.0E+00	1E-08	6800	1	0.01	45	.33	70	25500	7.0E-09	4.0E-10	0%	0.050	1.9E-08	0.0E+00	0.0E+00	0%	
2-ethoxyethane	1.0E-01	1E-08	6800	1	0.01	45	.33	70	25500	0.0E+00	0.0E+00	0%	0.050	1.0E-07	0.0E+00	0.0E+00	0%	
Dimethyl	1.0E+00	1E-08	6800	1	0.01	45	.33	70	25500	6.0E-08	1.6E-02	0%	0.050	1.6E-07	0.0E+00	0.0E+00	0%	
Dimethyl	1.0E+00	1E-08	6800	1	0.01	45	.33	70	25500	6.0E-08	1.6E-02	0%	0.050	1.6E-07	0.0E+00	0.0E+00	0%	
Dimethyl	6.0E-02	1E-08	6800	1	0.01	45	.33	70	25500	2.0E-09	0.0E+00	0%	0.050	5.1E-09	0.0E+00	0.0E+00	0%	
Fluoranthene	1.0E+00	1E-08	6800	1	0.01	45	.33	70	25500	8.0E-08	0.0E+00	0%	0.050	2.0E-07	2.0E-02	8.4E-08	0%	
hexachlorobutadiene	3.0E-01	1E-08	6800	1	0.01	45	.33	70	25500	1.3E-08	1.0E-08	2%	0.050	3.1E-08	0.0E+00	0.0E+00	0%	
Thiophene	7.7E-01	1E-08	6800	1	0.01	45	.33	70	25500	3.4E-08	0.0E+00	0%	0.050	7.9E-08	1.6E-02	6.3E-08	0%	
Tyrene	1.1E+00	1E-08	6800	1	0.01	45	.33	70	25500	4.8E-08	0.0E+00	0%	0.050	1.1E-07	1.6E-02	7.4E-08	0%	
alpha-Chordane	6.0E-03	1E-08	6800	1	0.01	45	.33	70	25500	2.9E-10	7.6E-10	0%	0.050	6.7E-10	3.0E-05	2.2E-05	0%	
gamma-Chordane	4.0E-03	1E-08	6800	1	0.01	45	.33	70	25500	5.2E-10	5.2E-10	0%	0.050	4.7E-10	3.0E-05	1.6E-05	0%	
UFC										6.0E-07					3.7E-02			

SEDIMENT INGESTION EXPOSURE ASSESSMENT  
OPERABLE UNIT NO. 7 SITE 2B - NEW RIVER AND COOGLES CREEK  
REMEDIAL INVESTIGATION CTD-021  
MCB CAMP LEUELNE, NORTH CAROLINA  
CURRENT AND FUTURE RESIDENTIAL ADULT

Intake from ingestion of sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * IR * CF * EF * ED / BW * ATc \text{ or } ATne * DY$$

CHANGE

$$\text{Risk} = \text{Intake} * CSF \text{ or } RD^*$$

Where:

C = contaminant concentration in sediment (mg/kg)

CF = conversion factor for kg to mg

EF = exposure frequency (days/yr)

ED = exposure duration (y)

IR = soil ingestion rate (mg/day)

BW = body weight (kg)

ATc = averaging time for carcinogen (yr)

ATne = averaging time for noncarcinogen (yr)

DY = days per year (days/year)

CSF = cancer slope factor (mg/kg-day)-1

RD = reference dose (mg/kg-day)

INPUTS

1E-08

45

30

100

1E-08

70

25500

1.0E+03

0.0E+00

0.0E+00

0%

1.0E-05

7.0E-02

1.0E-04

4.0E-08

1.0E+00

2.4E-04

1%

1.0E-05

1.4E-01

2.0E-04

6.0E-03

3.0E-03

0%

1.0E-05

6.0E-05

0.0E+00

0%

1.0E-05

3.0E-01

1.0E-04

0.0E+00

0%

1.0E-05

7.0E-08

0.0E+00

0%

1.0E-05

0.0E+00

0%

1.0E-05

3.0E-01

1.0E-04

0.0E+00

0%

1.0E-05

0.0E+00

0%

1.0E-05

3.0E-01

1.0E-04

0.0E+00

0%

1.0E-05

7.0E-08

0.0E+00

0%

1.0E-05

&lt;p

SEDIMENT/DERMAL CONTACT ASSESSMENT  
OPERABLE UNIT NO.7 (SITE 2B - NEW RIVER AND COOGELS CREEK)  
REMEDIAl INVESTIGATION CTO-0201  
MCB CAMP LEJEUNE, NORTH CAROLINA  
CURRENT AND FUTURE RESIDENTIAL ADULT

The intake from dermal contact to sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times EDBW \times ATc \text{ or } ATnc \times DY$$

Risk = Intake  $\times$  CSF or /RfD

Where:

C = contaminant concentration in soil (mg/kg)

CF = conversion factor (kg/mg)

SA = exposed skin surface area (cm<sup>2</sup>)

AF = sediment to skin adherence factor (mg/cm<sup>2</sup>)

Abs = fraction absorbed (unless) (contaminant specific)

EF = exposure frequency (events/yr)

ED = exposure duration (years)

BW = body weight (kg)

ATc = averaging time for carcinogen (yr)

ATnc = averaging time for noncarcinogen (yr)

DY = day per year (day/yr)

CSF = cancer slope factor (mg/kg-day)-1

RfD = reference dose (mg/kg-day)

INPUTS

Specific

Note: Inputs are scenario and site specific

COOGELS CREEK

Contaminant	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> )	Adherence Factor (mg/cm <sup>2</sup> )	AFS Factor (%)	Exposure Frequency (events/yr)	Exposure Duration (yr)	Body Weight (kg)	Average Carb Time (days)	Carb Dose (mg/kg/day)	Carb Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carcinogenic	Average Noncarb Time (days)	Noncarb Dose (mg/kg/day)	Carb Adjust Reference Dose (mg/kg-day)	Noncarb Adjust Reference Dose (mg/kg-day)	Percent Noncarbogenic Risk	Percent Noncarbogenic Risk
Antimony	2.0E+04	1E-08	6000	1	0.001	45	30	70	25000	7.1E-05	0.0E+00	0.0E+00	0%	10000	2.0E-04	2.0E-01	1.0E-03	20%	
Asbestos	7.4E+00	1E-08	6000	1	0.001	45	30	70	25000	3.2E-08	2.0E-07	2.0E-07	2.0E-07	10000	7.0E-08	0.0E-05	1.0E-03	25%	
Boron	5.9E+01	1E-08	6000	1	0.001	45	30	70	25000	2.0E-07	0.0E+00	0.0E+00	0%	10000	6.1E-07	1.4E-02	4.0E-05	1%	
Chromium	2.9E+01	1E-08	6000	1	0.001	45	30	70	25000	1.2E-07	0.0E+00	0.0E+00	0%	10000	2.0E-07	2.0E-01	1.4E-03	0%	
Copper	5.1E+01	1E-08	6000	1	0.001	45	30	70	25000	2.2E-07	0.0E+00	0.0E+00	0%	10000	5.2E-07	7.4E-03	7.0E-05	1%	
Lead	1.6E+02	1E-08	6000	1	0.001	45	30	70	25000	7.0E-07	0.0E+00	0.0E+00	0%	10000	1.0E-08	0.0E+00	0.0E+00	0%	
Manganese (soil)	2.5E+02	1E-08	6000	1	0.001	45	30	70	25000	9.0E-07	0.0E+00	0.0E+00	0%	10000	2.3E-06	2.0E-02	8.2E-05	2%	
Mercury	3.0E+01	1E-08	6000	1	0.001	45	30	70	25000	1.7E-08	0.0E+00	0.0E+00	0%	10000	3.0E-08	0.0E-05	0.4E-05	1%	
Thallium	2.0E+00	1E-08	6000	1	0.001	45	30	70	25000	1.0E-08	0.0E+00	0.0E+00	0%	10000	2.0E-08	1.0E-05	1.0E-03	37%	
Vanadium	3.0E+01	1E-08	6000	1	0.001	45	30	70	25000	1.0E-07	0.0E+00	0.0E+00	0%	10000	3.7E-07	1.4E-03	2.0E-04	5%	
Zinc	1.7E+02	1E-08	6000	1	0.001	45	30	70	25000	7.0E-07	0.0E+00	0.0E+00	0%	10000	1.0E-06	0.0E-02	3.0E-06	1%	
1,3-Dichlorobenzene	4.1E-01	1E-08	6000	1	0.01	45	30	70	25000	1.0E-08	0.0E-01	1.0E-08	2%	10000	4.2E-08	0.0E+00	0.0E+00	0%	
1,4-DCDE	4.0E-01	1E-08	6000	1	0.01	45	30	70	25000	2.0E-08	4.0E-01	0.0E-09	1%	10000	4.0E-08	0.0E+00	0.0E+00	0%	
Benzo(a)anthracene	1.0E-01	1E-08	6000	1	0.01	45	30	70	25000	7.0E-09	0.0E-01	0.0E-09	0%	10000	1.0E-08	0.0E+00	0.0E+00	0%	
Benzo(a)pyrene	1.4E-01	1E-08	6000	1	0.01	45	30	70	25000	6.1E-09	1.0E+00	0.0E-09	1%	10000	1.4E-08	0.0E+00	0.0E+00	0%	
Benzo(b)fluoranthene	1.1E-01	1E-08	6000	1	0.01	45	30	70	25000	4.7E-08	1.0E+00	0.0E-07	88%	10000	1.1E-07	0.0E+00	0.0E+00	0%	
Benzo(k)fluoranthene	0.9E-02	1E-08	6000	1	0.01	45	30	70	25000	2.0E-08	1.0E+00	0.0E-08	0%	10000	6.4E-09	0.0E+00	0.0E+00	0%	
Styrene	4.0E-02	1E-08	6000	1	0.01	45	30	70	25000	1.0E-08	1.0E-01	2.7E-10	0%	10000	4.0E-09	0.0E+00	0.0E+00	0%	
Styrene-2-ethylhexylphthalate	7.0E-01	1E-08	6000	1	0.01	45	30	70	25000	3.1E-08	2.0E-02	0.0E-10	0%	10000	7.2E-08	1.0E-02	7.2E-08	0%	
Carbon Disulfide	1.3E-02	1E-08	6000	1	0.01	45	30	70	25000	5.7E-10	0.0E+00	0.0E-08	0%	10000	1.3E-09	0.0E-02	1.7E-08	0%	
Chrysene	1.0E-01	1E-08	6000	1	0.01	45	30	70	25000	7.0E-09	1.0E-02	1.0E-10	0%	10000	1.0E-08	0.0E+00	0.0E+00	0%	
Fluoranthene	3.4E-01	1E-08	6000	1	0.01	45	30	70	25000	1.0E-08	0.0E+00	0.0E+00	0%	10000	3.0E-08	2.0E-02	1.7E-08	0%	
Fluoranthene	2.0E-01	1E-08	6000	1	0.01	45	30	70	25000	1.1E-08	0.0E+00	0.0E+00	0%	10000	2.7E-08	1.0E-02	1.8E-08	0%	
Tyrene	2.0E-01	1E-08	6000	1	0.01	45	30	70	25000	1.1E-08	0.0E+00	0.0E+00	0%	10000	2.0E-08	1.0E-02	1.7E-08	0%	
alpha-Chlordane	5.0E-03	1E-08	6000	1	0.01	45	30	70	25000	2.0E-10	0.0E+00	0.0E-10	0%	10000	6.0E-10	3.0E-05	3.0E-05	0%	
gamma-Chlordane	8.0E-03	1E-08	6000	1	0.01	45	30	70	25000	3.7E-10	0.0E+00	0.0E-10	0%	10000	8.0E-10	3.0E-05	3.0E-05	1%	
OTAL										1.0E-03						3.0E-03			

SURFACE WATER INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO.7 (SITE 28 - NEW RIVER AND COOGELS CREEK)  
 REMEDIAL INVESTIGATION CTD-0221  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT AND FUTURE ADULT RESIDENT

The intake from the ingestion of surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = \text{Cw} * \text{CR} * \text{ET} * \text{EF} * \text{ED/BW} * \text{ATc or ATnc} * \text{DY}$$

$$\text{Risk} = \text{Intake} * \text{CSF or RfD}$$

CHANGE!!

Where:	INPUT
Cw = contaminant concentration in surface water (mg/l)	0.005
CR = ingestion rate (L/person-hour)	28
ET = exposure time (hours/event)	45
EF = exposure frequency (events/yr)	35
ED = exposure duration (yr)	70
BW = body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	35
DY = days per year (days)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RfD = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

NEW RIVER

COMP	Concentration Carcinogen (mg/l)	Contact Rate (hour)	Exposure Time (hrs/event)	Exposure Frequency (events/yr)	Exposure Duration (years)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg-day)	Cancer Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Averaging time Noncarc. (days)	Noncarc. Dose (mg/kg-day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Nutrient	7.7E+00	0.005	28	45	30	70	25500	1.0E-05	0.0E+00	0%	1.0E-05	1.0E-05	1.0E-05	1.0E-05	4%	
Arsenic	4.3E-03	0.005	28	45	30	70	25500	4.2E-08	1.0E+00	7.4E-08	100%	1.0E-05	3.0E-04	3.0E-04	35%	
Cadmium (water)	4.2E-03	0.005	28	45	30	70	25500	4.1E-08	0.0E+00	0.0E+00	0%	1.0E-05	9.0E-08	5.0E-04	23%	
Lead	1.8E-02	0.005	28	45	30	70	25500	1.8E-07	0.0E+00	0.0E+00	0%	1.0E-05	4.1E-07	3.7E-02	1.1E-05	
Manganese (water)	8.0E-02	0.005	28	45	30	70	25500	4.8E-07	0.0E+00	0.0E+00	0%	1.0E-05	1.1E-08	6.0E-03	2.0E-04	
Vanadium	8.1E-03	0.005	28	45	30	70	25500	8.0E-08	0.0E+00	0.0E+00	0%	1.0E-05	1.4E-07	7.0E-03	2.0E-05	
Zinc	3.0E-01	0.005	28	45	30	70	25500	3.0E-08	0.0E+00	0.0E+00	0%	1.0E-05	8.0E-08	3.0E-01	3%	
AT-DOE	4.0E-05	0.005	28	45	30	70	25500	3.0E-10	3.4E-01	1.3E-10	0%	1.0E-05	9.0E-10	0.0E+00	0.0E+00	
TOTAL	8.0E-05	0.005	28	45	30	70	25500	4.9E-10	2.4E-01	1.3E-10	0%	1.0E-05	0.0E+00	0.0E+00	0%	

COOGELS CREEK

COMP	Concentration Carcinogen (mg/l)	Contact Rate (hour)	Exposure Time (hrs/event)	Exposure Frequency (events/yr)	Exposure Duration (years)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg-day)	Cancer Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Averaging time Noncarc. (days)	Noncarc. Dose (mg/kg-day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Nutrient	9.4E-01	0.005	28	45	30	70	25500	9.4E-08	0.0E+00	0.0E+00	0%	1.0E-05	2.1E-05	1.0E+00	2.1E-05	5%
Arsenic	2.6E-03	0.005	28	45	30	70	25500	2.6E-08	1.0E+00	4.4E-08	100%	1.0E-05	3.0E-04	2.0E-04	44%	
Lead	3.7E-03	0.005	28	45	30	70	25500	3.7E-08	0.0E+00	0.0E+00	0%	1.0E-05	8.0E-08	0.0E+00	0%	
Manganese (water)	4.9E-02	0.005	28	45	30	70	25500	4.9E-07	0.0E+00	0.0E+00	0%	1.0E-05	1.1E-08	6.0E-03	2.0E-04	
Vanadium	2.4E-03	0.005	28	45	30	70	25500	2.4E-08	0.0E+00	0.0E+00	0%	1.0E-05	5.0E-08	7.0E-03	8.0E-03	
Zinc	1.2E-02	0.005	28	45	30	70	25500	1.2E-07	0.0E+00	0.0E+00	0%	1.0E-05	2.7E-07	3.0E-01	9.1E-07	
TOTAL									4.9E-08						4.3E-04	

SURFACE WATER DERMAL DOSE ASSESSMENT  
 OPERABLE UNIT NO. 1 (SITE 2B - NEW RIVER AND COOKS CREEK)  
 REMEDIAL INVESTIGATION - CTO-0201  
 MC2 CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT AND FUTURE ADULT RESIDENT

The intake from dermal contact with surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = Cw \times SA \times PC \times ET \times EF \times ED \times CF/BW \times ATc \text{ or } ATn \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RID$$

CHANGE!

INPUTS	
CW = contaminant concentration in water (mg/l)	
SA = skin surface available for contact (cm <sup>2</sup> )	5000
PC = contaminant specific dermal permeability (cm/h)	Specific
ET = exposure time (hours/day)	2.6
EF = exposure frequency (days/yr)	45
ED = exposure duration (years)	30
CF = volumetric conversion factor for water (liter/1000 cm <sup>3</sup> )	0.001
BW = body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATn = averaging time for noncarcinogen (yr)	30
DY = days per year (days)	365
CSF = cancer slope factor (mg/kg/day)-1	Specific
RID = reference dose (mg/kg-day)	Specific

Note: Inputs are site and scenario specific

NEW RIVER

OPC	Concentration (mg/l)	Surface Area (cm <sup>2</sup> )	Dermal Permeability (cm/h)	Exposure Time (hours/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	Volumetric Conversion (l/m <sup>3</sup> )	Body Weight (kg)	Averaging Cmt Time (days)	Carc. Dose (mg/kg-day)	Derm. Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Average Noncarc. Time (days)	Noncarc. Dose (mg/kg-day)	Derm. Adjust Reference Dose (mg/kg-day)	Reference Risk	Percent Noncarcinogenic Risk
chromium	1.7E+00	5000	1.0E-03	2.6	45	30	0.001	70	2500	1.9E-05	0.0E+00	0%	1.0E-05	4.4E-05	2.0E-07	2.2E-04	0%	
nickel	4.0E-03	5000	1.0E-03	2.6	45	30	0.001	70	2500	4.9E-08	0.0E+00	4.3E-07	22%	1.0E-05	1.1E-07	6.0E-05	1.0E-03	30%
cadmium (water)	4.2E-03	5000	1.0E-03	2.6	45	30	0.001	70	2500	4.8E-08	0.0E+00	0%	1.0E-05	1.1E-07	1.0E-04	1.1E-03	23%	
copper	1.0E-02	5000	1.0E-03	2.6	45	30	0.001	70	2500	2.1E-07	0.0E+00	0%	1.0E-05	4.6E-07	7.4E-03	8.8E-06	1%	
lead	2.3E-02	5000	4.0E-08	2.6	45	30	0.001	70	2500	1.1E-09	0.0E+00	0%	1.0E-05	2.6E-09	0.0E+00	0.0E+00	0%	
manganese (water)	8.0E-02	5000	1.0E-03	2.6	45	30	0.001	70	2500	8.7E-07	0.0E+00	0%	1.0E-05	1.3E-06	1.0E-03	1.3E-03	27%	
vanadium	8.1E-03	5000	1.0E-03	2.6	45	30	0.001	70	2500	6.9E-08	0.0E+00	0%	1.0E-05	1.6E-07	1.4E-03	1.2E-04	2%	
Zinc	3.0E-01	5000	6.0E-04	2.6	45	30	0.001	70	2500	2.9E-08	0.0E+00	0%	1.0E-05	5.8E-08	6.0E-02	8.8E-05	2%	
M-00E	4.0E-05	5000	2.0E-01	2.6	45	30	0.01	70	2500	1.1E-08	0.0E+00	7.4E-07	38%	1.0E-05	2.6E-08	0.0E+00	0.0E+00	0%
U-00D	5.0E-05	5000	2.0E-01	2.6	45	30	0.01	70	2500	1.6E-05	4.0E-01	7.0E-07	40%	1.0E-05	3.7E-08	0.0E+00	0.0E+00	0%
<b>TOTAL</b>																		

SURFACE WATER DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO.7 (BTR 28 - Ne... - VEH AND COOGLES CREEK)  
 REMEDIAL INVESTIGATION - CTD-0201  
 MOB CAMP LEELINE, NORTH CAROLINA  
 CURRENT AND FUTURE ADULT RESIDENT

The intake from dermal contact with surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = \text{Cw} * \text{SA} * \text{PC} * \text{ET} * \text{EF} * \text{ED} = \text{CF/BW} * \text{ATc or ATnc} * \text{DY}$$

Risk = Intake + CSF or RD

CHANGE!

INPUTS

CW = contaminant concentration in water (mg/l)	
SA = skin surface available for contact (cm <sup>2</sup> )	5800
PC = contaminant specific dermal permeability (cm/hr)	Specific
ET = exposure time (hours/day)	2.6
EF = exposure frequency (days/yr)	45
ED = exposure duration (years)	30
CF = volumetric conversion factor for water (liter/1000 cm <sup>3</sup> )	0.001
BW = body weight (kg)	70
ATc = averaging time for carcinogen (y)	70
ATnc = averaging time for noncarcinogen (y)	30
DY = days per year (days)	365
CSF = cancer slope factor (mg/kg-day) <sup>-1</sup>	Specific
RD = reference dose (mg/kg-day)	Specific

Note: Inputs are site and scenario specific

COOGLES CREEK

CP/C	Concentration Carcinogen (mg/l)	Source Area (m <sup>2</sup> )	Dermal Permeability (cm/hr)	Exposure Time (hours/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	Volumetric Conversion (L/m <sup>3</sup> )	Body Weight (kg)	Averaging Carc Time (days)	Carc Dose (mg/kg-day)	Slope Factor (mg/kg-day) <sup>-1</sup>	Carcinogenic Risk	Percent Carcinogenic Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg-day)	Reference Dose (mg/kg-day)	NONC/R Risk	Percent Noncarcinogenic Risk
Ruthenium	3.4E-01	5800	1.0E-03	2.6	45	30	0.001	70	22600	1.1E-06	0.0E+00	0%	10650	2.3E-03	2.0E-07	1.3E-01	0%	
Intrinsic	2.6E-03	5800	1.0E-03	2.6	45	30	0.001	70	22600	2.6E-08	8.6E+00	2.6E-07	100%	10650	6.0E-08	6.0E-05	1.1E-03	44%
Lead	3.7E-03	5800	4.0E-08	2.6	45	30	0.001	70	22600	1.7E-10	0.0E+00	0.0E+00	0%	10650	3.0E-10	0.0E+00	0.0E+00	0%
Vanadocene (water)	4.9E-02	5800	1.0E-03	2.6	45	30	0.001	70	22600	6.6E-07	0.0E+00	0.0E+00	0%	10650	1.3E-08	1.3E-03	1.3E-03	50%
Vanadium	2.4E-03	5800	1.0E-03	2.6	45	30	0.001	70	22600	2.8E-08	0.0E+00	0.0E+00	0%	10650	6.0E-08	1.4E-03	4.6E-05	2%
Iron	1.2E-02	5800	6.0E-04	2.6	45	30	0.001	70	22600	8.1E-08	0.0E+00	0.0E+00	0%	10650	1.0E-07	6.0E-02	3.0E-08	0%
TOTAL																	2.0E-03	

SEDIMENT INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B - ODE POND AND NEW RIVER)  
 EMERGENCY INVESTIGATION CTO-0221  
 NC3 CAMP LEJEUNE, NORTH CAROLINA  
 FISHERMAN

Intake from ingestion of sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times IR \times CF \times EF \times ED / BW \times ATC \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RD$$

Where:

	INPUTS
C = contaminant concentration in sediment (mg/kg)	1E-05
CF = conversion for kg to mg	48
EF = exposure frequency (days/yr)	30
ED = exposure duration (yr)	100
IR = soil ingestion rate (mg/day)	70
BW = body weight (kg)	70
ATC = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	30
DY = days per year (days/yr)	365
CSF = cancer slope factor (mg/kg-day)-1	Specific
RD = reference dose (mg/kg-day)	Specific

Note: Inputs are scenario and site specific

ODE POND

COPC	Concentration (mg/kg)	Exposure Frequency (days/yr)	Exposure Duration (yr)	Ingestion Rate (mg/day)	Conversion Factor (kg/mg)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Antimony	4.9E+03	48	30	100	1E-05	70	25500	3.9E-04	1.00E+00	0%	1.0E-04	1.0E-04	1.00E+00	9.2E-04	17%	
Vanadium	6.6E+00	48	30	100	1E-05	70	25500	8.2E-07	1.79E+00	8%	1.0E-03	3.00E-04	4.0E-03	70%	0%	
Chromium	3.2E-01	48	30	100	1E-05	70	25500	2.6E-08	4.30E+00	1.1E-07	11%	1.0E-03	6.00E-03	1.2E-05	0%	
Cobalt	1.2E+01	48	30	100	1E-05	70	25500	9.8E-07	0.00E+00	0%	1.0E-03	2.2E-05	1.00E+00	2.2E-05	0%	
Copper	1.7E+00	48	30	100	1E-05	70	25500	1.4E-07	0.00E+00	0%	1.0E-03	3.2E-07	6.00E-02	6.3E-08	0%	
Lead	8.3E+00	48	30	100	1E-05	70	25500	8.7E-07	0.00E+00	0%	1.0E-03	3.7E-02	8.6E-08	0%	0%	
Manganese (soil)	9.8E+00	48	30	100	1E-05	70	25500	7.9E-07	0.00E+00	0%	1.0E-03	1.8E-06	1.40E-01	1.3E-05	0%	
nickel	2.2E+00	48	30	100	1E-05	70	25500	1.8E-07	0.00E+00	0%	1.0E-03	4.1E-07	2.00E-02	2.1E-05	0%	
Radium	1.2E+01	48	30	100	1E-05	70	25500	9.3E-07	0.00E+00	0%	1.0E-03	2.2E-06	7.00E-03	3.1E-04	0%	
U-238	8.3E-03	48	30	100	1E-05	70	25500	8.7E-10	2.40E-01	1.0E-10	0%	1.0E-03	0.00E+00	0.00E+00	0%	0%
UTAC								1.0E-08						5.3E-03		

SEDIMENT DERMAL CONTACT ASSESSMENT  
OPERABLE UNIT NO. 7 (SITE 2B - CHUC POND AND NEW RIVER)  
REMEDIATION INVESTIGATION CTD-0201  
MCB CAMP LEJEUNE, NORTH CAROLINA  
FISHERMAN

The intake from dermal contact to sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \times ATnc \times DY$$

Risk = Intake  $\times$  CSF or RfD

Where:

C = contaminant concentration in soil (mg/kg)

INPUTS

CF = conversion factor (kg/mg) 1.0E-06

6800

SA = exposed skin surface area (cm<sup>2</sup>) 1

Specific

AF = sediment to skin adherence factor (mg/cm<sup>2</sup>) 48

Specific

Abs = fraction absorbed (unless) (contaminant specific) 33

Specific

EF = exposure frequency (events/day) 33

Specific

ED = exposure duration (year) 70

Specific

BW = body weight (kg) 70

Specific

ATc = averaging time for carcinogen (yr) 33

Specific

ATnc = averaging time for noncarcinogen (yr) 33

Specific

DY = day per year (day/yr) 365

Specific

CSF = cancer slope factor (mg/kg-day)-1 1

Specific

RfD = reference dose (mg/kg-day)

Note: Inputs are scenario and site specific

CHOC POND

Contaminant	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> )	Adherence Factor (mg/cm <sup>2</sup> )	ATcS Factor (%)	Exposure Frequency (events/day)	Exposure Duration (yr)	Body Weight (kg)	Average Carc Time (days)	Carc. Dose (mg/kg/day)	Dermal Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Average Noncarc Time (days)	Noncarc. Dose (mg/kg/day)	Dermal Adjust Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Antimony	8.0E-03	1E-06	5800	1	0.001	48	30	70	22500	2.2E-03	0.00E+00	0%	10850	3.0E-03	2.00E-01	2.7E-04	17%	
Arsenic	6.4E+00	1E-06	5800	1	0.001	48	30	70	22500	3.0E-08	8.75E+00	85%	10850	7.0E-08	6.00E-05	1.2E-03	70%	
Beryllium	3.2E-01	1E-06	5800	1	0.001	48	30	70	22500	1.6E-09	2.1E+01	32%	10850	3.8E-09	1.00E-03	3.8E-06	0%	
Chromium	1.2E+01	1E-06	5800	1	0.001	48	30	70	22500	5.8E-08	0.00E+00	0%	10850	1.3E-07	2.00E-01	6.4E-07	0%	
Cobalt	1.7E+00	1E-06	5800	1	0.001	48	30	70	22500	7.0E-09	0.00E+00	0%	10850	1.8E-08	1.20E-02	1.8E-06	0%	
Copper	1.7E+00	1E-06	5800	1	0.001	48	30	70	22500	7.0E-09	0.00E+00	0%	10850	1.8E-08	7.42E-03	2.8E-06	0%	
Lead	8.3E+00	1E-06	5800	1	0.001	48	30	70	22500	3.0E-08	0.00E+00	0%	10850	9.0E-08	0.00E+00	0.0E+00	0%	
Manganese (soil)	9.8E+00	1E-06	5800	1	0.001	48	30	70	22500	4.0E-08	0.00E+00	0%	10850	1.1E-07	2.80E-02	3.8E-06	0%	
Nickel	2.2E+00	1E-06	5800	1	0.001	48	30	70	22500	1.0E-08	0.00E+00	0%	10850	2.4E-08	4.00E-03	6.0E-06	0%	
Platinum	1.2E+01	1E-06	5800	1	0.001	48	30	70	22500	6.4E-08	0.00E+00	0%	10850	1.3E-07	1.40E-03	9.0E-05	5%	
Uranium	8.2E-03	1E-06	5800	1	0.01	48	30	70	22500	3.8E-10	4.80E-01	1.8E-10	0%	10850	9.0E-10	0.00E+00	0.0E+00	0%
TOTAL											2.0E-07					1.5E-11		

SEDIMENT INGESTION EXPOSURE - ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B - ORDE POND AND NEW RIVER)  
 REMEDIAL INVESTIGATION CTC-0231  
 MC8 CAMP LEJEUNE, NORTH CAROLINA  
 FISHERMAN

Intake from ingestion of sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times R \times CF \times EF \times ED / BW \times ATC \text{ or } ATNC \times DY$$

Risk = Intake \* CSF or RD

Where:

C = contaminant concentration in sediment (mg/kg)

CF = conversion factor kg to mg

EF = exposure frequency (days/yr)

ED = exposure duration (yr)

R = ACR ingestion rate (mg/day)

BW = body weight (kg)

ATC = averaging time for carcinogen (yr)

ATNC = averaging time for noncarcinogen (yr)

DY = days per year (days/year)

CSF = cancer slope factor (mg/kg-day)-1

RD = reference dose (mg/kg-day)

INPUTS

1E-08

48

30

100

1E-08

70

25560

2.1E-05

1.7E-07

1.3E-08

72%

10860

4.0E-05

3.0E-04

6.6E-03

4.2E-05

7.0E-02

0%

10860

2.6E-04

3.71E-02

6.6E-03

5%

10860

0.00E+00

0.0E+00

0%

10860

1.7E-05

3.0E-01

5.6E-05

0%

10860

2.6E-03

0.0E+00

0%

10860

1.6E-03

0.0E+00

0%

10860

5.0E-04

1.1E-04

0%

10860

6.2E-07

3.0E-01

1.7E-07

0%

10860

2.2E-07

0.00E+00

0%

10860

1.1E-07

0.0E+00

0%

10860

6.0E-04

0.0E+00

0%

10860

2.4E-07

4.0E-02

8.6E-05

0%

10860

5.0E-08

0.0E+00

0%

10860

1.4E-07

3.0E-02

4.6E-08

0%

10860

2.0E-07

3.0E-02

8.6E-08

0%

10860

1.2E-08

2.1E-05

2.1E-05

1.4E-05

0%

TOTAL

1.8E-08

1.4E-01

Note: Inputs are scenario and site specific

NEW RIVER

POPC	Concentration (mg/kg)	Exposure Frequency (days/yr)	Exposure Duration (yr)	Ingestion Rate (mg/day)	Conversion Factor (kg/mg)	Body Weight (kg)	Average Carb Time (days)	CSC Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Average Noncancer Time (days)	Noncancer Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Molybdenum	2.0E+02	48	30	100	1E-08	70	25560	2.1E-05	0.00E+00	0%	10860	4.0E-05	4.0E-04	1.2E-01	91%	
Arsenic	9.0E+00	48	30	100	1E-08	70	25560	7.2E-07	1.7E-08	72%	10860	3.0E-04	6.6E-03	4%		
Boron	1.0E+01	48	30	100	1E-08	70	25560	1.2E-06	0.0E+00	0%	10860	2.6E-05	7.0E-02	4.2E-05	0%	
Copper	1.3E+03	48	30	100	1E-08	70	25560	1.1E-04	0.00E+00	0%	10860	2.6E-04	3.71E-02	6.6E-03	5%	
Lead	3.0E+04	48	30	100	1E-08	70	25560	3.1E-03	0.00E+00	0%	10860	0.00E+00	0.0E+00	0%		
Silver	2.1E+00	48	30	100	1E-08	70	25560	1.7E-07	0.00E+00	0%	10860	4.0E-07	6.0E-03	8.0E-05	0%	
Zinc	9.0E+01	48	30	100	1E-08	70	25560	7.2E-08	0.00E+00	0%	10860	1.7E-05	3.0E-01	5.6E-05	0%	
1,4-DOD	1.0E-02	48	30	100	1E-08	70	25560	1.2E-09	2.4E-01	2.9E-10	10860	3.0E-09	0.0E+00	0.0E+00	0%	
1,4-CDE	8.0E-03	48	30	100	1E-08	70	25560	8.0E-10	3.4E-01	2.3E-10	10860	1.6E-09	0.00E+00	0.0E+00	0%	
1,4-COT	3.0E-01	48	30	100	1E-08	70	25560	2.4E-08	3.4E-01	8.2E-09	10860	5.0E-08	5.0E-04	1.1E-04	0%	
Anthracene	2.0E-07	48	30	100	1E-08	70	25560	2.2E-08	0.00E+00	0%	10860	6.2E-07	3.0E-01	1.7E-07	0%	
Benzo(a)anthracene	1.2E+00	48	30	100	1E-08	70	25560	9.0E-08	7.3E-01	6.6E-08	10860	2.2E-07	0.00E+00	0.0E+00	0%	
Benzo(a)pyrene	8.0E-01	48	30	100	1E-08	70	25560	4.0E-08	7.0E-01	3.5E-07	10860	1.1E-07	0.00E+00	0.0E+00	0%	
Benzo(b)fluoranthene	1.1E+00	48	30	100	1E-08	70	25560	8.0E-08	7.3E-01	6.6E-08	10860	2.1E-07	0.00E+00	0.0E+00	0%	
Benzo(g,h,perylene	9.0E-01	48	30	100	1E-08	70	25560	2.4E-08	0.00E+00	0%	10860	6.0E-08	3.0E-02	1.9E-08	0%	
Benzo(k)fluoranthene	4.0E-01	48	30	100	1E-08	70	25560	3.0E-08	7.3E-02	2.9E-08	10860	9.1E-08	0.00E+00	0.0E+00	0%	
Benzo(2-methoxy)phthalate	1.3E+00	48	30	100	1E-08	70	25560	1.0E-07	1.4E-02	1.4E-08	10860	2.4E-07	2.0E-02	1.2E-05	0%	
Carbazole	1.0E-01	48	30	100	1E-08	70	25560	1.3E-08	2.0E-02	2.0E-10	10860	3.0E-09	0.00E+00	0.0E+00	0%	
Chrysene	1.0E+00	48	30	100	1E-08	70	25560	1.0E-07	7.3E-03	9.1E-10	10860	2.0E-07	0.00E+00	0.0E+00	0%	
Dibenzofuran	6.0E-02	48	30	100	1E-08	70	25560	4.0E-09	0.00E+00	0%	10860	1.1E-08	0.00E+00	0.0E+00	0%	
Fluorene	1.0E-01	48	30	100	1E-08	70	25560	1.0E-07	0.00E+00	0%	10860	2.4E-07	4.0E-02	8.6E-08	0%	
Indeno(1,2,3-cd)pyrene	3.0E-01	48	30	100	1E-08	70	25560	2.4E-08	7.3E-01	1.1E-08	10860	5.0E-08	0.00E+00	0.0E+00	0%	
Phenanthrene	7.7E-01	48	30	100	1E-08	70	25560	6.0E-08	0.00E+00	0%	10860	1.4E-07	3.0E-02	4.6E-08	0%	
Tyrene	1.1E+00	48	30	100	1E-08	70	25560	8.0E-08	0.00E+00	0%	10860	2.0E-07	3.0E-02	8.6E-08	0%	
alpha-Chlordane	6.0E-03	48	30	100	1E-08	70	25560	5.0E-10	1.3E-02	6.6E-10	10860	1.2E-09	0.00E-05	2.1E-05	0%	
gamma-Chlordane	4.0E-03	48	30	100	1E-08	70	25560	3.7E-10	4.8E-10	0%	10860	8.0E-10	0.00E-05	1.4E-05	0%	

SEDIMENT DERIVATIVE  
OPERABLE UNIT NO.7 (SITE 2a)  
POND AND NEW RIVER  
PRELIMINARY INVESTIGATION CTD-0201  
MCB CAMP LEJEUNE, NORTH CAROLINA  
FISHERMAN

The intake from dermal contact to sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RD$$

Where:

C = contaminant concentration in soil (mg/kg)

CF = conversion factor (kg/mg)

SA = exposed skin surface area (cm<sup>2</sup>)

INPUTS

AF = sediment to skin availability factor (mg/cm<sup>2</sup>)

Abs = fraction absorbed (unless) (contaminant specific)

EF = exposure frequency (events/yr)

ED = exposure duration (years)

BW = body weight (kg)

ATc = averaging time for carcinogen (yr)

ATnc = averaging time for noncarcinogen (yr)

DY = day per year (day/yr)

CSF = cancer slope factor (mg/kg-day)-1

RD = reference dose (mg/kg-day)

Specific

Specific

Specific

Specific

Specific

Note: Inputs are scenario and site specific.

NEW RIVER

CONTAMINANT	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> )	Availability Factor (%)	Abs Factor (%)	Exposure Frequency (events/yr)	Exposure Duration (yr)	Body Weight (kg)	Average Carc Time (day)	Carc Dose (mg/kg/day)	Dermal Adjust. Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Dermal Adjust. Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Amidopyrine	2.0E+02	1E-05	8800	1	0.001	48	30	70	28500	1.2E+00	0.00E+00	0%	10800	2.8E-05	0.00E+00	3.0E-02	9.0E-05	90%
Anisole	8.0E+00	1E-05	8800	1	0.001	48	30	70	28500	4.2E-08	8.7E+00	3.7E-07	30%	10800	9.8E-08	8.00E-05	1.8E-03	4%
Benzene	1.0E+01	1E-05	8800	1	0.001	48	30	70	28500	7.2E-08	0.00E+00	0.0E+00	0%	10800	1.7E-07	1.0E-02	1.8E-03	0%
Copper	1.3E+03	1E-05	8800	1	0.001	48	30	70	28500	6.3E-05	0.00E+00	0.0E+00	0%	10800	1.8E-05	7.4E-03	2.0E-03	5%
Lead	3.0E+04	1E-05	8800	1	0.001	48	30	70	28500	1.8E-04	0.00E+00	0.0E+00	0%	10800	4.2E-04	0.0E+00	0.0E+00	0%
River	2.1E+00	1E-05	8800	1	0.001	48	30	70	28500	1.0E-08	0.00E+00	0.0E+00	0%	10800	2.3E-08	1.0E-03	2.3E-05	0%
Phenol	9.0E+01	1E-05	8800	1	0.001	48	30	70	28500	4.2E-07	0.00E+00	0.0E+00	0%	10800	9.8E-07	8.00E-02	1.8E-05	0%
1,4-EDDO	1.0E+02	1E-05	8800	1	0.001	48	30	70	28500	7.0E-10	4.0E-01	3.4E-10	0%	10800	1.8E-09	0.00E+00	0.0E+00	0%
1,4-EDDE	8.0E+03	1E-05	8800	1	0.001	48	30	70	28500	4.0E-10	8.0E-01	2.7E-10	0%	10800	9.3E-10	0.00E+00	0.0E+00	0%
1,4-DDT	3.0E+01	1E-05	8800	1	0.001	48	30	70	28500	1.4E-08	8.0E-01	9.8E-09	1%	10800	3.8E-08	3.8E-04	1.3E-04	0%
Anthracene	2.0E+01	1E-05	8800	1	0.001	48	30	70	28500	1.3E-08	0.00E+00	0.0E+00	0%	10800	3.0E-08	1.0E-01	2.0E-07	0%
Benz(a)anthracene	1.0E+00	1E-05	8800	1	0.001	48	30	70	28500	8.0E-09	1.4E+00	8.0E-08	0%	10800	1.8E-07	0.00E+00	0.0E+00	0%
Benz(a)pyrene	6.0E-01	1E-05	8800	1	0.001	48	30	70	28500	2.0E-08	1.4E+01	3.8E-07	41%	10800	8.1E-08	0.00E+00	0.0E+00	0%
Benz(b)fluoranthene	1.1E+00	1E-05	8800	1	0.001	48	30	70	28500	5.1E-08	1.4E+00	7.5E-08	0%	10800	1.2E-07	0.00E+00	0.0E+00	0%
Benz(b)phenanthrene	3.0E+01	1E-05	8800	1	0.001	48	30	70	28500	1.4E-08	0.00E+00	0.0E+00	0%	10800	3.8E-08	1.8E-02	2.3E-05	0%
Benz(k)fluoranthene	4.0E-01	1E-05	8800	1	0.001	48	30	70	28500	2.3E-08	1.4E+01	3.3E-08	0%	10800	5.5E-08	0.00E+00	0.0E+00	0%
Bi(2-naphthyl)phthalate	1.0E+00	1E-05	8800	1	0.001	48	30	70	28500	6.0E-08	2.0E+02	1.7E-08	0%	10800	1.4E-07	1.0E-02	1.4E-05	0%
Carbazole	1.0E+01	1E-05	8800	1	0.001	48	30	70	28500	7.0E-09	4.0E+02	3.0E-10	0%	10800	1.7E-08	0.00E+00	0.0E+00	0%
Chrysene	1.0E+00	1E-05	8800	1	0.001	48	30	70	28500	7.3E-08	1.4E+02	1.1E-08	0%	10800	1.8E-07	0.00E+00	0.0E+00	0%
Dibenzofuran	8.0E-01	1E-05	8800	1	0.001	48	30	70	28500	2.8E-08	0.00E+00	0.0E+00	0%	10800	6.8E-09	0.00E+00	0.0E+00	0%
Tetrahydro-	1.0E+00	1E-05	8800	1	0.001	48	30	70	28500	5.8E-08	0.00E+00	0.0E+00	0%	10800	1.4E-07	2.0E-02	8.8E-08	0%
Tetrahydro-	3.0E+01	1E-05	8800	1	0.001	48	30	70	28500	1.4E-08	1.4E+00	2.0E-08	0%	10800	3.8E-08	0.00E+00	0.0E+00	0%
Therphene	7.7E+01	1E-05	8800	1	0.001	48	30	70	28500	3.6E-08	0.00E+00	0.0E+00	0%	10800	8.4E-08	1.8E-02	6.8E-06	0%
Pyrene	1.1E+00	1E-05	8800	1	0.001	48	30	70	28500	6.0E-08	0.00E+00	0.0E+00	0%	10800	1.3E-07	1.8E-02	7.8E-08	0%
alpha-Chordane	8.0E-03	1E-05	8800	1	0.001	48	30	70	28500	3.1E-10	2.8E+00	8.0E-10	0%	10800	7.2E-10	3.0E-03	2.4E-05	0%
gamma-Chordane	4.0E-03	1E-05	8800	1	0.001	48	30	70	28500	2.1E-10	2.8E+00	8.0E-10	0%	10800	8.0E-10	3.0E-03	1.7E-05	0%
TOTAL																		

SURFACE WATER INGESTION DOSE/EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO.7 (SITE 2B - ODE POND AND NEW RIVER)  
 REMEDIAL INVESTIGATION CTD-021  
 MCN CAMP LEJEUNE, NORTH CAROLINA  
 FISHERMAN

The intake from the ingestion of surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = C_w \times CR \times ET \times EF \times ED/SW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RD$$

CHANGE!!!!

Where:

	INPUT
Cw = contaminant concentration in surface water (mg/l)	0.005
CR = contact rate (hour/week)	2.6
ET = exposure time (hours/week)	48
EF = exposure frequency (events/yr)	50
ED = exposure duration (yr)	70
BW = body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	50
DY = days per year (days)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RD = reference dose (mg/kg-day)	specific

ODE POND

COPC	Concentration (mg/l)	Contact Rate (hour)	Exposure Time (hours/week)	Exposure Frequency (events/yr)	Exposure Duration (years)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg-day)	Cancer Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Averaging Time Noncarc. (years)	Noncarc. Dose (mg/kg-day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
chromium	1.2E-03	0.005	2.6	48	30	70	25500	1.7E-05	0.00E+00	0%	1.0E-02	2.0E-05	1.0E-02	2.0E-05	0%	
Nickel	0.3E-03	0.005	2.6	48	30	70	25500	9.7E-06	0.00E+00	0%	1.0E-02	2.3E-07	2.0E-02	1.1E-05	1%	
Thallium	6.0E-03	0.005	2.6	48	30	70	25500	6.9E-06	0.00E+00	0%	1.0E-02	1.6E-07	8.0E-05	2.0E-03	0%	
<b>TOTAL</b>									0.0E+00						2.0E-03	

NEW RIVER

COPC	Concentration Carcinogen (mg/l)	Contact Rate (hour)	Exposure Time (hours/week)	Exposure Frequency (events/yr)	Exposure Duration (years)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg-day)	Cancer Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Averaging Time Noncarc. (years)	Noncarc. Dose (mg/kg-day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Chromium	1.7E+00	0.005	2.6	48	30	70	25500	1.7E-05	0.00E+00	0%	1.0E-02	4.1E-05	1.0E+00	4.1E-05	4%	
Arsenic	4.0E-03	0.005	2.6	48	30	70	25500	4.0E-06	1.7E+00	100%	1.0E-02	1.1E-07	3.0E-04	3.0E-04	39%	
Chromium (water)	4.2E-03	0.005	2.6	48	30	70	25500	4.4E-06	0.00E+00	0%	1.0E-02	1.0E-07	6.0E-04	2.1E-04	23%	
Copper	1.0E-02	0.005	2.6	48	30	70	25500	1.0E-07	0.00E+00	0%	1.0E-02	4.4E-07	3.7E-02	1.2E-05	1%	
Lead	2.0E-02	0.005	2.6	48	30	70	25500	2.0E-07	0.00E+00	0%	1.0E-02	8.7E-07	0.00E+00	0.0E+00	0%	
Manganese (water)	5.0E-02	0.005	2.6	48	30	70	25500	5.0E-07	0.00E+00	0%	1.0E-02	1.2E-06	5.0E-03	2.4E-04	27%	
Vanadium	8.1E-03	0.005	2.6	48	30	70	25500	8.1E-08	0.00E+00	0%	1.0E-02	1.6E-07	7.0E-03	2.1E-05	2%	
Zinc	3.0E-01	0.005	2.6	48	30	70	25500	3.0E-08	0.00E+00	0%	1.0E-02	8.9E-09	3.0E-01	3.0E-05	3%	
4,4'-DDT	4.0E-05	0.005	2.6	48	30	70	25500	4.2E-10	3.4E-01	1.4E-10	0%	1.0E-02	0.00E+00	0.0E+00	0%	
4,4'-DDD	6.0E-05	0.005	2.6	48	30	70	25500	6.2E-10	2.4E-01	1.3E-10	0%	1.0E-02	0.00E+00	0.0E+00	0%	
<b>TOTAL</b>									7.0E-08						9.0E-04	

SURFACE WATER DERMAL DOSE EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO.7 (SITE 2B - CNOE POND AND NEW RIVER)  
 REMEDIAL INVESTIGATION - CTD-0221  
 MCIS CAMP LEJEUNE, NORTH CAROLINA  
 FISHERMAN

The intake from dermal contact with surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = \text{CV} * \text{SA} * \text{PC} * \text{ET} * \text{EF} * \text{ED} * \text{CF}_{SW} * \text{ATc or ATnc} * \text{CY}$$

$\text{Px} = \text{Intake} * \text{CSF or RID}$

CHANGE III

Where:	INPUTS
CV = contaminant concentration in water (mg/l)	6800
SA = skin surface available for contact (cm <sup>2</sup> )	Specific
PC = contaminant specific dermal permeability (cm/h)	2.6
ET = exposure time (hours/day)	48
EF = exposure frequency (days/yr)	30
ED = exposure duration (years)	0.001
CF = volumetric conversion factor for water (liter/1000 cm <sup>3</sup> )	0.001
BW = body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	30
CY = days per year (days)	365
CSF = cancer slope factor (mg/kg-day)-1	Specific
RID = reference dose (mg/kg-day)	Specific

Note: Inputs are site and scenario specific

CNOE POND

POPs	Concentration Carcinogen (mg/l)	Surface Area (cm <sup>2</sup> )	Dermal Permeability (cm/h)	Exposure Time (hours/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	Volumetric Conversion (l/m <sup>3</sup> )	Body Weight (kg)	Averaging Carr Time (days)	Carr Dose (mg/kg-day)	Dermal Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carr Carcinogenic Risk	Average Noncarr Time (days)	Noncarr Dose (mg/kg-day)	Dermal Adjust Reference Dose (mg/kg-day)	Noncarr Risk	Percent Noncarcinogenic Risk
Naphthalene	1.7E+01	6800	1.0E-03	2.6	48	30	0.001	70	22650	2.1E-06	0.00E+00	0%	10650	4.8E-06	2.0E-01	2.4E-05	0%	
Nickel	1.2E-02	6800	1.0E-04	2.6	48	30	0.001	70	22650	1.6E-06	0.00E+00	0%	10650	3.6E-06	4.0E-03	9.1E-06	0%	
Thallium	4.7E-03	6800	1.0E-03	2.6	48	30	0.001	70	22650	5.7E-06	0.00E+00	0%	10650	1.3E-07	1.0E-05	8.3E-03	100%	
<b>TOTAL:</b>																		

NEW RIVER

POPs	Concentration Carcinogen (mg/l)	Surface Area (cm <sup>2</sup> )	Dermal Permeability (cm/h)	Exposure Time (hours/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	Volumetric Conversion (l/m <sup>3</sup> )	Body Weight (kg)	Averaging Carr Time (days)	Carr Dose (mg/kg-day)	Dermal Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carr Carcinogenic Risk	Average Noncarr Time (days)	Noncarr Dose (mg/kg-day)	Dermal Adjust Reference Dose (mg/kg-day)	Noncarr Risk	Percent Noncarcinogenic Risk
Naphthalene	1.7E+00	6800	1.0E-03	2.6	48	30	0.001	70	22650	2.0E-06	0.00E+00	0%	10650	4.7E-05	2.0E-01	2.4E-05	0%	
Asenic	4.5E-03	6800	1.0E-03	2.6	48	30	0.001	70	22650	6.2E-06	9.7E-06	4.6E-07	74%	10650	1.2E-07	6.0E-06	2.0E-03	
Cadmium (water)	4.2E-03	6800	1.0E-03	2.6	48	30	0.001	70	22650	8.1E-06	0.00E+00	0%	10650	1.3E-07	1.0E-04	1.2E-03	23%	
Copper	1.8E-02	6800	1.0E-03	2.6	48	30	0.001	70	22650	2.2E-07	0.00E+00	0%	10650	5.1E-07	7.4E-03	8.6E-05	1%	
Lead	2.3E-02	6800	4.0E-06	2.6	48	30	0.001	70	22650	1.1E-06	0.00E+00	0%	10650	2.7E-09	0.00E+00	0.0E+00	0%	
Manganese (water)	6.0E-02	6800	1.0E-03	2.6	48	30	0.001	70	22650	6.0E-07	0.00E+00	0%	10650	1.4E-08	1.0E-03	1.4E-03	27%	
Vanadium	6.1E-03	6800	1.0E-03	2.6	48	30	0.001	70	22650	7.4E-06	0.00E+00	0%	10650	1.7E-07	1.4E-03	1.2E-04	2%	
Zinc	3.8E-01	6800	8.0E-04	2.6	48	30	0.001	70	22650	2.6E-06	0.00E+00	0%	10650	6.2E-08	6.0E-02	1.0E-04	2%	
Hg-DOE	4.0E-05	6800	2.4E-01	2.6	48	30	0.001	70	22650	1.2E-07	8.8E-01	7.9E-08	13%	10650	3.7E-07	0.00E+00	0.0E+00	0%
Hg-DOO	6.0E-05	6800	2.0E-01	2.6	48	30	0.001	70	22650	1.7E-07	4.8E-01	8.2E-08	13%	10650	4.0E-07	0.00E+00	0.0E+00	0%
<b>TOTAL:</b>																		

FISH INGESTION EXPOSURE ASSESSMENT  
 SITE 2B - OZONE POND  
 REMEDIAL INVESTIGATION CTD-0231  
 MCAMP LEJEUNE, NORTH CAROLINA  
 FISHERMAN

Intake (mg/kg-day) = CF \* IR \* FI \* EF \* ED/BW \* ATc or ATnc \* DY

Risk = Intake \* CSF or /RfD

Where:

	INPUTS
CF = contaminant concentration in fish (mg/kg)	
IR = adult ingestion rate (kg/meat)	0.284
FI = fraction ingested from contaminated source (unitless)	1.00
EF = adult exposure frequency (meals/yr)	48
ED = adult exposure duration (years)	30
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (years)	70
ATnc = averaging time for noncarcinogen (years)	30
DY = days per year (days/yr)	365

Note: Inputs are scenario and site specific

OZONE POND

Contaminant	Concentration (mg/kg)	Ingestion Rate (kg/meat)	Fraction Ingestion (%)	Exposure Frequency (meals/yr)	Exposure Duration (years)	Average Weight (kg)	Carc Time (days)	Calc Dose (mg/kg-day)	Dose Factor (mg/kg-day)-1	Carcinogenic Risk Adult	Noncarcinogenic Risk Adult	Average Noncarc Time (days)	Noncarc Dose (mg/kg-day)	Harmless Dose (mg/kg-day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Barium	0.0E+01	0.284	1	48	30	70	25560	1.4E-04	0.0E+00	0%	10650	3.2E-01	7.0E-02	4.0E-02	5%	
Manganese (soil)	2.2E+00	0.284	1	48	30	70	25560	6.0E-04	0.0E+00	0%	10650	1.2E-03	1.4E-01	8.4E-03	10%	
Selenium	3.3E-01	0.284	1	48	30	70	25560	7.3E-05	0.0E+00	0%	10650	1.7E-04	5.0E-03	3.4E-02	30%	
Zinc	2.2E+01	0.284	1	48	30	70	25560	8.2E-03	0.0E+00	0%	10650	1.2E-02	3.0E-01	4.1E-02	40%	
<b>TOTAL</b>									0.0E+00	0				8.8E-02		

FISH INGESTION EXPOSURE ASSESSMENT  
SITE: THE NEW RIVER  
REMEDIATION INVESTIGATION CTD-0201  
MCB CAMP LEJEUNE, NORTH CAROLINA  
FISHERMAN

Intake (mg/kg-day) = CF \* IR \* EF \* ED/BW \* ATc or ATnc \* DY

Risk = Intake \* CSF or RfD

Where:

	INPUTS
CF = contaminant concentration in fish (mg/kg)	
IR = adult ingestion rate (kg/meal)	0.284
FI = fraction ingested from contaminated source (unless)	100
EF = adult exposure frequency (meals/yr)	48
ED = adult exposure duration (years)	30
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (years)	70
ATnc = averaging time for noncarcinogen (years)	30
DY = days per year (days/yr)	365

Note: Inputs are scenario and site specific

COPC	Concentration Carcinogen (mg/kg) Adult	Ingestion Rate (kg/meal) Adult	Fraction Ingestion (%)	Exposure Frequency (meals/yr) Adult	Exposure Duration (years) Adult	Body Weight (kg) Adult	Average Carb Time (days)	Carb Dose (mg/kg-day)-1 Adult	Slope Factor (mg/kg-day)-1 Adult	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarb Time (days)	Noncarb Dose (mg/kg-day) Adult	Reference Dose (mg/kg-day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Manganese	2.4E-01	0.284	1	48	30	70	25500	6.6E-05	1.0E+00	0.0E+00	0%	10850	1.3E-04	4.0E-04	3.2E-01	78%
Barium	9.8E-01	0.284	1	48	30	70	25500	2.2E-04	0.0E+00	0.0E+00	0%	10850	8.2E-04	7.0E-02	7.8E-03	2%
Cobalt	2.0E-02	0.284	1	48	30	70	25500	6.9E-05	0.0E+00	0.0E+00	0%	10850	1.6E-05	8.0E-02	2.7E-04	0%
Copper	6.1E-01	0.284	1	48	30	70	25500	1.4E-04	0.0E+00	0.0E+00	0%	10850	3.3E-04	3.7E-02	8.8E-03	2%
Selenium	3.9E-01	0.284	1	48	30	70	25500	8.6E-05	0.0E+00	0.0E+00	0%	10850	2.1E-04	6.0E-03	4.2E-02	10%
U-238	4.2E-02	0.284	1	48	30	70	25500	9.8E-05	3.4E-01	3.3E-05	89%	10850	2.2E-05	0.0E+00	0.0E+00	0%
U-234	2.3E-02	0.284	1	48	30	70	25500	6.3E-08	2.4E-01	1.3E-05	23%	10850	1.2E-05	0.0E+00	0.0E+00	0%
Alpha-Chlordene	3.6E-03	0.284	1	48	30	70	25500	8.2E-07	1.3E+00	1.1E-05	19%	10850	1.8E-05	8.0E-05	3.2E-02	8%
TOTAL																4.1E-01

## SURFACE SOIL INGESTION EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 26)

REMEDIAl INVESTIGATION CTO-0231

MCB CAMP LEJEUNE, NORTH CAROLINA

CURRENT MILITARY PERSONNEL

Intake from ingestion of soil is calculated as follows:

Intake (mg/kg-day) = C \* CF \* EF \* ED \* IR/BW \* ATc or ATnc \* DY

Risk = Intake \* CSF or /RID

Where:

C = contaminant concentration in soil (mg/kg)

CF = conversion for kg to mg

EF = adult exposure frequency (days/yr)

## INPUTS

ED = adult exposure duration (yr)

1E-06

IR = adult soil ingestion rate (mg/day)

250

BW = adult body weight (kg)

4

ATc = averaging time for carcinogen (yr)

100

ATnc = averaging time for noncarcinogen (yr)

70

DY = days per year (days/year)

4

CSF = cancer slope factor (mg/kg-day)-1

365

RID = reference dose (mg/kg-day)

specific

specific

Note: Inputs are scenario and site specific

COPC	Concentration Carcinogen (mg/kg)	Exposure Frequency (days/yr) Adult	Exposure Duration (yr) Adult	Conversion Factor (kg/mg)	Ingestion Rate (mg/day) Adult	Body Weight (kg) Adult	Average Carc Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	3.4E+03	250	4	1E-06	100	70	25550	1.9E-04	0.0E+00	0%	1460	3.3E-03	1.0E+00	3.3E-03	7%	
Antimony	5.7E+00	250	4	1E-06	100	70	25550	3.2E-07	0.0E+00	0%	1460	5.5E-08	4.0E-04	1.4E-02	31%	
Arsenic	1.7E+00	250	4	1E-06	100	70	25550	9.7E-08	1.7E+00	51%	1460	1.7E-08	3.0E-04	5.7E-03	13%	
Barium	3.5E+01	250	4	1E-06	100	70	25550	1.9E-08	0.0E+00	0%	1460	3.4E-05	7.0E-02	4.9E-04	1%	
Cadmium (soil)	1.6E+00	250	4	1E-06	100	70	25550	6.7E-08	0.0E+00	0%	1460	1.8E-08	1.0E-03	1.8E-03	3%	
Chromium	1.0E+01	250	4	1E-06	100	70	25550	5.8E-07	0.0E+00	0%	1460	1.0E-05	1.0E+00	1.0E-05	0%	
Cobalt	9.1E-01	250	4	1E-06	100	70	25550	6.1E-08	0.0E+00	0%	1460	8.9E-07	6.0E-02	1.8E-05	0%	
Copper	1.3E+02	250	4	1E-06	100	70	25550	7.9E-08	0.0E+00	0%	1460	1.3E-04	3.7E-02	3.4E-03	8%	
Lead	1.7E+02	250	4	1E-06	100	70	25550	9.5E-08	0.0E+00	0%	1460	1.7E-04	0.0E+00	0.0E+00	0%	
Manganese (soil)	3.3E+02	250	4	1E-06	100	70	25550	1.8E-05	0.0E+00	0%	1460	3.2E-04	1.4E-01	2.3E-03	5%	
Mercury	2.8E-01	250	4	1E-06	100	70	25550	1.6E-08	0.0E+00	0%	1460	2.8E-07	3.0E-04	9.3E-04	2%	
Nickel	8.6E+00	250	4	1E-06	100	70	25550	3.1E-07	0.0E+00	0%	1460	5.4E-08	2.0E-02	2.7E-04	1%	
Silver	1.1E+00	250	4	1E-06	100	70	25550	6.3E-08	0.0E+00	0%	1460	1.1E-08	5.0E-03	2.2E-04	0%	
Thallium	5.1E-01	250	4	1E-06	100	70	25550	2.9E-08	0.0E+00	0%	1460	5.0E-07	8.0E-06	6.3E-03	14%	
Vanadium	7.9E+00	250	4	1E-06	100	70	25550	4.4E-07	0.0E+00	0%	1460	7.8E-06	7.0E-03	1.1E-03	2%	
Zinc	1.1E+03	250	4	1E-06	100	70	25550	6.0E-06	0.0E+00	0%	1460	1.0E-03	3.0E-01	3.5E-03	8%	
4,4'-DDD	6.4E-02	250	4	1E-06	100	70	25550	3.6E-09	2.4E-01	8.6E-10	0%	1460	6.3E-08	0.0E+00	0.0E+00	0%
4,4'-DDE	3.6E-01	250	4	1E-06	100	70	25550	1.9E-08	3.4E-01	6.6E-09	2%	1460	3.4E-07	0.0E+00	0.0E+00	0%
4,4'-DDT	1.8E-01	250	4	1E-06	100	70	25550	8.3E-09	3.4E-01	2.8E-09	1%	1460	1.5E-07	5.0E-04	2.9E-04	1%
Anthracene	2.1E-01	250	4	1E-06	100	70	25550	1.1E-08	0.0E+00	0.0E+00	0%	1460	2.0E-07	3.0E-01	6.7E-07	0%
Benz(a)anthracene	2.8E-01	250	4	1E-06	100	70	25550	1.6E-08	7.3E-01	1.1E-08	3%	1460	2.7E-07	0.0E+00	0.0E+00	0%
Benz(a)pyrene	2.7E-01	250	4	1E-06	100	70	25550	1.6E-08	7.3E+00	1.1E-07	33%	1460	2.6E-07	0.0E+00	0.0E+00	0%
Benz(b)fluoranthene	2.9E-01	250	4	1E-06	100	70	25550	1.6E-08	7.3E-01	1.2E-08	4%	1460	2.9E-07	0.0E+00	0.0E+00	0%
Benz(o,h,p)perylene	2.5E-01	250	4	1E-06	100	70	25550	1.4E-08	0.0E+00	0.0E+00	0%	1460	2.6E-07	3.0E-02	8.3E-06	0%
Carbazole	1.7E-01	250	4	1E-06	100	70	25550	9.5E-09	2.0E-02	1.9E-10	0%	1460	1.7E-07	0.0E+00	0.0E+00	0%
Chrysene	2.8E-01	250	4	1E-06	100	70	25550	1.6E-08	7.3E-03	1.1E-10	0%	1460	2.7E-07	0.0E+00	0.0E+00	0%
Heptachlor epoxide	6.9E-03	250	4	1E-06	100	70	25550	3.9E-10	9.1E+00	3.5E-09	1%	1460	6.8E-09	1.3E-05	5.2E-04	1%
Indeno(1,2,3-cd)pyrene	2.5E-01	250	4	1E-06	100	70	25550	1.4E-08	7.3E-01	1.0E-08	3%	1460	2.5E-07	0.0E+00	0.0E+00	0%
Phenanthrene	2.8E-01	250	4	1E-06	100	70	25550	1.6E-08	0.0E+00	0.0E+00	0%	1460	2.8E-07	3.0E-02	9.3E-08	0%
Alpha-Chlordane	3.8E-02	250	4	1E-06	100	70	25550	2.1E-09	1.3E+00	2.8E-09	1%	1460	3.7E-08	6.0E-05	6.2E-04	1%
gamma-Chlordane	1.3E-02	250	4	1E-06	100	70	25550	7.2E-10	9.3E+00	9.4E-10	0%	1460	1.3E-08	6.0E-05	2.1E-04	0%
TOTAL								3.3E-07							4.8E-02	

SURFACE SOIL DERMAL CONTAMINATION ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B)  
 REMEDIAL INVESTIGATION CTD-0291  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT MILITARY PERSONNEL

Dermal contact with soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

Risk = Intake  $\times$  CSF or /RD

Where:

C = contaminant concentration in soil (mg/kg)

CF = conversion factor (kg/mg)

SA = adult exposed skin surface area (cm<sup>2</sup>)

AF = soil to skin adherence factor (mg/cm<sup>2</sup>)

Abs = fraction absorbed (unitless)

EF = adult exposure frequency (events/yr)

ED = adult exposure duration (years)

BW = adult body weight (kg)

ATc = averaging time for carcinogen (yr)

ATnc = averaging time for noncarcinogen (yr)

DY = day per year (day/yr)

CSF = cancer slope factor (mg/kg-day)<sup>-1</sup>

RD = reference dose (mg/kg-day)

INPUTS

Specific

1

250

4

70

4

305

specific

specific

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> )	Adherence Factor (mg/cm <sup>2</sup> ) Adult	Fraction Absorbed (%)	Exposure Frequency (events/yr) - Adult	Exposure Duration (yr) - Adult	Body Weight (kg) Adult	Average Cst Time (days)	Carc Dose (mg/kg/day) Adult	Dermal Adult Slope Factor (mg/kg-day)-1 Adult	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Dose (mg/kg/day) Adult	Noncarc Dose (mg/kg/day) Adult	Dermal Adult Reference Dose (mg/kg/day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	3.4E+03	1E-08	4000	1	0.001	250	4	70	25500	8.2E-08	0.00E+00	0.0E+00	0%	1400	1.4E-08	2.0E-01	7.2E-04	7%
Antimony	5.7E+00	1E-08	4000	1	0.001	250	4	70	25500	1.4E-08	0.00E+00	0.0E+00	0%	1400	2.4E-07	8.0E-05	3.0E-03	28%
Arsenic	1.7E+00	1E-08	4000	1	0.001	250	4	70	25500	4.3E-08	0.75E+00	3.7E-08	21%	1400	7.3E-08	8.0E-05	1.2E-03	11%
Barium	3.9E+01	1E-08	4000	1	0.001	250	4	70	25500	8.4E-08	0.00E+00	0.0E+00	0%	1400	1.8E-08	1.4E-02	1.0E-04	1%
Chromium (soil)	1.0E+00	1E-08	4000	1	0.001	250	4	70	25500	3.7E-08	0.00E+00	0.0E+00	0%	1400	6.9E-08	2.0E-04	3.3E-04	0%
Chromium	1.0E+01	1E-08	4000	1	0.001	250	4	70	25500	2.6E-08	0.00E+00	0.0E+00	0%	1400	4.4E-07	2.0E-01	2.2E-03	0%
Cobalt	9.1E-01	1E-08	4000	1	0.001	250	4	70	25500	2.3E-08	0.00E+00	0.0E+00	0%	1400	3.8E-08	1.2E-02	3.2E-03	0%
Copper	1.3E+02	1E-08	4000	1	0.001	250	4	70	25500	3.1E-07	0.00E+00	0.0E+00	0%	1400	6.8E-08	7.4E-03	7.4E-04	7%
Lead	1.7E+02	1E-08	4000	1	0.001	250	4	70	25500	4.1E-07	0.00E+00	0.0E+00	0%	1400	7.1E-08	8.0E+00	0.0E+00	0%
Manganese (soil)	3.3E+02	1E-08	4000	1	0.001	250	4	70	25500	7.0E-07	0.00E+00	0.0E+00	0%	1400	1.4E-05	2.8E-02	4.9E-04	8%
Mercury	2.8E-01	1E-08	4000	1	0.001	250	4	70	25500	6.0E-10	0.00E+00	0.0E+00	0%	1400	1.2E-08	8.0E-05	2.0E-04	2%
Nickel	8.0E+00	1E-08	4000	1	0.001	250	4	70	25500	1.3E-08	0.00E+00	0.0E+00	0%	1400	2.3E-07	4.0E-03	5.9E-05	1%
Silver	1.1E+00	1E-08	4000	1	0.001	250	4	70	25500	2.7E-08	0.00E+00	0.0E+00	0%	1400	4.8E-08	1.0E-03	4.8E-05	0%
Thallium	5.1E-01	1E-08	4000	1	0.001	250	4	70	25500	1.2E-08	0.00E+00	0.0E+00	0%	1400	2.2E-08	1.8E-05	1.4E-03	13%
Vanadium	7.9E+00	1E-08	4000	1	0.001	250	4	70	25500	1.8E-08	0.00E+00	0.0E+00	0%	1400	3.3E-07	1.4E-03	2.4E-04	2%
Zinc	1.1E+03	1E-08	4000	1	0.001	250	4	70	25500	2.0E-08	0.00E+00	0.0E+00	0%	1400	4.6E-08	8.0E-02	7.8E-04	7%
1,4-DOD	6.4E-02	1E-08	4000	1	0.001	250	4	70	25500	1.0E-08	4.80E-01	7.4E-08	0%	1400	2.7E-08	8.0E+00	0.0E+00	0%
1,4-DDE	3.8E-01	1E-08	4000	1	0.001	250	4	70	25500	8.4E-09	8.80E-01	8.7E-08	3%	1400	1.8E-07	8.0E+00	0.0E+00	0%
1,4-DDT	1.8E-01	1E-08	4000	1	0.001	250	4	70	25500	3.0E-08	8.80E-01	2.4E-09	1%	1400	6.3E-08	2.9E-04	2.8E-04	2%
Anthracene	2.1E-01	1E-08	4000	1	0.001	250	4	70	25500	4.5E-08	0.00E+00	0.0E+00	0%	1400	8.8E-08	1.8E-01	5.8E-07	0%
benzo(a)anthracene	2.8E-01	1E-08	4000	1	0.001	250	4	70	25500	6.7E-08	1.4E+00	9.8E-09	0%	1400	1.2E-07	8.0E+00	0.0E+00	0%
benzo(a)pyrene	2.8E-01	1E-08	4000	1	0.001	250	4	70	25500	6.7E-08	1.4E+00	9.8E-09	0%	1400	1.2E-07	8.0E+00	0.0E+00	0%
benzo(b)fluoranthene	2.7E-01	1E-08	4000	1	0.001	250	4	70	25500	8.4E-09	1.4E+01	8.4E-09	54%	1400	1.1E-07	8.0E+00	0.0E+00	0%
benzo(g,h,i)perylene	2.9E-01	1E-08	4000	1	0.001	250	4	70	25500	6.1E-09	0.00E+00	0.0E+00	0%	1400	1.2E-07	8.0E+00	0.0E+00	0%
benzofuran	1.7E-01	1E-08	4000	1	0.001	250	4	70	25500	4.1E-09	4.0E-02	1.6E-10	0%	1400	7.2E-08	8.0E+00	0.0E+00	0%
Chrysene	2.8E-01	1E-08	4000	1	0.001	250	4	70	25500	8.7E-09	1.4E-02	9.8E-11	0%	1400	1.3E-07	8.0E+00	0.0E+00	0%
Heptachlor epoxide	6.0E-03	1E-08	4000	1	0.001	250	4	70	25500	1.7E-10	1.8E+01	3.0E-09	2%	1400	2.9E-08	8.0E+00	4.8E-04	4%
Indeno(1,2,3-cd)pyrene	2.8E-01	1E-08	4000	1	0.001	250	4	70	25500	8.1E-09	1.4E+00	8.9E-09	0%	1400	1.1E-07	8.0E+00	0.0E+00	0%
Phenanthrene	2.8E-01	1E-08	4000	1	0.001	250	4	70	25500	8.6E-09	0.00E+00	0.0E+00	0%	1400	1.2E-07	1.6E-02	8.0E-03	0%
alpha-Chlordane	3.8E-02	1E-08	4000	1	0.001	250	4	70	25500	9.2E-10	2.0E+00	2.4E-09	1%	1400	1.6E-08	3.0E-05	8.4E-04	6%
gamma-Chlordane	1.3E-02	1E-08	4000	1	0.001	250	4	70	25500	3.1E-10	2.0E+00	8.1E-10	0%	1400	8.4E-09	3.0E-05	1.8E-04	2%
TOTAL										1.7E-07					1.1E-02			

SURFACE SOIL PARTICULATE INHALATION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 26)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT MILITARY PERSONNEL

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (C * EF * ED * IR * 1/PEF) / (BW * ATc or ATnc * DY)$$

Risk = Intake \* CSF or /RfD

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	Calculated
CSF = carcinogenic slope factor	Specific
RfD = reference dose for noncarcinogen	Specific
IR = inhalation rate (m³)	30
EF = adult exposure frequency (days)	250
ED = adult exposure duration (years)	4
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	4
DY = day per year (day/yr)	365
PEF = particulate emission factor (m³/kg)	4.63E+09

Note: Inputs are scenario and site specific

COPC	Concentration Carcinogen (mg/kg)	Particulate Emission Factor (m³/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m³/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Manganese	3.4E+03	4.6E+09	250	30	4	70	25550	1.2E-08	4.0E+00	0.0E+00	0%	1460	2.2E-07	0.00E+00	0.0E+00	0%
Antimony	5.7E+00	4.6E+09	250	30	4	70	25550	2.1E-11	0.0E+00	0.0E+00	0%	1460	3.6E-10	0.00E+00	0.0E+00	0%
Arsenic	1.7E+00	4.6E+09	250	30	4	70	25550	6.5E-12	1.5E+01	9.6E-11	92%	1460	1.1E-10	0.00E+00	0.0E+00	0%
Barium	3.6E+01	4.6E+09	250	30	4	70	25550	1.3E-10	0.0E+00	0.0E+00	0%	1460	2.2E-09	1.43E-04	1.5E-05	99%
Dcadmium (soil)	1.0E+00	4.6E+09	250	30	4	70	25550	5.6E-12	0.0E+00	0.0E+00	0%	1460	9.8E-11	0.00E+00	0.0E+00	0%
Chromium	1.0E+01	4.6E+09	250	30	4	70	25550	3.8E-11	0.0E+00	0.0E+00	0%	1460	6.6E-10	0.00E+00	0.0E+00	0%
Cobalt	9.1E-01	4.6E+09	250	30	4	70	25550	3.3E-12	0.0E+00	0.0E+00	0%	1460	5.8E-11	0.00E+00	0.0E+00	0%
Copper	1.3E+02	4.6E+09	250	30	4	70	25550	4.7E-10	0.0E+00	0.0E+00	0%	1460	8.2E-09	0.00E+00	0.0E+00	0%
Lead	1.7E+02	4.6E+09	250	30	4	70	25550	6.1E-10	0.0E+00	0.0E+00	0%	1460	1.1E-08	0.00E+00	0.0E+00	0%
Manganese (soil)	3.3E+02	4.6E+09	250	30	4	70	25550	1.2E-09	0.0E+00	0.0E+00	0%	1460	2.1E-08	0.00E+00	0.0E+00	0%
Mercury	2.8E-01	4.6E+09	250	30	4	70	25550	1.0E-12	0.0E+00	0.0E+00	0%	1460	1.8E-11	8.57E-05	2.1E-07	1%
Nickel	6.6E+00	4.6E+09	250	30	4	70	25550	2.0E-11	0.0E+00	0.0E+00	0%	1460	3.5E-10	0.00E+00	0.0E+00	0%
Silver	1.1E+00	4.6E+09	250	30	4	70	25550	4.1E-12	0.0E+00	0.0E+00	0%	1460	7.2E-11	0.00E+00	0.0E+00	0%
Thallium	6.1E-01	4.6E+09	250	30	4	70	25550	1.0E-12	0.0E+00	0.0E+00	0%	1460	3.3E-11	0.00E+00	0.0E+00	0%
Vanadium	7.9E+00	4.6E+09	250	30	4	70	25550	2.9E-11	0.0E+00	0.0E+00	0%	1460	5.0E-10	0.00E+00	0.0E+00	0%
Zinc	1.1E+03	4.6E+09	250	30	4	70	25550	3.9E-09	0.0E+00	0.0E+00	0%	1460	6.8E-08	0.00E+00	0.0E+00	0%
1,4'-DDD	6.4E-02	4.6E+09	250	30	4	70	25550	2.3E-13	0.0E+00	0.0E+00	0%	1460	4.1E-12	0.00E+00	0.0E+00	0%
1,4'-DDE	3.6E-01	4.6E+09	250	30	4	70	25550	1.3E-12	0.0E+00	0.0E+00	0%	1460	2.2E-11	0.00E+00	0.0E+00	0%
1,4'-DDT	1.6E-01	4.6E+09	250	30	4	70	25550	5.4E-13	3.4E-01	1.8E-13	0%	1460	9.4E-12	0.00E+00	0.0E+00	0%
Anthracene	2.1E-01	4.6E+09	250	30	4	70	25550	7.4E-13	0.0E+00	0.0E+00	0%	1460	1.3E-11	0.00E+00	0.0E+00	0%
Benz(a)anthracene	2.0E-01	4.6E+09	250	30	4	70	25550	1.0E-12	8.1E-01	6.1E-13	1%	1460	1.8E-11	0.00E+00	0.0E+00	0%
Benz(a)pyrene	2.7E-01	4.6E+09	250	30	4	70	25550	9.7E-13	8.10E+00	6.9E-12	6%	1460	1.7E-11	0.00E+00	0.0E+00	0%
Benz(b)fluoranthene	2.9E-01	4.6E+09	250	30	4	70	25550	1.1E-12	6.10E-01	6.8E-13	1%	1460	1.9E-11	0.00E+00	0.0E+00	0%
Benz(g,h,i)perylene	2.8E-01	4.6E+09	250	30	4	70	25550	9.2E-13	0.0E+00	0.0E+00	0%	1460	1.6E-11	0.00E+00	0.0E+00	0%
Carbazole	1.7E-01	4.6E+09	250	30	4	70	25550	6.2E-13	0.0E+00	0.0E+00	0%	1460	1.1E-11	0.00E+00	0.0E+00	0%
Chrysene	2.8E-01	4.6E+09	250	30	4	70	25550	1.0E-12	6.10E-03	6.2E-15	0%	1460	1.8E-11	0.00E+00	0.0E+00	0%
Heptachlor epoxide	6.9E-03	4.6E+09	250	30	4	70	25550	2.6E-14	9.10E+00	2.3E-13	0%	1460	4.4E-13	0.00E+00	0.0E+00	0%
Indeno(1,2,3-cd)pyrene	2.8E-01	4.6E+09	250	30	4	70	25550	9.2E-13	6.10E-01	5.6E-13	1%	1460	1.6E-11	0.00E+00	0.0E+00	0%
Phenanthrene	2.8E-01	4.6E+09	250	30	4	70	25550	1.0E-12	0.0E+00	0.0E+00	0%	1460	1.8E-11	0.00E+00	0.0E+00	0%
alpha-Chlordane	3.8E-02	4.6E+09	250	30	4	70	25550	1.4E-13	1.20E+00	1.8E-13	0%	1460	2.4E-12	0.00E+00	0.0E+00	0%
gamma-Chlordane	1.3E-02	4.6E+09	250	30	4	70	25550	4.7E-14	1.20E+00	6.0E-14	0%	1460	8.2E-13	0.00E+00	0.0E+00	0%
TOTAL												1.0E-10				1.6E-05

SEDIMENT INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B - ODE POND)  
 REMEDIAL INVESTIGATION CTD-0201  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT MILITARY PERSONNEL.

Intake from ingestion of sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * IR * CF * EF * ED / BW * ATC \text{ or } ATnc * DY$$

Risk = Intake \* CSF or RfD

Where:

C = contaminant concentration in sediment (mg/kg)

CF = conversion factor (kg to mg)

INPUTS

1E-08

EF = exposure frequency (days/yr)

250

ED = exposure duration (yr)

4

IR = soil ingestion rate (mg/day)

100

BW = body weight (kg)

70

ATC = averaging time for carcinogen (yr)

70

ATnc = averaging time for noncarcinogen (yr)

4

DY = days per year (days/year)

365

CSF = cancer slope factor (mg/kg-day)-1

Specific

RfD = reference dose (mg/kg-day)

Specific

Note: Inputs are scenario and site specific

ODE POND

Contaminant	Concentration (mg/kg)	Exposure Frequency (days/yr)	Exposure Duration (yr)	Ingestion Rate (mg/day)	Conversion Factor (kg/mg)	Body Weight (kg)	Average Carc Time (days)	Carc. Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Average Noncarc Time (days)	Noncarc. Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Antimony	4.0E+03	250	4	100	1E-08	70	25500	2.7E-04	0.00E+00	0.0E+00	0%	1400	4.0E-03	1.0E-03	4.0E-03	17%
Arsenic	6.4E+03	250	4	100	1E-08	70	25500	3.6E-07	1.7E-06	0.0E-07	0%	1400	6.3E-03	3.0E-04	2.1E-02	70%
Beryllium	3.2E+01	250	4	100	1E-08	70	25500	1.0E-08	4.0E-06	7.7E-08	11%	1400	3.1E-07	5.0E-03	6.3E-05	0%
Chromium	1.2E+01	250	4	100	1E-08	70	25500	9.6E-07	0.00E+00	0.0E+00	0%	1400	1.2E-05	1.0E+00	1.2E-05	0%
Cobalt	1.7E+00	250	4	100	1E-08	70	25500	9.6E-08	0.00E+00	0.0E+00	0%	1400	1.7E-05	6.0E-02	2.0E-05	0%
Copper	1.7E+00	250	4	100	1E-08	70	25500	9.0E-08	0.00E+00	0.0E+00	0%	1400	1.7E-05	3.7E-02	4.5E-05	0%
Lead	6.3E+03	250	4	100	1E-08	70	25500	4.0E-07	0.00E+00	0.0E+00	0%	1400	9.1E-08	0.00E+00	0.0E+00	0%
Manganese (soil)	9.0E+03	250	4	100	1E-08	70	25500	6.0E-07	0.00E+00	0.0E+00	0%	1400	9.0E-08	1.4E-01	9.0E-05	0%
Nickel	2.2E+03	250	4	100	1E-08	70	25500	1.2E-07	0.00E+00	0.0E+00	0%	1400	2.2E-08	2.0E-02	1.1E-04	0%
Vanadium	1.2E+01	250	4	100	1E-08	70	25500	6.4E-07	0.00E+00	0.0E+00	0%	1400	1.1E-05	7.0E-03	1.0E-03	0%
U-238	8.3E-03	250	4	100	1E-08	70	25500	4.0E-10	1.1E-10	0%	1400	8.1E-09	0.00E+00	0.0E+00	0%	
TOTAL									7.0E-07						2.0E-02	

SEDIMENT DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO.7 (SITE 2B - ODE POND)  
 REMEDIAL INVESTIGATION CTD-001  
 MCS CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT MILITARY PERSONNEL

The intake from dermal contact to sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

Risk = Intake  $\times$  CSF or RD

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	1.00E-05
CF = conversion factor (kg/mg)	4300
SA = exposed skin surface area (cm <sup>2</sup> )	1
AF = sediment to skin adherence factor (mp/cm <sup>2</sup> )	Specific
Abs = fraction absorbed (unless) (contaminant specific)	250
EF = exposure frequency (events/yr)	4
ED = exposure duration (year)	70
BW = body weight (kg)	70
ATc = averaging time for carcinogen (yr)	4
ATnc = averaging time for noncarcinogen (yr)	365
DY = day per year (day/yr)	Specific
CSF = cancer slope factor (mg/kg-day) <sup>-1</sup>	Specific
RD = reference dose (mg/kg-day)	Specific

Note: Inputs are scenario and site specific

ODE POND

COPC	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> )	Adherence Factor (mg/m <sup>2</sup> )	AF/S Factor (%)	Exposure Frequency (events/yr)	Exposure Duration (yr)	Body Weight (kg)	Average Conc Time (days)	Carc. Dose (mg/kg/day)	Dermal Adapt Slope Factor (mg/kg-day) <sup>-1</sup>	Carcinogenic Risk	Percent Carcinogenic Risk	Average Noncarc Time (days)	Reference Dose (mg/kg/day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Arsenic	4.0E+03	1E-05	4300	1	0.001	250	4	70	25500	1.4E-05	8.7E+00	0%	1400	2.1E-04	2.00E+01	1.1E-03	11%	
Antimony	6.4E+03	1E-05	4300	1	0.001	250	4	70	25500	1.4E-05	8.7E+00	0%	1400	6.0E-05	4.00E+01	4.4E-03	70%	
Beryllium	3.2E-01	1E-05	4300	1	0.001	250	4	70	25500	7.7E-10	2.1E+01	1.7E-08	11%	1400	1.3E-08	1.00E+03	1.3E-05	0%
Chromium	1.2E+01	1E-05	4300	1	0.001	250	4	70	25500	2.9E-08	0.005+00	0.05+00	0%	1400	6.0E-09	2.00E+01	2.00E-09	0%
Cobalt	1.7E+00	1E-05	4300	1	0.001	250	4	70	25500	4.1E-08	0.005+00	0.05+00	0%	1400	7.5E-08	1.20E+03	6.0E-08	0%
Copper	1.7E+00	1E-05	4300	1	0.001	250	4	70	25500	2.0E-08	0.005+00	0.05+00	0%	1400	7.2E-09	7.40E+03	8.4E-09	0%
Lead	8.3E+00	1E-05	4300	1	0.001	250	4	70	25500	2.4E-08	0.005+00	0.05+00	0%	1400	4.1E-07	0.005+00	0.05+00	0%
Manganese (soil)	9.8E+00	1E-05	4300	1	0.001	250	4	70	25500	8.3E-09	0.005+00	0.05+00	0%	1400	8.3E-08	4.00E+03	2.2E-09	0%
Nickel	2.2E+00	1E-05	4300	1	0.001	250	4	70	25500	2.8E-08	0.005+00	0.05+00	0%	1400	4.8E-07	1.40E+03	3.8E-04	0%
Vanadium	1.2E+01	1E-05	4300	1	0.001	250	4	70	25500	2.0E-10	4.00E-01	8.6E-11	0%	1400	3.8E-08	0.005+00	0.0E+00	0%
TOTAL																		

SURFACE WATER INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO.7 (SITE 26 - OZONE POND)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCAMP LEJEUNE, NORTH CAROLINA  
 CURRENT MILITARY PERSONNEL

The intake from the ingestion of surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = \text{Cw} * \text{CR} * \text{ET} * \text{EF} * \text{ED/BW} * \text{ATc or ATnc} * \text{DY}$$

$$\text{Risk} = \text{Intake} * \text{CSF or RID}$$

CHANGE!!!!

Where:	INPUT
Cw = contaminant concentration in surface water (mg/l)	
CR = contact rate (Lhr/hour)	0.005
ET = exposure time (hours/event)	2.6
EF = exposure frequency (events/yr)	45
ED = exposure duration (yr)	4
BW = body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	4
DY = days per year (days)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RID = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

COPC	Concentration (mg/l)	Contact Rate (hour)	Exposure Time (hrs/event)	Exposure Frequency (events/yr)	Exposure Duration (years)	Body Weight (kg)	Averaging Cenc. Time (days)	Carc. Dose (mg/kg-day)	Cancer Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Noncarcinogenic Risk	Averaging Time Noncarc. (days)	Reference Dose (mg/kg-day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Naphthalene	1.7E-01	0.005	2.6	45	4	70	25550	2.2E-07	0.00E+00	0%	1460	3.0E-06	1.0E+00	3.0E-06	0%	
Total	1.3E-02	0.005	2.6	45	4	70	25550	1.7E-08	0.00E+00	0%	1460	2.9E-07	2.0E-02	1.0E-05	1%	
Thallium	4.7E-03	0.005	2.6	45	4	70	25550	6.1E-09	0.00E+00	0%	1460	1.1E-07	8.0E-05	1.3E-03	98%	

TOTAL

SURFACE WATER DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO.7 (SITE 2B - DRIDE POND)  
 REMEDIAL INVESTIGATION - CTO-K0207  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT MILITARY PERSONNEL

The intake from dermal contact with surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = Cw \times SA \times PC \times ET \times EF \times ED \times CF/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times \text{CSF or RID}$$

Where:	INPUTS
CW = contaminant concentration in water (mg/l)	
SA = skin surface available for contact (cm <sup>2</sup> )	4300
PC = contaminant specific dermal permeability (cm/hr)	Specific
ET = exposure time (hours/day)	2.6
EF = exposure frequency (days/yr)	45
ED = exposure duration (years)	4
CF = volumetric conversion factor for water (liter/1000 cm <sup>3</sup> )	0.001
BW = body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	4
DY = days per year (days)	365
CSF = cancer slope factor (mg/kg-day)-1	Specific
RID = reference dose (mg/kg-day)	Specific

Note: Inputs are site and scenario specific

POPC	Concentration (mg/l)	Surface Area (cm <sup>2</sup> )	Dermal Permeability (cm/hr)	Exposure Time (hours/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	Volumetric Conversion (L/m <sup>3</sup> )	Body Weight (kg)	Averaging/Carc Time (day)	Carc. Dose (mg/kg-day)	Dermal Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Averaging Noncarc Time (day)	Noncarc. Dose (mg/kg-day)	Derm. Adjust Reference Dose (mg/kg-day)	Noncarc. Risk	Percent Noncarcinogenic Risk
Nomium	0.017	4300	1.0E-03	2.6	45	4	0.001	70	26600	1.1E-07	0.001+00	0%	1460	0.0E+00	2.0E-07	1.7E-03	0%	
Nickel	0.0129	4300	1.0E-04	2.6	45	4	0.001	70	26600	1.4E-09	0.005+00	0%	1460	2.0E-08	4.0E-09	0.3E-03	0%	
Thallium	0.0047	4300	1.0E-03	2.6	45	4	0.001	70	25500	8.3E-08	0.0E+00	0%	1460	0.0E+00	1.0E-03	0.0E+00	100%	
TOTAL																		

## SUBSURFACE SOIL INGESTION EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 2B)

REMEDIAl INVESTIGATION CTO-0231

MCB CAMP LEJEUNE, NORTH CAROLINA

FUTURE CONSTRUCTION WORKER

Intake from ingestion of soil is calculated as follows:

Intake (mg/kg-day) = C \* CF \* EF \* ED \* IR/BW \* ATc or ATnc \* DY

Risk = Intake \* CSF or /RfD

Where:

C = contaminant concentration in soil (mg/kg)

CF = conversion for kg to mg

EF = adult exposure frequency (days/yr)

ED = adult exposure duration (yr)

IR = adult soil ingestion rate (mg/day)

BW = adult body weight (kg)

ATc = averaging time for carcinogen (yr)

ATnc = averaging time for noncarcinogen (yr)

DY = days per year (days/year)

CSF = cancer slope factor (mg/kg-day)-1

RfD = reference dose (mg/kg-day)

## INPUTS

1E-06

90

1

480

70

25550

1.9E-06

0.0E+00

0%

365

1.3E-02

1.0E+00

1.3E-02

24%

5.0E-02

22%

4.8E-02

1%

1.9E-03

0%

5.0E-03

4.9E-05

0%

3%

6.4E-03

0%

5.8E-05

0%

7.8E-05

0%

0%

6.0E-02

0%

7.8E-05

0%

0%

0.0E+00

0%

1.4E-02

7%

1%

1.3E-03

0%

2.0E-03

1%

3.8E-07

0.0E+00

0.0E+00

0%

2.7E-06

0.0E+00

0.0E+00

0%

2.0E-07

0.0E+00

0.0E+00

0%

5.0E-07

0.0E+00

0.0E+00

0%

4.0E-04

0%

4.0E-02

0%

3.0E-02

0%

0.0E+00

0%

0.0E+00

0%

1.2E-06

0%

4.0E-02

0%

1.4E-06

0%

3.0E-05

0%

3.3E-04

0%

1.9E-04

0%

0.0E+00

0%

2.0E-06

0%

6.0E-05

0%

1.9E-04

0%

0.0E+00

0%

5.0E-07

0.0E+00

0.0E+00

0%

2.1E-01

0%

365

1.1E-08

0.0E+00

0.0E+00

0%

2.1E-01

0%

365

5.0E-07

0.0E+00

0.0E+00

0%

4.0E-07

0.0E+00

0.0E+00

0%

1.2E-06

0%

4.0E-02

0%

1.4E-06

0%

3.0E-05

0%

3.3E-04

0%

1.9E-04

0%

0.0E+00

0%

0.0E+00

0%

5.0E-07

0.0E+00

0.0E+00

0%

2.1E-01

0%

365

5.0E-07

0.0E+00

0.0E+00

0%

4.0E-07

0.0E+00

0.0E+00

0%

1.2E-06

0%

4.0E-02

0%

1.4E-06

0%

3.0E-05

0%

3.3E-04

0%

1.9E-04

0%

0.0E+00

0%

0.0E+00

0%

5.0E-07

0.0E+00

0.0E+00

0%

2.1E-01

0%

365

5.0E-07

0.0E+00

0.0E+00

0%

4.0E-07

0.0E+00

0.0E+00

0%

1.2E-06

0%

4.0E-02

0%

1.4E-06

0%

3.0E-05

0%

3.3E-04

0%

1.9E-04

0%

0.0E+00

0%

0.0E+00

0%

5.0E-07

0.0E+00

0.0E+00

0%

2.1E-01

0%

365

5.0E-07

0.0E+00

0.0E+00

0%

4.0E-07

0.0E+00

0.0E+00

0%

1.2E-06

0%

4.0E-02

0%

1.4E-06

0%

3.0E-05

0%

3.3E-04

0%

1.9E-04

0%

0.0E+00

0%

0.0E+00

0%

5.0E-07

0.0E+00

0.0E+00

0%

2.1E-01

0%

365

5.0E-07

0.0E+00

0.0E+00

0%

4.0E-07

0.0E+00

0.0E+00

0%

1.2E-06

0%

4.0E-02

0%

1.4E-06

0%

3.0E-05

0%

3.3E-04

0%

1.9E-04

0%

0.0E+00

0%

0.0E+00

0%

5.0E-07

0.0E+00

0.0E+00

0%

2.1E-01

0%

365

5.0E-07

0.0E+00

0.0E+00

0%

4.0E-07

0.0E+00

0.0E+00

0%

1.2E-06

0%

4.0E-02

0%

1.4E-06

0%

3.0E-05

0%

3.3E-04

0%

1.9E-04

0%

0.0E+00

0%

0.0E+00

0%

5.0E-07

0.0E+00

0.0E+00

0%

2.1E-01

0%

365

5.0E-07

0.0E+00

0.0E+00

0%

4.0E-07

0.0E+00

## SUBSURFACE SOIL DERMAL CONTACT EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 2B)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE CONSTRUCTION WORKER

Dermal contact with soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RID$$

Where:

C = contaminant concentration in soil (mg/kg)	1E-06	INPUTS
CF = conversion factor (kg/mg)	4300	
SA = adult exposed skin surface area (cm <sup>2</sup> )	1	
AF = soil to skin adherence factor (mg/cm <sup>2</sup> )	Specific	
Abs = fraction absorbed (unless)	90	
EF = adult exposure frequency (events/yr)	1	
ED = adult exposure duration (years)	70	
BW = adult body weight (kg)	1	
ATc = averaging time for carcinogen (yr)	70	
ATnc = averaging time for noncarcinogen (yr)	1	
DY = day per year (day/yr)	365	
CSF = cancer slope factor (mg/kg-day)-1	specific	
RID = reference dose (mg/kg-day)	specific	

Note: Inputs are scenario and site specific

POPC	Concentration Carcinogen (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> )	Adherence Factor (mg/cm <sup>2</sup> )	Fraction Absorbed (%)	Exposure Frequency (events/yr)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg/day)	Dermal Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	7.8E+03	1E-06	4300	1	0.001	90	1	70	25550	1.7E-06	0.0E+00	0%	365	1.2E-04	2.0E-01	5.9E-04	6%	
Antimony	1.2E+01	1E-06	4300	1	0.001	90	1	70	25550	2.6E-09	0.0E+00	0%	365	1.8E-07	0.0E-05	2.2E-03	24%	
Aseno	7.9E+00	1E-06	4300	1	0.001	90	1	70	25550	1.7E-09	8.8E+00	1.5E-08	29%	365	1.2E-07	0.0E-05	2.0E-03	21%
Barium	7.8E+01	1E-06	4300	1	0.001	90	1	70	25550	1.7E-08	0.0E+00	0.0E+00	0%	365	1.2E-06	1.4E-02	8.8E-05	1%
Beryllium	1.4E-01	1E-06	4300	1	0.001	90	1	70	25550	3.1E-11	2.1E+01	8.7E-10	1%	365	2.2E-09	1.0E-03	2.2E-06	0%
Cadmium (soil)	3.2E+00	1E-06	4300	1	0.001	90	1	70	25550	6.9E-10	0.0E+00	0.0E+00	0%	365	4.9E-08	2.0E-04	2.4E-04	3%
Chromium	3.4E+01	1E-06	4300	1	0.001	90	1	70	25550	7.4E-09	0.0E+00	0.0E+00	0%	365	6.2E-07	2.0E-01	2.8E-06	0%
Cobalt	2.7E+00	1E-06	4300	1	0.001	90	1	70	25550	6.8E-10	0.0E+00	0.0E+00	0%	365	4.1E-08	1.2E-02	3.4E-06	0%
Copper	8.9E+02	1E-06	4300	1	0.001	90	1	70	25550	1.9E-07	0.0E+00	0.0E+00	0%	365	1.3E-06	7.4E-03	1.8E-03	19%
Lead	1.7E+03	1E-06	4300	1	0.001	90	1	70	25550	3.7E-07	0.0E+00	0.0E+00	0%	365	2.6E-06	0.0E+00	0.0E+00	0%
Manganese (soil)	1.2E+03	1E-06	4300	1	0.001	90	1	70	25550	2.6E-07	0.0E+00	0.0E+00	0%	365	1.8E-06	2.8E-02	6.4E-04	7%
Mercury	2.3E-01	1E-06	4300	1	0.001	90	1	70	25550	4.9E-11	0.0E+00	0.0E+00	0%	365	3.5E-09	8.0E-05	5.8E-05	1%
Nickel	2.3E+01	1E-06	4300	1	0.001	90	1	70	25550	5.0E-09	0.0E+00	0.0E+00	0%	365	3.5E-07	4.0E-03	8.8E-05	1%
Silver	2.8E+00	1E-06	4300	1	0.001	90	1	70	25550	5.4E-10	0.0E+00	0.0E+00	0%	365	3.8E-08	1.0E-03	3.8E-05	0%
Vanadium	2.4E+01	1E-06	4300	1	0.001	90	1	70	25550	6.3E-09	0.0E+00	0.0E+00	0%	365	3.7E-07	1.4E-03	2.6E-04	3%
Zinc	4.3E+03	1E-06	4300	1	0.001	90	1	70	25550	9.4E-07	0.0E+00	0.0E+00	0%	365	6.6E-06	6.0E-02	1.1E-03	12%
2-Methylnaphthalene	8.9E-02	1E-06	4300	1	0.01	90	1	70	25550	1.9E-10	0.0E+00	0.0E+00	0%	365	1.3E-06	2.0E-02	6.7E-07	0%
4,4'-DDD	8.8E-01	1E-06	4300	1	0.01	90	1	70	25550	1.9E-09	4.8E-01	9.1E-10	2%	365	1.3E-07	0.0E+00	0.0E+00	0%
4,4'-DDE	1.6E+00	1E-06	4300	1	0.01	90	1	70	25550	3.5E-09	6.8E-01	2.4E-09	5%	365	2.4E-07	0.0E+00	0.0E+00	0%
4,4'-DDT	1.2E-01	1E-06	4300	1	0.01	90	1	70	25550	2.6E-10	6.8E-01	1.7E-10	0%	365	1.8E-08	2.6E-04	7.1E-05	1%
Benzo(a)anthracene	5.9E-01	1E-06	4300	1	0.01	90	1	70	25550	1.3E-09	1.5E+00	1.8E-09	4%	365	8.8E-08	0.0E+00	0.0E+00	0%
Benzo(a)pyrene	5.8E-01	1E-06	4300	1	0.01	90	1	70	25550	1.2E-09	1.5E+01	1.8E-08	35%	365	8.5E-08	0.0E+00	0.0E+00	0%
Benzo(b)fluoranthene	6.2E-01	1E-06	4300	1	0.01	90	1	70	25550	1.3E-09	1.5E+00	1.8E-09	4%	365	9.3E-08	0.0E+00	0.0E+00	0%
Benzo(g,h)perylene	4.4E-01	1E-06	4300	1	0.01	90	1	70	25550	9.8E-10	0.0E+00	0.0E+00	0%	365	8.7E-08	1.5E-02	4.5E-06	0%
Benzo(k)fluoranthene	6.4E-01	1E-06	4300	1	0.01	90	1	70	25550	1.2E-09	1.5E-01	1.7E-10	0%	365	8.1E-08	0.0E+00	0.0E+00	0%
Bis(2-ethylhexyl)phthalate	4.9E-01	1E-06	4300	1	0.01	90	1	70	25550	1.0E-09	2.8E-02	2.9E-11	0%	365	7.2E-08	1.0E-02	7.2E-06	0%
Chrysene	5.7E-01	1E-06	4300	1	0.01	90	1	70	25550	1.2E-09	1.5E-02	1.8E-11	0%	365	8.7E-08	0.0E+00	0.0E+00	0%
Diben(a,h)anthracene	2.8E-01	1E-06	4300	1	0.01	90	1	70	25550	6.1E-10	1.5E+01	8.8E-09	17%	365	4.3E-08	0.0E+00	0.0E+00	0%
Fluorene	2.8E-01	1E-06	4300	1	0.01	90	1	70	25550	6.1E-10	0.0E+00	0.0E+00	0%	365	4.2E-08	2.0E-02	2.1E-06	0%
Indeno(1,2,3-cd)pyrene	4.3E-01	1E-06	4300	1	0.01	90	1	70	25550	9.4E-10	1.5E+00	1.4E-09	3%	365	6.6E-08	0.0E+00	0.0E+00	0%
Naphthalene	3.4E-01	1E-06	4300	1	0.01	90	1	70	25550	7.4E-10	0.0E+00	0.0E+00	0%	365	5.2E-08	2.0E-02	2.6E-06	0%
Phenanthrene	6.4E-01	1E-06	4300	1	0.01	90	1	70	25550	1.4E-09	0.0E+00	0.0E+00	0%	365	8.7E-08	1.5E-02	6.5E-06	0%
alpha-Chlordane	1.2E-02	1E-06	4300	1	0.01	90	1	70	25550	2.5E-11	2.6E+00	6.5E-11	0%	365	1.7E-09	3.0E-05	5.8E-05	1%
gamma-Chlordane	6.7E-03	1E-06	4300	1	0.01	90	1	70	25550	1.4E-11	2.6E+00	3.8E-11	0%	365	1.0E-09	3.0E-05	3.4E-05	0%
TOTAL										5.1E-08						9.3E-03		

SUBSURFACE SOIL PARTICULATE INHALATION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 2B)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE CONSTRUCTION WORKER

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (C * EF * ED * IR * I/PEF) / (BW * ATc or ATnc * DY)$$

Risk = Intake \* CSF or /RfD

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	Specific
CSF = carcinogenic slope factor	Specific
RfD = reference dose for noncarcinogen	Specific
IR = inhalation rate (m3)	20
EF = adult exposure frequency (days)	90
ED = adult exposure duration (years)	1
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	1
DY = day per year (day/yr)	365
PEF = particulate emission factor (m3/kg)	4.63E+09

Note: Inputs are scenario and site specific

COPC	Concentration Carcinogen (mg/kg)	Particulate Emission Factor (m3/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m3/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Aluminum	7.8E+03	4.6E+09	90	20	1	70	25550	1.7E-07	0.0E+00	0%	365	1.2E-07	0.0E+00	0.0E+00	0%	
Antimony	1.2E+01	4.6E+09	90	20	1	70	25550	2.6E-12	0.0E+00	0%	365	1.8E-10	0.0E+00	0.0E+00	0%	
Arsenic	7.9E+00	4.6E+09	90	20	1	70	25550	1.7E-12	1.5E+01	94%	365	1.2E-10	0.0E+00	0.0E+00	0%	
Barium	7.8E+01	4.6E+09	90	20	1	70	25550	1.7E-11	0.0E+00	0%	365	1.2E-09	1.4E-04	8.3E-06	100%	
Beryllium	1.4E-01	4.6E+09	90	20	1	70	25550	3.1E-14	8.4E+00	2%	365	2.2E-12	0.0E+00	0.0E+00	0%	
Cadmium (soil)	3.2E+00	4.6E+09	90	20	1	70	25550	7.0E-13	0.0E+00	0%	365	4.9E-11	0.0E+00	0.0E+00	0%	
Chromium	3.4E+01	4.6E+09	90	20	1	70	25550	7.4E-12	0.0E+00	0%	365	5.2E-10	0.0E+00	0.0E+00	0%	
Cobalt	2.7E+00	4.6E+09	90	20	1	70	25550	5.8E-13	0.0E+00	0%	365	4.1E-11	0.0E+00	0.0E+00	0%	
Copper	8.9E+02	4.6E+09	90	20	1	70	25550	1.9E-10	0.0E+00	0%	365	1.3E-08	0.0E+00	0.0E+00	0%	
Lead	1.7E+03	4.6E+09	90	20	1	70	25550	3.7E-10	0.0E+00	0%	365	2.6E-06	0.0E+00	0.0E+00	0%	
Manganese (soil)	1.2E+03	4.6E+09	90	20	1	70	25550	2.6E-10	0.0E+00	0%	365	1.8E-06	0.0E+00	0.0E+00	0%	
Mercury	2.3E+01	4.6E+09	90	20	1	70	25550	5.0E-14	0.0E+00	0%	365	3.5E-12	8.6E-05	4.0E-08	0%	
Nickel	2.3E+01	4.6E+09	90	20	1	70	25550	8.0E-12	0.0E+00	0%	365	3.5E-10	0.0E+00	0.0E+00	0%	
Silver	2.6E+00	4.6E+09	90	20	1	70	25550	6.4E-13	0.0E+00	0%	365	3.8E-11	0.0E+00	0.0E+00	0%	
Vanadium	2.4E+01	4.6E+09	90	20	1	70	25550	5.3E-12	0.0E+00	0%	365	3.7E-10	0.0E+00	0.0E+00	0%	
Zinc	4.3E+03	4.6E+09	90	20	1	70	25550	8.4E-10	0.0E+00	0%	365	6.6E-08	0.0E+00	0.0E+00	0%	
2-Methylnaphthalene	8.9E-02	4.6E+09	90	20	1	70	25550	1.0E-14	0.0E+00	0%	365	1.4E-12	0.0E+00	0.0E+00	0%	
4,4'-DDD	8.8E-01	4.6E+09	90	20	1	70	25550	1.0E-13	0.0E+00	0%	365	1.3E-11	0.0E+00	0.0E+00	0%	
4,4'-DDE	1.6E+00	4.6E+09	90	20	1	70	25550	3.6E-13	0.0E+00	0%	365	2.4E-11	0.0E+00	0.0E+00	0%	
4,4'-DDT	1.2E-01	4.6E+09	90	20	1	70	25550	2.6E-14	3.4E-01	0%	365	1.8E-12	0.0E+00	0.0E+00	0%	
Benzo(a)anthracene	5.8E-01	4.6E+09	90	20	1	70	25550	1.3E-13	6.1E-01	7.7E-14	0%	365	8.8E-12	0.0E+00	0.0E+00	0%
Benzo(a)pyrene	5.6E-01	4.6E+09	90	20	1	70	25550	1.2E-13	6.1E-00	7.6E-13	3%	365	8.6E-12	0.0E+00	0.0E+00	0%
Benzo(b)fluoranthene	6.2E-01	4.6E+09	90	20	1	70	25550	1.3E-13	6.1E-01	8.2E-14	0%	365	9.4E-12	0.0E+00	0.0E+00	0%
Benzo(g,h,i)perylene	4.4E-01	4.6E+09	90	20	1	70	25550	9.7E-14	0.0E+00	0.0E+00	0%	365	6.8E-12	0.0E+00	0.0E+00	0%
Benzo(k)fluoranthene	5.4E-01	4.6E+09	90	20	1	70	25550	1.2E-13	6.1E-02	7.1E-16	0%	365	8.1E-12	0.0E+00	0.0E+00	0%
Bis(2-ethylhexyl)phthalate	4.8E-01	4.6E+09	90	20	1	70	25550	1.0E-13	0.0E+00	0.0E+00	0%	365	7.3E-12	0.0E+00	0.0E+00	0%
Chrysene	6.7E-01	4.6E+09	90	20	1	70	25550	1.2E-13	8.1E-03	7.8E-16	0%	365	8.7E-12	0.0E+00	0.0E+00	0%
Dibenz(a,h)anthracene	2.9E-01	4.6E+09	90	20	1	70	25550	6.1E-14	6.1E+00	3.7E-13	1%	365	4.3E-12	0.0E+00	0.0E+00	0%
Fluorene	2.8E-01	4.6E+09	90	20	1	70	25550	6.1E-14	0.0E+00	0.0E+00	0%	365	4.3E-12	0.0E+00	0.0E+00	0%
Indeno(1,2,3-cd)pyrene	4.5E-01	4.6E+09	90	20	1	70	25550	9.6E-14	8.1E-01	6.6E-14	0%	365	6.6E-12	0.0E+00	0.0E+00	0%
Naphthalene	3.4E-01	4.6E+09	90	20	1	70	25550	7.6E-14	0.0E+00	0.0E+00	0%	365	5.2E-12	0.0E+00	0.0E+00	0%
Phenanthrene	6.4E-01	4.6E+09	90	20	1	70	25550	1.4E-13	0.0E+00	0.0E+00	0%	365	9.7E-12	0.0E+00	0.0E+00	0%
alpha-Chlordane	1.2E-02	4.6E+09	90	20	1	70	25550	2.6E-15	1.3E+00	3.2E-15	0%	365	1.0E-13	0.0E+00	0.0E+00	0%
beta-Chlordane	6.7E-03	4.6E+09	90	20	1	70	25550	1.6E-15	1.9E+00	1.9E-15	0%	365	8.4E-08	0.0E+00	0.0E+00	0%
<b>TOTAL</b>								2.0E-11								

**SITE 30**

SUBSURFACE SOIL INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 30)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL CHILD

Intake from ingestion of soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times EF \times ED \times IR/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RD$$

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	1E-06
CF = conversion for kg to mg	350
EF = child exposure frequency (days/yr)	6
ED = child exposure duration (yr)	200
IR = child soil ingestion rate (mg/day)	15
BW = child body weight (kg)	70
ATc = averaging time for carcinogen (yr)	8
ATnc = averaging time for noncarcinogen (yr)	365
DY = days per year (days/year)	specific
CSF = cancer slope factor (mg/kg-day)-1	specific
RD = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

COPC	Concentration Carcinogen (mg/kg)	Exposure Frequency (days/yr) Child	Exposure Duration (yr) Child	Conversion Factor (kg/mg)	Ingestion Rate (mg/day) Child	Body Weight (kg) Child	Averaging Carr Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Averaging Noncarc Time (days)	Noncarc Dose (mg/kg/day) Child	Reference Dose (mg/kg/day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child
Aluminum	9567.1	350	6	1E-06	200	15	25550	1.0E+02	0.0E+00	0%	2190	1.2E-01	1.0E+00	1.2E-01	68%	
Arsenic	0.9	350	6	1E-06	200	15	25550	9.7E-07	1.8E+00	1.7E-06	100%	2190	1.1E-05	3.0E-04	3.8E-02	21%
Chromium	10.1	350	6	1E-06	200	15	25550	1.1E-06	0.0E+00	0.0E+00	0%	2190	1.3E-04	1.0E+00	1.3E-04	0%
Cobalt	0.5	350	6	1E-06	200	15	25550	6.9E-07	0.0E+00	0.0E+00	0%	2190	6.9E-06	6.0E-02	1.1E-04	0%
Copper	3.5	350	6	1E-06	200	15	25550	3.9E-06	0.0E+00	0.0E+00	0%	2190	4.6E-05	3.7E-02	1.2E-03	1%
Manganese (soil)	6.8	350	6	1E-06	200	15	25550	7.4E-06	0.0E+00	0.0E+00	0%	2190	8.7E-05	1.4E-01	6.2E-04	0%
Mercury	0.1	350	6	1E-06	200	15	25550	1.4E-07	0.0E+00	0.0E+00	0%	2190	1.7E-06	3.0E-04	5.5E-03	3%
Nickel	2.2	350	6	1E-06	200	15	25550	2.4E-06	0.0E+00	0.0E+00	0%	2190	2.8E-05	2.0E-02	1.4E-03	1%
Vanadium	8.1	350	6	1E-06	200	15	25550	8.9E-06	0.0E+00	0.0E+00	0%	2190	1.0E-04	7.0E-03	1.5E-02	8%
TOTAL								1.7E-06							1.8E-01	

SUBSURFACE SOIL DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 30)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL CHILD

Dermal contact with soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * CF * SA * AF * Abs * EF * ED/BW * ATc \text{ or } ATnc * DY$$

Risk = Intake \* CSF or /RD

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	1E-06
CF = conversion factor (kg/mg)	2300
SA = child exposed skin surface area (cm <sup>2</sup> )	1
AF = soil to skin adherence factor (mg/cm <sup>2</sup> )	Specific
Abs = fraction absorbed (unitless)	350
EF = child exposure frequency (events/yr)	6
ED = child exposure duration (years)	15
BW = child body weight (kg)	15
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	8
DY = day per year (day/yr)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RD = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

POPC	Concentration Carcinogen (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> ) Child	Adherence Factor (mg/cm <sup>2</sup> )	Fraction Absorbed (%)	Exposure Frequency (events/yr) Child	Exposure Duration (yrs) Child	Body Weight (kg) Child	Average Carc Time (days)	Carc Dose (mg/kg/day) Child	Dermal Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Averaging Noncarc Time (days)	Noncarc Dose (mg/kg/day) Child	Dermal Adjust Reference Dose (mg/kg-day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child
Aluminum	3.6E+03	1E-06	2300	1	0.001	350	6	15	25550	1.2E-04	0.0E+00	0%	2190	1.4E-03	2.0E-01	7.0E-03	66%	
Anerobic	8.0E-01	1E-06	2300	1	0.001	350	6	15	25550	1.1E-06	8.8E+00	9.8E-08	100%	2190	1.3E-07	6.0E-05	2.2E-03	21%
Chromium	1.0E+01	1E-06	2300	1	0.001	350	6	15	25550	1.3E-07	0.0E+00	0.0E+00	0%	2190	1.5E-06	2.0E-01	7.4E-06	0%
Cobalt	5.4E-01	1E-06	2300	1	0.001	350	6	15	25550	6.8E-09	0.0E+00	0.0E+00	0%	2190	7.9E-06	1.2E-02	8.6E-06	0%
Copper	3.8E+00	1E-06	2300	1	0.001	350	6	15	25550	4.4E-08	0.0E+00	0.0E+00	0%	2190	8.2E-07	7.4E-03	7.0E-05	1%
Manganese (soil)	6.8E+00	1E-06	2300	1	0.001	350	6	15	25550	8.6E-08	0.0E+00	0.0E+00	0%	2190	1.0E-08	2.8E-02	3.6E-05	0%
Mercury	1.0E-01	1E-06	2300	1	0.001	350	6	15	25550	1.6E-09	0.0E+00	0.0E+00	0%	2190	1.9E-08	6.0E-05	3.2E-04	3%
Nickel	2.2E+00	1E-06	2300	1	0.001	350	6	15	25550	2.7E-08	0.0E+00	0.0E+00	0%	2190	3.2E-07	4.0E-03	8.0E-05	1%
Vanadium	8.1E+00	1E-06	2300	1	0.001	350	6	15	25550	1.0E-07	0.0E+00	0.0E+00	0%	2190	1.2E-06	1.4E-03	8.5E-04	8%
<b>TOTAL</b>										8.8E-08								1.1E-02

## SUBSURFACE SOIL PARTICULATE INHALATION EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 30)

REMEDIAl INVESTIGATION CTO-0231

MCB CAMP LEJEUNE, NORTH CAROLINA

FUTURE RESIDENTIAL CHILD

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (C \times EF \times ED \times IR \times 1/\text{PEF}) / (\text{BW} \times ATc \text{ or } ATnc \times DY)$$

$$\text{Risk} = \text{Intake} \times \text{CSF or } / \text{RID}$$

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	Calculated
CSF = carcinogenic slope factor	Specific
RID = reference dose for noncarcinogen	Specific
IR = inhalation rate (m3)	10
EF = child exposure frequency (days)	360
ED = child exposure duration (years)	6
BW = child body weight (kg)	15
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	6
DY = day per year (day/yr)	365
PEF = particulate emission factor (m3/kg)	4.63E+09

Note: Inputs are scenario and site specific

COPC	Concentration Carcogen (mg/kg)	Particulate Emission Factor (m3/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m3/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
As (dust)	9.0E+03	4.6E+09	360	10	6	15	25550	1.1E-07	0.0E+00	0%	2190	1.3E-06	0.0E+00	0.0E+00	0%	
Arsenic	8.9E-01	4.6E+09	360	10	6	15	25550	1.0E-11	1.6E+01	100%	2190	1.2E-10	0.0E+00	0.0E+00	0%	
Chromium	1.0E+01	4.6E+09	360	10	6	15	25550	1.2E-10	0.0E+00	0%	2190	1.4E-09	0.0E+00	0.0E+00	0%	
Cobalt	5.4E-01	4.6E+09	360	10	6	15	25550	6.3E-12	0.0E+00	0%	2190	7.4E-11	0.0E+00	0.0E+00	0%	
Copper	3.6E+00	4.6E+09	360	10	6	15	25550	4.2E-11	0.0E+00	0%	2190	4.9E-10	0.0E+00	0.0E+00	0%	
Manganese (soil)	6.6E+00	4.6E+09	360	10	6	15	25550	8.0E-11	0.0E+00	0%	2190	9.4E-10	0.0E+00	0.0E+00	0%	
Mercury	1.3E-01	4.6E+09	360	10	6	15	25550	1.6E-12	0.0E+00	0%	2190	1.6E-11	8.6E-06	2.1E-07	100%	
Nickel	2.2E+00	4.6E+09	360	10	6	15	25550	2.6E-11	0.0E+00	0%	2190	3.0E-10	0.0E+00	0.0E+00	0%	
Vanadium	8.1E+00	4.6E+09	360	10	6	15	25550	9.6E-11	0.0E+00	0%	2190	1.1E-09	0.0E+00	0.0E+00	0%	
<b>TOTAL</b>																
																<b>2.1E-07</b>

SURFACE WATER INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO.7 (SITE 30 - FRENCH CREEK)  
 REMEDIAL INVESTIGATION CTO-029  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT AND FUTURE CHILD RESIDENT

The intake from the ingestion of surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = \text{Cw} * \text{CR} * \text{ET} * \text{EF} * \text{ED/BW} * \text{ATc or ATnc} * \text{DY}$$

Risk = Intake \* CSF or /RDO

Where:

	INPUT
Cw = contaminant concentration in surface water (mg/l)	0.005
CR = contact rate (Liter/hour)	0.005
ET = child exposure time (hours/event)	2.6
EF = child exposure frequency (events/yr)	45
ED = child exposure duration (yr)	6
BW = child body weight (kg)	15
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	6
DY = days per year (days)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RDO = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

FRENCH CREEK

COPC	Concentration Carcinogen (mg/l)	Contact Rate (hour)	Exposure Time (hrs/event) Child	Exposure Frequency (events/yr) Child	Exposure Duration (year) Child	BW/BV Weight (kg) Child	Averaging Carr. Time (days) Child	Carc. Dose (mg/kg-day) Child	Cancer Slope Factor (mg/kg-day)-1	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Averaging Time Noncarc (days)	Noncarc. Dose (mg/kg-day) Child	Reference Dose (mg/kg-day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child
Nutrient	1.48	0.005	2.6	45	6	15	25550	1.4E-05	1.0E+00	0.0E+00	0%	2160	1.0E-04	1.0E+00	1.0E-04	35%
Lead	0.0023	0.005	2.6	45	6	15	25550	2.1E-06	0.0E+00	0.0E+00	0%	2160	2.6E-07	0.0E+00	0.0E+00	0%
Vaseline (water)	0.0111	0.005	2.6	45	6	15	25550	1.0E-07	0.0E+00	0.0E+00	0%	2160	1.2E-08	6.0E-03	2.4E-04	53%
Mercury	0.00015	0.005	2.6	45	6	15	25550	1.4E-08	0.0E+00	0.0E+00	0%	2160	1.0E-08	3.0E-04	5.3E-05	12%
UTL									0.0E+00					4.3E-04		

## SURFACE WATER DERMAL CONTACT EXPOSURE ASSESSMENT

OPERABLE UNIT NO.7 (SITE 30 - FRENCHS CREEK)

REMEDIAl INVESTIGATION - CTO-0201

VETS CAMP LEJEUNE, NORTH CAROLINA

CURRENT AND FUTURE CHILD RESIDENT

The intake from dermal contact with surface water is calculated as follows:

$$\text{intake (mg/kg-day)} = \text{Cw} \times \text{SA} \times \text{PC} \times \text{ET} \times \text{EF} \times \text{ED} \times \text{CF/BW} \times \text{ATc or ATnc} \times \text{DF}$$

$$\text{Risk} = \text{intake} \times \text{CSF or RfD}$$

INPUTS	
CW = contaminant concentration in water (mg/l)	
SA = child skin surface available for contact (cm <sup>2</sup> )	2300
PC = contaminant specific dermal permeability (cm/hr)	Specific
ET = child exposure time (hours/day)	2.6
EF = child exposure frequency (days/yr)	45
ED = child exposure duration (years)	6
CF = volumetric conversion factor for water (liter/1000 cm <sup>3</sup> )	0.001
BW = child body weight (kg)	15
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	6
DT = days per year (days)	365
CSF = cancer slope factor (mg/kg-day) <sup>-1</sup>	Specific
RfD = reference dose (mg/kg-day)	Specific

Note: Inputs are site and scenario specific

## FRENCHS CREEK

DPC	concentration Carcinogen (mg/l)	Surface Area (cm <sup>2</sup> )	Dermal Permeability (cm/hr)	Exposure Time (hours/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	Volumetric Conversion (l/m <sup>3</sup> )	body Weight (kg) Child	Averaging Carc Time (days)	Carc Dose (mg/kg-day) Child	Dermat Adjust Slope Factor (mg/kg-day) <sup>-1</sup>	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc. Time (days)	Noncarc. Dose (mg/kg-day) Child	Dermat Adjust Reference Dose (mg/kg-day)	Noncarc. Risk Child	Percent Noncarcinogenic Risk Child
Vaseline	1.0E+00	2300	1.0E-03	2.6	45	6	0.001	15	25000	6.2E-08	0.0E+00	0.0E+00	0%	2190	7.3E-05	2.0E-01	3.0E-04	30%
Lead	2.3E-03	2300	4.0E-05	2.6	45	6	0.001	15	25000	3.6E-11	0.0E+00	0.0E+00	0%	2190	4.5E-10	0.0E+00	0.0E+00	0%
Vanillin (water)	1.1E-02	2300	1.0E-03	2.6	45	6	0.001	15	25000	4.7E-08	0.0E+00	0.0E+00	0%	2190	5.6E-07	1.0E-03	5.6E-04	53%
Mercury	1.0E-04	2300	1.0E-03	2.6	45	6	0.001	15	25000	6.3E-10	0.0E+00	0.0E+00	0%	2190	7.4E-09	0.0E-05	1.2E-04	12%
OTAL										0.0E+00							1.0E-03	

SEDIMENT INGESTION EXPOSURE ASSESSMENT  
OPERABLE UNIT NO. 7 (SITE 30 - FRENCHS CREEK)  
REMEDIAL INVESTIGATION CTO-0231  
MCB CAMP LEJEUNE, NORTH CAROLINA  
CURRENT AND FUTURE RESIDENTIAL CHILD

Intake from ingestion of sediment is calculated as follows:

$$\text{intake (mg/kg-day)} = C \times IR \times CF \times ED / BW \times ATC \text{ or } ATnc \times DY$$

Risk = Intake \* CSF or /RD

Where:

C = contaminant concentration in sediment (mg/kg)  
CF = conversion for kg to mg  
EF = exposure frequency for child (days/yr)  
ED = exposure duration for child (yr)  
IR = soil ingestion rate for child (mg/day)  
BW = body weight for child (kg)  
ATC = averaging time for carcinogen (yr)  
ATnc = averaging time for noncarcinogen (yr)  
DY = days per year (days/year)  
CSF = cancer slope factor (mg/kg-day)-1  
RD = reference dose (mg/kg-day)

INPUTS

1E-06  
45  
6  
200  
15  
70  
5  
365  
Specific  
Specific

Note: Inputs are scenario and site specific

FRENCHS CREEK

Contaminant	Concentration (mg/kg)	Exposure Frequency (days/yr) Child	Exposure Duration (yr) Child	Ingestion Rate (mg/day) Child	Conversion Factor (kg/mg)	Body Weight (kg) Child	Average Caco Time (day)	Carc. Dose (mg/kg/day)-1 Child	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Child	Percent Carcinogenic Risk Child	Average Noncarc. Time (days)	Noncarc. Dose (mg/kg/day) Child	Reference Dose (mg/kg/day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk Child
ALUMINUM	3.4E+03	45	6	200	1E-06	15	25500	4.3E-04	0.0E+00	0%	2100	5.8E-03	1.0E+00	5.8E-03	0%	
Chromium	7.8E+00	45	6	200	1E-06	15	25500	1.1E-06	0.0E+00	0%	2100	1.3E-05	1.0E+00	1.3E-05	0%	
Copper	9.8E+00	45	6	200	1E-06	15	25500	1.3E-06	0.0E+00	0%	2100	1.6E-05	3.7E-02	4.2E-04	5%	
Lead	1.0E+01	45	6	200	1E-06	15	25500	2.2E-05	0.0E+00	0%	2100	2.8E-05	0.0E+00	0.0E+00	0%	
Manganese (soil)	1.4E+01	45	6	200	1E-06	15	25500	2.7E-05	0.0E+00	0%	2100	3.1E-05	1.4E-01	2.2E-04	3%	
Nickel	8.8E+00	45	6	200	1E-06	15	25500	8.2E-07	0.0E+00	0%	2100	9.8E-06	2.0E-02	4.8E-04	6%	
Vanadium	6.0E+00	45	6	200	1E-06	15	25500	8.8E-07	0.0E+00	0%	2100	9.8E-06	7.0E-03	1.4E-03	17%	
Zinc	1.1E+01	45	6	200	1E-06	15	25500	1.0E-06	0.0E+00	0%	2100	1.0E-05	3.0E-01	8.2E-05	1%	
<b>TOTAL</b>									0.0E+00							

SEDIMENT DERMAL CONTACT EXPOSURE ASSESSMENT  
OPERABLE UNIT NO.7 (SITE 30 - FRENCHS CREEK)  
REMEDIATION INVESTIGATION CTO-0201  
MCB CAMP LEJEUNE, NORTH CAROLINA  
CURRENT AND FUTURE RESIDENTIAL CHILD

The intake from dermal contact to sediment is calculated as follows:

$$\text{Intake (mg/kg day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

Risk = Intake  $\times$  CSF or RID

Where:

C = contaminant concentration in soil (mg/kg)

CF = conversion factor (kg/mg)

SA = child exposed skin surface area (cm<sup>2</sup>)

AF = sediment to skin adherence factor (mg/cm<sup>2</sup>)

Abs = fraction absorbed (unitless) (contaminant specific)

EF = child exposure frequency (events/yr)

ED = child exposure duration (years)

BW = child body weight (kg)

ATc = averaging time for carcinogen (yr)

ATnc = averaging time for noncarcinogen (yr)

DY = day per year (days/yr)

CSF = cancer slope factor (mg/kg-day)-1

RID = reference dose (mg/kg-day)

INPUTS

1.00E-08

2300

1

Specific

45

6

15

70

8

365

Specific

Specific

Note: Inputs are scenario and site specific

FRENCHS CREEK

Contaminant	Concen/Ref(2001) Carcinogen (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> ) Child	Adherence Factor (mg/cm <sup>2</sup> )	Ass Factor (%)	Exposure Frequency (events/yr) Child	Exposure Duration (yr) Child	Body Weight (kg) Child	Average Carc Time (days) Child	Carc. Dose (mg/kg/day) Child	Dermal Adjust. Slope Factor (mg/kg-day)-1	Carcinogenic Risk Child	Percent Carcinogenic Risk	Average Noncarc Time (days)	Noncarc. Dose (mg/kg/day) Child	Dermal Adjust. Reference Dose (mg/kg-day)	Noncarcinogenic Risk Child	Percent Noncarcinogenic Risk
Arsenium	3.4E+02	1E-08	2300	1	0.001	45	6	15	22500	3.3E-08	0.0E+00	0%	2100	6.3E-05	2.0E-01	3.2E-04	0%	
Chromium	7.8E+00	1E-08	2300	1	0.001	45	6	15	22500	1.3E-08	0.0E+00	0%	2100	1.6E-07	2.0E-01	7.6E-07	0%	
Copper	9.8E+00	1E-08	2300	1	0.001	45	6	15	22500	1.6E-08	0.0E+00	0%	2100	1.6E-07	7.4E-03	2.4E-05	0%	
Lead	1.8E+01	1E-08	2300	1	0.001	45	6	15	22500	2.8E-08	0.0E+00	0%	2100	2.8E-07	0.0E+00	0.0E+00	0%	
Manganese (soil)	1.9E+01	1E-08	2300	1	0.001	45	6	15	22500	3.1E-08	0.0E+00	0%	2100	3.0E-07	2.0E-02	1.3E-05	3%	
Nickel	8.8E+00	1E-08	2300	1	0.001	45	6	15	22500	9.4E-09	0.0E+00	0%	2100	1.1E-07	4.0E-03	2.7E-05	6%	
Vanadium	8.0E+00	1E-08	2300	1	0.001	45	6	15	22500	9.7E-09	0.0E+00	0%	2100	1.1E-07	1.4E-03	8.1E-05	17%	
Zinc	1.1E+01	1E-08	2300	1	0.001	45	6	15	22500	1.8E-08	0.0E+00	0%	2100	2.1E-07	6.0E-02	3.8E-05	1%	
Total											0.0E+00					4.7E-04		

SUBSURFACE SOIL INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 30)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL ADULT

Intake from ingestion of soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * CF * EF * ED * IR/BW * ATc \text{ or } ATnc * DY$$

$$\text{Risk} = \text{Intake} * CSF \text{ or } RfD$$

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	1E-06
CF = conversion for kg to mg	350
EF = adult exposure frequency (days/yr)	30
ED = adult exposure duration (yr)	100
IR = adult soil ingestion rate (mg/day)	70
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	30
DY = days per year (days/year)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RfD = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

COPC	Concentration Carcinogen (mg/kg)	Exposure Frequency (days/yr)	Exposure Duration (yr)	Conversion Factor (kg/mg)	Ingestion Rate (mg/day)	Body Weight (kg)	Averaging Carr Time (years)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk	Percent Carrnogeni	Averaging Noncarr Time (days)	Noncarr Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Aluminum	957.1	350	30	1E-06	100	70	25550	6.5E-03	0.0E+00	0%	10950	1.3E-02	1.0E+00	1.5E-02	55%	
Arsenic	0.9	350	30	1E-06	100	70	25550	5.2E-07	1.8E+00	9.1E-07	100%	10950	1.2E-06	3.0E-04	4.0E-03	21%
Chromium	10.1	350	30	1E-06	100	70	25550	5.9E-08	0.0E+00	0%	10950	1.4E-05	1.0E+00	1.4E-05	0%	
Cobalt	0.6	350	30	1E-06	100	70	25550	3.1E-07	0.0E+00	0.0E+00	0%	10950	7.3E-07	6.0E-02	1.2E-05	0%
Copper	3.5	350	30	1E-06	100	70	25550	2.1E-08	0.0E+00	0.0E+00	0%	10950	4.8E-08	3.7E-02	1.3E-04	1%
Manganese (soil)	6.8	350	30	1E-06	100	70	25550	4.0E-08	0.0E+00	0.0E+00	0%	10950	9.3E-06	1.4E-01	6.7E-05	0%
Mercury	0.1	350	30	1E-06	100	70	25550	7.6E-08	0.0E+00	0.0E+00	0%	10950	1.8E-07	3.0E-04	5.0E-04	3%
Nickel	2.2	350	30	1E-06	100	70	25550	1.3E-08	0.0E+00	0.0E+00	0%	10950	3.0E-08	2.0E-02	1.5E-04	1%
Vanadium	8.1	350	30	1E-06	100	70	25550	4.8E-08	0.0E+00	0.0E+00	0%	10950	1.1E-05	7.0E-03	1.0E-03	8%
TOTAL								9.1E-07						2.0E-02		

## SUBSURFACE SOIL DERMAL CONTACT EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 30)

REMEDIAL INVESTIGATION CTO-0231

MCB CAMP LEJEUNE, NORTH CAROLINA

FUTURE RESIDENTIAL ADULT

Dermal contact with soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RfD$$

Where:

C = contaminant concentration in soil (mg/kg)

CF = conversion factor (kg/mg)

SA = adult exposed skin surface area (cm<sup>2</sup>)AF = soil to skin adherence factor (mg/cm<sup>2</sup>)

Abs = fraction absorbed (unitless)

EF = adult exposure frequency (events/yr)

ED = adult exposure duration (years)

BW = adult body weight (kg)

ATc = averaging time for carcinogen (yr)

ATnc = averaging time for noncarcinogen (yr)

DY = day per year (day/yr)

CSF = cancer slope factor (mg/kg-day)-1

RfD = reference dose (mg/kg-day)

## INPUTS

1E-06

5800

1

Specific

350

30

70

70

30

365

specific

specific

Note: Inputs are scenario and site specific

COPC	Concentration Carcinogen (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> ) Adult	Adherence Factor (mg/cm <sup>2</sup> )	Fraction Absorbed (%)	Exposure Frequency (events/yr) Adult	Exposure Duration (yrs) Adult	Body Weight (kg) Adult	Averaging Carc Time (days)	Carc Dose (mg/kg/day) Adult	Dermal Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Averaging Noncarc Time (days)	Noncarc Dose (mg/kg/day) Adult	Dermal Adjust Reference Dose (mg/kg-day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	4557.1	1E-06	5800	1	0.001	350	30	70	25500	3.3E-04	0.0E+00	0%	10950	7.6E-04	2.0E-01	3.8E-03	68%	
Arsenic	0.9	1E-06	5800	1	0.001	350	30	70	25500	3.0E-06	0.0E+00	2.6E-07	100%	10950	7.0E-08	6.0E-05	1.2E-03	21%
Chromium	10.1	1E-06	5800	1	0.001	350	30	70	25500	3.4E-07	0.0E+00	0.0E+00	0%	10950	8.0E-07	2.0E-01	4.0E-06	0%
Cobalt	0.5	1E-06	5800	1	0.001	350	30	70	25500	1.8E-06	0.0E+00	0.0E+00	0%	10950	4.3E-08	1.2E-02	3.5E-06	0%
Copper	3.5	1E-06	5800	1	0.001	350	30	70	25500	1.2E-07	0.0E+00	0.0E+00	0%	10950	2.9E-07	7.4E-03	3.8E-05	1%
Manganese (soil)	6.8	1E-06	5800	1	0.001	350	30	70	25500	2.3E-07	0.0E+00	0.0E+00	0%	10950	6.4E-07	2.8E-02	1.9E-05	0%
Mercury	0.1	1E-06	5800	1	0.001	350	30	70	25500	4.4E-09	0.0E+00	0.0E+00	0%	10950	1.0E-08	6.0E-05	1.7E-04	3%
Nickel	2.2	1E-06	5800	1	0.001	350	30	70	25500	7.4E-08	0.0E+00	0.0E+00	0%	10950	1.7E-07	4.0E-03	4.3E-05	1%
Vanadium	8.1	1E-06	5800	1	0.001	350	30	70	25500	2.8E-07	0.0E+00	0.0E+00	0%	10950	6.5E-07	1.4E-03	4.6E-04	8%
TOTAL											2.6E-07						5.7E-03	

SUBSURFACE SOIL PARTICULATE INHALATION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 30)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE RESIDENTIAL ADULT

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (\text{C} * \text{EF} * \text{ED} * \text{IR} * 1/\text{PEF}) / (\text{BW} * \text{ATc or ATnc} * \text{DY})$$

$$\text{Risk} = \text{Intake} * \text{CSF or /RD}$$

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	Specific
CSF = carcinogenic slope factor	Specific
RID = reference dose for noncarcinogen	Specific
IR = Inhalation rate (m³)	20
EF = adult exposure frequency (days)	360
ED = adult exposure duration (years)	30
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	30
DY = day per year (day/yr)	365
PEF = particulate emission factor (m³/kg)	4.63E+09

Note: Inputs are scenario and site specific

COMP	Concentration Carcinogen (mg/kg)	Particulate Emission Factor (m³/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m³/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Cerc Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncerc Time (days)	Noncerc Dose (mg/kg/day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Aluminum	9.8E+03	4.6E+09	350	20	30	70	25550	2.4E-07	0.0E+00	0%	10950	5.7E-07	0.0E+00	0.0E+00	0%	
Asenic	8.9E-01	4.6E+09	350	20	30	70	25550	2.2E-11	1.5E+01	3.4E-10	100%	10950	5.2E-11	0.0E+00	0.0E+00	0%
Chromium	1.0E+01	4.6E+09	350	20	30	70	25550	2.0E-10	0.0E+00	0.0E+00	0%	10950	6.0E-10	0.0E+00	0.0E+00	0%
Cobalt	6.4E-01	4.6E+09	350	20	30	70	25550	1.4E-11	0.0E+00	0.0E+00	0%	10950	3.2E-11	0.0E+00	0.0E+00	0%
Copper	3.6E+00	4.6E+09	350	20	30	70	25550	8.9E-11	0.0E+00	0.0E+00	0%	10950	2.1E-10	0.0E+00	0.0E+00	0%
Manganese (soil)	6.8E+00	4.6E+09	350	20	30	70	25550	1.7E-10	0.0E+00	0.0E+00	0%	10950	4.0E-10	0.0E+00	0.0E+00	0%
Mercury	1.3E-01	4.6E+09	350	20	30	70	25550	3.3E-12	0.0E+00	0.0E+00	0%	10950	7.6E-12	8.0E-05	8.9E-08	100%
Nickel	2.2E+00	4.6E+09	350	20	30	70	25550	5.6E-11	0.0E+00	0.0E+00	0%	10950	1.3E-10	0.0E+00	0.0E+00	0%
Vanadium	8.1E+00	4.6E+09	350	20	30	70	25550	2.1E-10	0.0E+00	0.0E+00	0%	10950	4.8E-10	0.0E+00	0.0E+00	0%
<b>TOTAL</b>									3.4E-10						8.9E-08	

SURFACE WATER INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO.7 (SITE 30 - FRENCHS CREEK)  
 REMEDIAL INVESTIGATION CTD-0021  
 MCAS CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT AND FUTURE ADULT RESIDENT

The intake from the ingestion of surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = Cr * CR * ET * EF * ED/BW * ATc \text{ or } ATnc * DY$$

$$\text{Risk} = \text{Intake} * CSF \text{ or } RD$$

CHANGE!!

Where:

	INPUT
Cr = contaminant concentration in surface water (mg/l)	0.005
CR = ingestion rate (Liter/hour)	2.6
ET = exposure time (hours/interval)	45
EF = exposure frequency (events/yr)	30
ED = exposure duration (yrs)	70
BW = body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	30
DY = days per year (days)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RD = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

FRENCHS CREEK

POPC	Concentration Carcinogen (mg/l)	Contact Rate (hour)	Exposure Time (hour/interval)	Exposure Frequency (events/yr)	Exposure Duration (years)	Body Weight (kg)	AVERAGING Carr. Time (day)	Carc. Dose (mg/kg-day)	Cancer Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Averaging time Noncarc. (day)	NONCARC. Dose (mg/kg-day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Lead	1.0E+00	0.005	2.6	45	30	70	25500	1.0E-05	0.0E+00	0%	0%	30	1.0E-02	1.0E+00	1.0E-02	35%
Manganese (water)	2.3E-03	0.005	2.6	45	30	70	25500	2.3E-08	0.0E+00	0%	0%	30	1.0E-05	0.0E+00	0.0E+00	0%
Mercury	1.1E-02	0.005	2.6	45	30	70	25500	1.1E-07	0.0E+00	0%	0%	30	9.3E-05	6.0E-03	1.0E-02	63%
UTAC									0.0E+00						0.0E+00	

SURFACE WATER DERMAL CONTACT EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 SITE SO - FRENCHIS CREEK  
 REMEDIAL INVESTIGATION - CTD-021  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 CURRENT AND FUTURE ADULT RESIDENT

The intake from dermal contact with surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = \text{Cw} \times \text{SA} \times \text{PC} \times \text{ET} \times \text{EF} \times \text{ED} \times \text{CF/BW} \times \text{ATc or ATnc} \times \text{DY}$$

$$\text{Risk} = \text{Intake} \times \text{CSF or RD}$$

CHANGE1

Where:

	INPUTS
CW = contaminant concentration in water (mg/l)	
SA = skin surface available for contact (cm <sup>2</sup> )	5600
PC = contaminant specific dermal permeability (cm/hr)	Specific
ET = exposure time (hours/day)	2.6
EF = exposure frequency (days/yr)	45
ED = exposure duration (years)	30
CF = volumetric conversion factor for water (liter/1000 cm <sup>3</sup> )	0.001
BW = body weight (kg)	70
ATc = averaging time for carcinogen (y)	70
ATnc = averaging time for noncarcinogen (y)	30
DY = days per year (days)	365
CSF = cancer slope factor (mg/kg-day)-1	Specific
RD = reference dose (mg/kg-day)	Specific

Note: Inputs are site and scenario specific

FRENCHS CREEK

Contaminant	Concentration (mg/l)	Surface Area (cm <sup>2</sup> )	Dermal Permeability (cm/hr)	Exposure Time (hours/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	Volumetric Conversion (L/m <sup>3</sup> )	Body Weight (kg)	Averaging Cred Time (days)	Carc. Dose (mg/kg-day)	Dermal Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Average Noncarc Time (days)	Noncarc. Dose (mg/kg-day)	Dermal Adjust Reference Dose (mg/kg-day)	NONCARC Risk	Percent Noncarcinogenic Risk
Nominal	1.3E+00	5600	1.0E-03	2.6	45	30	0.001	70	25500	1.7E-05	0.0E+00	0%	10800	3.0E-05	2.0E-01	2.0E-01	35%	
Lead	2.3E-03	5600	4.0E-05	2.6	45	30	0.001	70	25500	1.0E-10	0.0E+00	0%	10800	2.4E-10	0.0E+00	0.0E+00	0%	
Manganese (water)	1.1E-02	5600	1.0E-03	2.6	45	30	0.001	70	25500	1.3E-07	0.0E+00	0%	10800	2.0E-07	1.0E-03	2.0E-04	62%	
Mercury	1.5E-04	5600	1.0E-03	2.6	45	30	0.001	70	25500	1.7E-09	0.0E+00	0%	10800	4.0E-09	0.0E+00	0.0E+00	12%	
TOTAL										0.0E+00						5.0E-04		

SEDIMENT INGESTION EXPOSURE / SEDIMENT  
OPERABLE UNIT NO. 7 (SITE 30 - FRENCHS CREEK)  
ENVIRONMENTAL INVESTIGATION CTO-0221  
KCB CAMP LEJEUNE, NORTH CAROLINA  
CURRENT AND FUTURE RESIDENTIAL ADULT

intake from ingestion of sediment is calculated as follows:

$$\text{intake (mg/kg-day)} = C * IR * CF * EF * ED / BW * ATc \text{ or } ATnc * DY$$

CHANGE

Risk = Intake \* CSF or RD

Where:

C = contaminant concentration in sediment (mg/kg)

INPUTS

CF = conversion factor kg to mg

1E+00

EF = exposure frequency (days/yr)

45

ED = exposure duration (y)

30

IR = soil ingestion rate (mg/day)

100

BW = body weight (kg)

70

ATc = averaging time for carcinogen (y)

70

ATnc = averaging time for noncarcinogen (y)

30

DY = days per year (days/year)

365

CSF = cancer slope factor (mg/kg-day)-1

Specific

RD = reference dose (mg/kg-day)

Specific

Note: Inputs are scenario and site specific

FRENCHS CREEK

OPC	Concentration (mg/kg)	Exposure Frequency (days/yr)	Exposure Duration (y)	Ingestion Rate (mg/day)	Conversion Factor (kg/mg)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Uranium	3.4E+03	45	30	100	1E-03	70	25500	2.0E-04	0.0E+00	0%	10000	8.0E-04	1.0E+00	8.0E-04	0%	
Zirconium	7.0E+00	45	30	100	1E-03	70	25500	8.0E-07	0.0E+00	0%	10000	1.4E-03	1.0E+00	1.4E-03	0%	
Copper	9.0E+00	45	30	100	1E-03	70	25500	7.0E-07	0.0E+00	0%	10000	1.7E-03	3.7E-02	4.6E-03	5%	
Lead	1.0E+01	45	30	100	1E-03	70	25500	1.2E-06	0.0E+00	0%	10000	2.7E-06	0.0E+00	0.0E+00	0%	
Asbestos (total)	1.0E+01	45	30	100	1E-03	70	25500	1.4E-06	0.0E+00	0%	10000	3.3E-06	1.4E-01	2.4E-05	3%	
As-3+	6.0E+00	45	30	100	1E-03	70	25500	4.0E-07	0.0E+00	0%	10000	1.0E-03	2.0E-02	8.1E-03	0%	
Amidium	6.0E+00	45	30	100	1E-03	70	25500	4.0E-07	0.0E+00	0%	10000	1.1E-03	7.0E-03	1.6E-04	17%	
Zinc	1.1E+01	45	30	100	1E-03	70	25500	8.0E-07	0.0E+00	0%	10000	2.0E-03	3.0E-01	8.0E-03	1%	
UTAC									0.0E+00					8.0E-04		

SEDIMENT DERMAL CONTACT EXPOSURE ASSESSMENT  
OPERABLE UNIT NO.7 (SITE 30 - FRENCHS CREEK)  
REMEDIAl INVESTIGATION CTO-0201  
MCB CAMP LEJEUNE, NORTH CAROLINA  
CURRENT AND FUTURE RESIDENTIAL ADULT

The intake from dermal contact to sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RD$$

Where:		INPUTS
C = contaminant concentration in soil (mg/kg)		1.00E-08
CF = conversion factor (kg/mg)		5800
SA = exposed skin surface area (cm <sup>2</sup> )		1
AF = sediment to skin advance factor (mg/cm <sup>2</sup> )		48
Abs = fraction absorbed (unitless) (contaminant specific)	Specific	0.01
EF = exposure frequency (events/yr)		30
ED = exposure duration (years)		70
BW = body weight (kg)		70
ATc = averaging time for carcinogen (yr)		30
ATnc = averaging time for noncarcinogen (yr)		30
DY = day per year (day/yr)		365
CSF = cancer slope factor (mg/kg-day)-1	Specific	3.0E-05
RD = reference dose (mg/kg-day)	Specific	0.0001

Note: Inputs are scenario and site specific

FRENCHS CREEK

Contaminant	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> )	Advance Factor (mg/cm <sup>2</sup> )	Abs Factor (%)	Exposure Frequency (events/yr)	Exposure Duration (yr)	Body Weight (kg)	Average Carc Time (days)	Carc. Dose (mg/kg/day)	Dermal Assess Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Potent Carcinogenic Risk	Average Noncarc Time (days)	Noncarc. Dose (mg/kg/day)	Dermal Assess Reference Dose (mg/kg/day)	Noncarcinogenic Risk	Potent Noncarcinogenic Risk
Nickel	3.4E+03	1E-08	5800	1	0.001	48	30	70	25000	1.0E-08	0.0E+00	0%	10000	3.0E-05	2.0E-01	1.7E-04	0%	
Chromium	7.0E+00	1E-08	5800	1	0.001	48	30	70	25000	3.0E-08	0.0E+00	0%	10000	8.0E-08	2.0E-01	4.0E-07	0%	
Copper	8.0E+00	1E-08	5800	1	0.001	48	30	70	25000	4.0E-08	0.0E+00	0%	10000	8.7E-08	7.4E-03	1.3E-06	0%	
Lead	1.0E+01	1E-08	5800	1	0.001	48	30	70	25000	6.0E-08	0.0E+00	0%	10000	1.0E-07	0.0E+00	0.0E+00	0%	
Manganese (soil)	1.0E+01	1E-08	5800	1	0.001	48	30	70	25000	8.0E-08	0.0E+00	0%	10000	1.0E-07	2.0E-02	8.0E-08	0%	
Nickel	8.0E+00	1E-08	5800	1	0.001	48	30	70	25000	2.0E-08	0.0E+00	0%	10000	8.0E-08	4.0E-03	1.0E-05	0%	
Vanadium	8.0E+00	1E-08	5800	1	0.001	48	30	70	25000	2.0E-08	0.0E+00	0%	10000	8.1E-08	1.0E-03	4.4E-03	17%	
Zinc	1.1E+01	1E-08	5800	1	0.001	48	30	70	25000	4.0E-08	0.0E+00	0%	10000	1.0E-07	0.0E+00	1.0E-05	0%	
<b>TOTAL</b>																		

## SURFACE WATER INGESTION EXPOSURE ASSESSMENT

OPERABLE UNIT NO.7 (SITE 30)

REMEDIAL INVESTIGATION CTD-0201

MCB CAMP LEUELNE, NORTH CAROLINA

CURRENT MILITARY PERSONNEL

The intake from the ingestion of surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = C_w \times CR \times ET \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RD$$

Where:

 $C_w$  = contaminant concentration in surface water (mg/l)

CR = contact rate (liters/hour)

## INPUT

ET = exposure time (hours/year)

0.005

EF = exposure frequency (events/yr)

26

ED = exposure duration (hrs)

45

BW = body weight (kg)

4

ATc = averaging time for carcinogen (yr)

70

ATnc = averaging time for noncarcinogen (yr)

70

DY = days per year (days)

4

CSF = cancer slope factor (mg/kg-day)-1

365

RD = reference dose (mg/kg-day)

specific

specific

Note: Inputs are scenario and site specific

OPC	Concentration Carcinogen (mg/l)	Contact Rate (hour)	Exposure Time (hr/year)	Exposure Frequency (events/yr)	Exposure Duration (years)	Body Weight (kg)	Averaging Time Car. (days)	Carc. Dose (mg/kg-day)	Cancer Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Noncarc. Risk	Averaging Time Noncarc. (years)	Noncarc. Dose (mg/kg-day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Potassium	1.48	0.005	26	45	4	70	25000	1.9E-05	0.0E+00	0.0E+00	0%	1460	3.4E-05	1.0E+00	3.4E-05	35%
Lead	0.0023	0.005	26	45	4	70	25000	3.0E-09	0.0E+00	0.0E+00	0%	1460	6.3E-08	0.0E+00	6.3E-08	0%
Manganese (water)	0.0111	0.005	26	45	4	70	25000	1.6E-08	0.0E+00	0.0E+00	0%	1460	2.3E-07	6.0E-03	5.1E-05	53%
Mercury	0.00015	0.005	26	45	4	70	25000	2.0E-10	0.0E+00	0.0E+00	0%	1460	3.4E-09	3.0E-04	1.1E-05	12%
TOTAL									0.0E+00						0.0E+00	

## SURFACE WATER DERMAL CON... EXPOSURE ASSESSMENT

OPERABLE UNIT NO.7 (SITE 30)

RISK EVALUATION - CTO-0231

MCB CAMP LEJEUNE, NORTH CAROLINA

CURRENT MILITARY PERSONNEL

The intake from dermal contact with surface water is calculated as follows:

$$\text{Intake (mg/kg-day)} = Cw \times SA \times PC \times ET \times EF \times ED \times CF/BW \times ATc \text{ or } ATn \times DT$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RD$$

Where:

Cw = contaminant concentration in water (mg/l)

SA = skin surface available for contact (cm<sup>2</sup>)

## INPUTS

5800

Specific

PC = contaminant specific dermal permeability (cm/hr)

2.6

ET = exposure time (hours/day)

4

EF = exposure frequency (day/yr)

45

ED = exposure duration (years)

4

CF = volumetric conversion factor for water (liter/1000 cm<sup>3</sup>)

0.001

BW = body weight (kg)

70

ATc = averaging time for carcinogen (yr)

70

ATn = averaging time for noncarcinogen (yr)

4

DT = days per year (days)

365

CSF = cancer slope factor (mg/kg-day)-1

Specific

RD = reference dose (mg/kg-day)

Specific

Note: Inputs are site and scenario specific

POPs	Concentration Carcinogen (mg/l)	Surface Area (cm <sup>2</sup> )	Dermal Permeability (cm/hr)	Exposure Time (hours/day)	Exposure Frequency (days/yr)	Exposure Duration (years)	volumetric Conversion (L-mg)	Body Weight (kg)	Averaging Carc Time (days)	Carc Dose (mg/kg-day)	Dermal Adjust Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg-day)	Dermal Adjust Reference Dose (mg/kg-day)	Noncarc Risk	Percent Noncarcinogenic Risk
Arsenic	1.40	5800	1.0E-03	2.6	45	4	0.001	70	26250	3.48E-06	0.0E+00	0%	1400	0.00E-05	5.1E+00	1.2E-05	35%	
Lead	0.0003	5800	1.0E-03	2.6	45	4	0.001	70	26250	6.41E-06	0.0E+00	0%	1400	9.47E-06	0.0E+00	0.0E+00	0%	
Manganese (water)	0.0111	5800	1.0E-03	2.6	45	4	0.001	70	26250	2.01E-06	0.0E+00	0%	1400	4.37E-07	2.6E-02	1.8E-05	63%	
Mercury	0.00015	5800	1.0E-03	2.6	45	4	0.001	70	26250	3.83E-10	0.0E+00	0%	1400	8.18E-09	1.8E-03	4.1E-08	12%	
TOTAL										0.0E+00						3.5E-05		

SEDIMENT INGESTION EXPOSURE ASSESSMENT  
OPERABLE UNIT NO. 7 (SITE 33)  
REMEDIAL INVESTIGATION CTO-0231  
MCB CAMP LEJEUNE, NORTH CAROLINA  
CURRENT MILITARY PERSONNEL

Intake from ingestion of sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * R * CF * EF * ED / BW * ATC \text{ or } ATnc * DY$$

Risk = Intake \* CSF or RfD

Where:

C = contaminant concentration in sediment (mg/kg)

CF = conversion for kg to mg

INPUTS

EF = exposure frequency (days/yr)

1E-08

ED = exposure duration (y)

360

R = soil ingestion rate (mg/day)

4

BW = body weight (kg)

100

ATC = averaging time for carcinogen (yr)

70

ATnc = averaging time for noncarcinogen (yr)

4

DY = days per year (days/year)

365

CSF = cancer slope factor (mg/kg-day)-1

Specific

RfD = reference dose (mg/kg-day)

Specific

Note: Inputs are scenario and site specific

COPC	Concentration (mg/kg)	Exposure Frequency (days/yr)	Exposure Duration (y)	Ingestion Rate (mg/day)	Conversion Factor (kg/mg)	Body Weight (kg)	Averaging Carr. Time (days)	Carc. Dose (mg/kg/day)	Slope Factor (mg/kg/day)-1	Carcinogenic Risk	Percent Carcinogenic Risk	Averaging Noncarc. Time (days)	Noncarc. Dose (mg/kg/day)	Reference Dose (mg/kg/day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Nickel	3.4E+03	360	4	100	1E-08	70	26560	2.7E-04	0.0E+00	0%	1400	4.5E-03	1.0E+00	4.5E-03	0%	
Chromium	7.9E+00	360	4	100	1E-08	70	26560	6.2E-07	0.0E+00	0%	1400	1.1E-05	1.0E+00	1.1E-05	0%	
Copper	9.5E+00	360	4	100	1E-08	70	26560	7.4E-07	0.0E+00	0%	1400	1.3E-05	3.7E-02	3.5E-04	5%	
Lead	1.0E+01	360	4	100	1E-08	70	26560	1.2E-06	0.0E+00	0%	1400	2.1E-05	0.0E+00	0.0E+00	0%	
Manganese (soil)	1.5E+01	360	4	100	1E-08	70	26560	1.8E-06	0.0E+00	0%	1400	2.6E-05	1.4E-01	1.5E-04	3%	
Nickel	5.5E+00	360	4	100	1E-08	70	26560	4.6E-07	0.0E+00	0%	1400	7.9E-05	2.0E-02	4.0E-04	6%	
Zinc	6.0E+00	360	4	100	1E-08	70	26560	4.7E-07	0.0E+00	0%	1400	8.2E-05	7.0E-03	1.2E-03	17%	
Zinc	1.1E+01	360	4	100	1E-08	70	26560	8.8E-07	0.0E+00	0%	1400	1.6E-05	3.0E-01	5.2E-05	1%	
TOTAL									0.0E+00					5.6E-03		

## SEDIMENT DERMAL CONTACT RISK ASSESSMENT

OPERABLE UNIT NO.1 (SITE 30)

REMEDIAL INVESTIGATION CTD-0201

MCB CAMP LEJEUNE, NORTH CAROLINA

CURRENT MILITARY PERSONNEL

The intake from dermal contact to sediment is calculated as follows:

$$\text{Intake (mg/kg-day)} = C \times CF \times SA \times AF \times Abs \times EF \times ED/BW \times ATc \text{ or } ATnc \times DY$$

$$\text{Risk} = \text{Intake} \times CSF \text{ or } RD$$

Where:

C = contaminant concentration in soil (mg/kg)

## INPUTS

1.0E-08

CF = conversion factor (kg/mg)

5800

SA = exposed skin surface area (cm<sup>2</sup>)

1

AF = sediment to skin adherence factor (mg/m<sup>2</sup>)

Specific

Abs = fraction absorbed (urine/urine) (contaminant specific)

350

EF = exposure frequency (events/yr)

4

ED = exposure duration (years)

70

BW = body weight (kg)

70

ATc = averaging time for carcinogen (yr)

4

ATnc = averaging time for noncarcinogen (yr)

365

DY = day per year (day/yr)

Specific

CSF = cancer slope factor (mg/kg-day)-1

Specific

RD = reference dose (mg/kg-day)

Specific

SUBSURFACE SOIL INGESTION EXPOSURE ASSESSMENT  
 OPERABLE UNIT NO. 7 (SITE 30)  
 REMEDIAL INVESTIGATION CTO-0231  
 MCB CAMP LEJEUNE, NORTH CAROLINA  
 FUTURE CONSTRUCTION WORKER

Intake from ingestion of soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * CF * EF * ED * IR/BW * ATc \text{ or } ATnc * DY$$

$$\text{Risk} = \text{Intake} * CSF \text{ or } RID$$

Where:

	INPUTS
C = contaminant concentration in soil (mg/kg)	
CF = conversion for kg to mg	1E-06
EF = adult exposure frequency (days/yr)	90
ED = adult exposure duration (yr)	1
IR = adult soil ingestion rate (mg/day)	480
BW = adult body weight (kg)	70
ATc = averaging time for carcinogen (yr)	70
ATnc = averaging time for noncarcinogen (yr)	1
DY = days per year (days/year)	365
CSF = cancer slope factor (mg/kg-day)-1	specific
RID = reference dose (mg/kg-day)	specific

Note: Inputs are scenario and site specific

COPC	Concentration Carcinogen (mg/kg)	Exposure Frequency (days/yr) Adult	Exposure Duration (yr) Adult	Conversion Factor (kg/mg)	Ingestion Rate (mg/day) Adult	Body Weight (kg) Adult	Averaging Carc Time (days)	Carc Dose (mg/kg/day) Adult	Slope Factor (mg/kg/day)-1	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Averaging Noncarc. Time (days)	Noncarc. Dose (mg/kg/day) Adult	Reference Dose (mg/kg/day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult	
Aluminum	9557.1	90	1	1E-06	480	70	25550	2.3E-04	0.0E+00	0%	365	1.6E-02	1.0E+00	1.6E-02	66%		
Arsenic	0.9	90	1	1E-06	480	70	25550	2.1E-08	1.8E+00	3.7E-08	100%	365	1.5E-08	3.0E-04	8.0E-03	21%	
Chromium	10.1	90	1	1E-06	480	70	25550	2.4E-07	0.0E+00	0%	365	1.7E-05	1.0E+00	1.7E-05	0%		
Cobalt	0.5	90	1	1E-06	480	70	25550	1.3E-06	0.0E+00	0.0E+00	0%	365	9.1E-07	6.0E-02	1.6E-05	0%	
Copper	3.5	90	1	1E-06	480	70	25550	8.5E-08	0.0E+00	0.0E+00	0%	365	6.0E-08	3.7E-02	1.6E-04	1%	
Manganese (soil)	6.8	90	1	1E-06	480	70	25550	1.6E-07	0.0E+00	0.0E+00	0%	365	1.1E-05	1.4E-01	8.2E-05	0%	
Mercury	0.1	90	1	1E-06	480	70	25550	3.1E-09	0.0E+00	0.0E+00	0%	365	2.2E-07	3.0E-04	7.3E-04	3%	
Nickel	2.2	90	1	1E-06	480	70	25550	5.3E-08	0.0E+00	0.0E+00	0%	365	3.7E-08	2.0E-02	1.6E-04	1%	
Vanadium	8.1	90	1	1E-06	480	70	25550	2.0E-07	0.0E+00	0.0E+00	0%	365	1.4E-05	7.0E-03	2.0E-03	8%	
TOTAL									3.7E-08						2.4E-02		

## SUBSURFACE SOIL DERMAL CONTACT EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 30)

REMEDIAl INVESTIGATION CTO-0231

MCB CAMP LEJEUNE, NORTH CAROLINA

FUTURE CONSTRUCTION WORKER

Dermal contact with soil is calculated as follows:

$$\text{Intake (mg/kg-day)} = C * CF * SA * AF * Abs * EF * ED/BW * ATc \text{ or } ATnc * DY$$

$$\text{Risk} = \text{Intake} * CSF \text{ or } RID$$

Where:

C = contaminant concentration in soil (mg/kg)

## INPUTS

CF = conversion factor (kg/mg)

1E+06

SA = adult exposed skin surface area (cm<sup>2</sup>)

4300

AF = soil to skin adherence factor (mg/cm<sup>2</sup>)

1

Abs = fraction absorbed (unitless)

Specific

EF = adult exposure frequency (events/yr)

90

ED = adult exposure duration (years)

1

BW = adult body weight (kg)

70

ATc = averaging time for carcinogen (yr)

70

ATnc = averaging time for noncarcinogen (yr)

1

DY = day per year (day/yr)

365

CSF = cancer slope factor (mg/kg-day)<sup>-1</sup>

specific

RID = reference dose (mg/kg-day)

specific

Note: Inputs are scenario and site specific

PCP	Concentration (mg/kg)	Conversion Factor (kg/mg)	Surface Area (cm <sup>2</sup> )	Adherence Factor (mg/cm <sup>2</sup> )	Fraction Absorbed (%)	Exposure Frequency (events/yr) Adult	Exposure Duration (yrs) Adult	Body Weight (kg) Adult	Averaging Carr. Time (days)	Carc. Dose (mg/kg/day) Adult	Dermal Adjust Slope Factor (mg/kg-day) <sup>-1</sup>	Carcinogenic Risk Adult	Percent Carcinogenic Risk Adult	Averaging Noncarc. Time (days)	Noncarc. Dose (mg/kg/day) Adult	Dermal Adjust Reference Dose (mg/kg-day)	Noncarcinogenic Risk Adult	Percent Noncarcinogenic Risk Adult
Aluminum	9.6E+03	1E+06	4300	1	0.001	90	1	70	25550	2.1E-06	0.0E+00	0%	365	1.4E-04	2.0E-01	7.2E-04	66%	
Arsenic	8.9E-01	1E+06	4300	1	0.001	90	1	70	25550	1.9E-10	8.8E+00	1.7E-09	100%	365	1.3E-06	6.0E-05	2.2E-04	21%
Chromium	1.0E+01	1E+06	4300	1	0.001	90	1	70	25550	2.2E-09	0.0E+00	0.0E+00	0%	365	1.5E-07	2.0E-01	7.6E-07	0%
Cobalt	6.4E-01	1E+06	4300	1	0.001	90	1	70	25550	1.2E-10	0.0E+00	0.0E+00	0%	365	8.1E-09	1.2E-02	6.8E-07	0%
Copper	3.6E+00	1E+06	4300	1	0.001	90	1	70	25550	7.6E-10	0.0E+00	0.0E+00	0%	365	5.3E-08	7.4E-03	7.2E-08	1%
Manganese (soil)	6.8E+00	1E+06	4300	1	0.001	90	1	70	25550	1.6E-09	0.0E+00	0.0E+00	0%	365	1.0E-07	2.8E-02	3.7E-08	0%
Mercury	1.3E-01	1E+06	4300	1	0.001	90	1	70	25550	2.8E-11	0.0E+00	0.0E+00	0%	365	2.0E-09	6.0E-05	3.3E-05	3%
Nickel	2.2E+00	1E+06	4300	1	0.001	90	1	70	25550	4.7E-10	0.0E+00	0.0E+00	0%	365	3.3E-08	4.0E-03	8.2E-06	1%
Vanadium	8.1E+00	1E+06	4300	1	0.001	90	1	70	25550	1.8E-09	0.0E+00	0.0E+00	0%	365	1.2E-07	1.4E-03	8.8E-05	8%
TOTAL										1.7E-05							1.7E-03	

## SUBSURFACE SOIL PAH...DUST INHALATION EXPOSURE ASSESSMENT

OPERABLE UNIT NO. 7 (SITE 30)

REMEDIAl INVESTIGATION CTO-0231

MCB CAMP LEJEUNE, NORTH CAROLINA

FUTURE CONSTRUCTION WORKER

Intake from the inhalation of particulates is calculated as follows:

$$\text{Intake (mg/kg-day)} = (C * EF * ED * IR * 1/PEF) / (BW * ATc or ATnc * DY)$$

$$\text{Risk} = \text{Intake} * \text{CSF or } / \text{RfD}$$

Where:

C = contaminant concentration in soil (mg/kg)

CSF = carcinogenic slope factor

RfD = reference dose for noncarcinogen

IR = inhalation rate (m³)

## INPUTS

Specific

Specific

Specific

20

EF = adult exposure frequency (days)

90

ED = adult exposure duration (years)

1

BW = adult body weight (kg)

70

ATc = averaging time for carcinogen (yr)

70

ATnc = averaging time for noncarcinogen (yr)

1

DY = day per year (day/yr)

365

PEF = particulate emission factor (m³/kg)

4.63E+09

Note: Inputs are scenario and site specific

COPC	Concentration Carcinogen (mg/kg)	Particulate Emission Factor (m³/kg)	Exposure Frequency (events/yr)	Inhalation Rate (m³/day)	Exposure Duration (yrs)	Body Weight (kg)	Average Carc Time (days)	Carc Dose (mg/kg/day)	Slope Factor (mg/kg-day)-1	Carcinogenic Risk	Percent Contribution to Risk	Average Noncarc Time (days)	Noncarc Dose (mg/kg/day)	Reference Dose (mg/kg-day)	Noncarcinogenic Risk	Percent Noncarcinogenic Risk
Methinkium	9.8E+03	4.6E+09	90	20	1	70	25550	2.1E-09	0.0E+00	0%	365	1.5E-07	0.0E+00	0.0E+00	0%	
Arsenic	8.8E-01	4.6E+09	90	20	1	70	25550	1.9E-13	1.5E+01	2.9E-12	100%	365	1.3E-11	0.0E+00	0.0E+00	0%
Chromium	1.0E+01	4.6E+09	90	20	1	70	25550	2.2E-12	0.0E+00	0.0E+00	0%	365	1.5E-10	0.0E+00	0.0E+00	0%
Cobalt	5.4E-01	4.6E+09	90	20	1	70	25550	1.2E-13	0.0E+00	0.0E+00	0%	365	8.2E-12	0.0E+00	0.0E+00	0%
Copper	3.6E+00	4.6E+09	90	20	1	70	25550	7.7E-13	0.0E+00	0.0E+00	0%	365	5.4E-11	0.0E+00	0.0E+00	0%
Manganese (soil)	8.8E+00	4.6E+09	90	20	1	70	25550	1.6E-12	0.0E+00	0.0E+00	0%	365	1.0E-10	0.0E+00	0.0E+00	0%
Mercury	1.3E-01	4.6E+09	90	20	1	70	25550	2.8E-14	0.0E+00	0.0E+00	0%	365	2.0E-12	8.6E-05	2.3E-08	100%
Nickel	2.2E+00	4.6E+09	90	20	1	70	25550	4.7E-13	0.0E+00	0.0E+00	0%	365	3.3E-11	0.0E+00	0.0E+00	0%
Vanadium	8.1E+00	4.6E+09	90	20	1	70	25550	1.8E-12	0.0E+00	0.0E+00	0%	365	1.2E-10	0.0E+00	0.0E+00	0%
TOTAL																

**APPENDIX S**  
**SUMMARY OF GROUNDWATER DATA AND AQUIFER**  
**CHARACTERISTICS AT MCB CAMP LEJEUNE**

---

## TECHNICAL MEMORANDUM

### SUMMARY OF GROUNDWATER DATA AND AQUIFER CHARACTERISTICS MARINE CORPS BASE, CAMP LEJEUNE JACKSONVILLE, NORTH CAROLINA

#### SUMMARY

This study examines the utility of exploratory aquifer tests (pump tests) at investigation sites across Marine Corps Base, Camp Lejeune (MCB-CL). The study reviews the available information on the relevant water-bearing layers, considers the general characteristics and applicability of aquifer tests, and concludes:

- That available information is satisfactorily complete to allow appropriate designs of groundwater systems in the main operating areas of MCB-CL;
- That quantified characterization of the water-bearing layers in explored areas of MCB-CL can be extended to other areas having similar geologic terrain;
- That exploratory tests are not longer routinely required or advisable;
- That reconnaissance testing (well-head tests or slug tests) of each newly installed or otherwise uncharacterized data station is highly advisable; and,
- That performance testing of groundwater extraction systems should be the recommended form of evaluating and adjusting withdrawal systems.

#### BACKGROUND

This study considers the aquifer characteristics (especially, the Coefficient of Transmissivity) and the production capacities (available discharge rates) of the two water-bearing layers relevant to the studies at MCB-CL. These water-bearing layers are the (shallow or surficial) water table and the Upper Castle Hayne Aquifer.

The water table at MCB-CL occupies the water-bearing zone within 25 to 35 feet of the surface; the Castle Hayne, immediately below this. However, the separation of the water table and the Castle Hayne is not always obvious. Usually, this separation is effected only by the low permeability material of the water table transiting to the significantly more permeable material of the Upper Castle Hayne; there is rarely an aquiclude or aquitard of vertically extensive clay separating the water table from the Castle Hayne.

The data available for this summary derive from three main sources:

- Assessment of Hydrologic and Hydrogeologic Data at Camp Lejeune Marine Corps Base, North Carolina; U. S. Geological Survey, Water-Resources Investigation Report 89-4096; 1989

## **COMPARABILITY OF DATA ACROSS MCB-CL**

The stratigraphic sequences of MCB-CL containing the water table and the Upper Castle Hayne have been well characterized. The available information indicates that the lithology and the hydrologic conditions can be correlated stratigraphically across the base (Tables 1 and 2). From these correlations, aquifer performance can be predicted sufficiently for an engineering design whose final criteria for suitability are performance-based.

The upper water-bearing zone is a highly variable layering and intercalation of clay, silt, and sand. This variability, however, is found within recognizable limits. These limits correspond to the range of hydrologic characteristics described previously. Similar correlation is available for the lithology and hydrology of the Upper Castle Hayne.

In areas not near stations catalogued in Tables 1, 2, and 3, a reconnaissance comparison of well-head tests (slug tests) and an examination of lithologic descriptions will likely be sufficient to support the engineering evaluation of the site. There is ample demonstration that lithology has a significant influence on the hydrology of a site, and that, for a given geologic terrain, the influence is fairly consistent. The geologic terrain of MCB-CL has been broadly characterized and correlated between lithologic (stratigraphic descriptions) and hydrologic (aquifer tests and well-head tests) sequences. Lithologic descriptions can now provide a good indication of hydrologic conditions at MCB-CL in areas of similar terrain.

## **GENERAL APPLICABILITY OF AQUIFER TESTS**

Aquifer (pump) tests are a problematic activity at contamination sites. While the information available from aquifer tests is required for engineering design of withdrawal systems, aquifer tests should not be a reconnaissance or an initial step in the investigation. Full consideration must be made of the redistribution of contaminants expectable from the test, of the change in structural support of disposal features by relaxation or increase of hydrostatic loading, and so forth.

Consideration must also be made of alternative sources of acceptable data on the aquifer. In the case of MCB-CL, alternatives to exploratory aquifer tests are available from the tabulation and correlation of aquifer characteristics, production performance and geologic terrain presently available.

From the available information and in light of the relative consistency of the geologic terrain of MCB-CL, exploratory tests at MCB-CL are not generally required. Therefore, exploratory tests are not advisable and should not form part of the initial investigation of a site. While they may be useful in certain circumstances after the initial investigation of a site, they should not, in the general case, be part of the investigation. Sufficiently satisfactory information is presently available to allow the initial engineering design of a groundwater response.

While exploratory aquifer tests are not advisable, performance tests of a newly installed system are highly recommended. These tests, to some extent, are a normal part of the initial operation of a system. Only minor additional monitoring and modification of the system during operation would provide data directly relevant to the long-term operation of that system.

In the Coastal Plain of MCB-CL, the information from a n exploratory data station not coincident with the long-term extraction system is not fully transferable. That is, if the test station and the recovery station are not the same, the aquifer parameters and calculations based on those parameters will differ. This means that data from an exploratory station are not more reliably usable than the data presently available, unless the

TABLE 1  
CAMP LEJEUNE PUMP TEST DATA

Well Number	Well Depth (ft,BGS)	Well Diameter (in)	Total Aquifer Thickness (ft)	Screened Length (ft)	Screened Interval (ft,BGS)	Water-level Drawdown During Pumping (ft,BGS)	Pumping Rate (Recovery wells) GPM	Duration of Pumping (min)	Specific Capacity (pumping rate/drawdown)	T (square ft/day)	K (ft/day)	S	Soils (ft,BGS)
013RW-01*	23	2	15	20	3-23	8.773	1	480	0.11	7.17	0.48	NA	0-10 silt/clay, 10-23 sand.
013MW-18	13	2	15	10	3-13	0.297	NA	480	NA	105.98	7.05	1.40E-02	0-7 silt/clay, 7-13 sand.
013MW-21	14	2	15	10	4-14	0.31	NA	480	NA	82.27	5.48	2.77E-02	0-4 silt/sand, 0-14 clay/silt
108RW-01*	15	2	9	9.1	2.45-11.55	6.38	0.5	485	0.08	5.30	0.59	NA	very fine sand
108MW-04		2	9				NA	485	NA	118.63	13.18	1.33E-02	
108MW-15	12.5	2	9	9.03	2.79-11.82		NA	485	NA	56.78	6.31	7.33E-03	0-8 sand/silt, 8-10 silt/clay
109MW-15		2	15			0.939	NA	460	NA	76.26	5.08	1.11E-02	
109MW-17	14.5	2	15	10	4.5-14.5	0.545	NA	460	NA	163.10	10.87	7.30E-03	0-15 fine sand
109RW-01*	15	2	15	9.5	2-11.5	6.265	3	460	0.48	7.80	0.52	NA	0-4 sand, 4-8 silt, 8-15 sand
110RW-01* (Drawdown,Theis)	21.8	2	50	19.2	2-21.2	9.53	3	475	0.31	200.02	4.00	NA	0-10.5 sand/silt, 10.5-15 sand/clay, 15-21.5 sand/clay, 21.5- sand
110RW-01* (Drawdown,Cooper)										161.86	3.24	NA	
110RW-01* Recovery(Theis)										106.06	2.12	NA	
110DW-01 (Drawdown,Theis)	30.3	2	50	4.8	24.9-29.7	0.02	3	475	NA	7080.48	142.00	4.52E-03	0-4 sand/silt, 4-10.5 clay, 10.5-15.5 sand/silt, 15.5-20.5 clay, 20.5-on sand
110DW-01 (Drawdown,Cooper)										7099.20	142	4.51E-03	
110DW-02 (Drawdown,Theis)	30	2	50	4.7	24.7-29.4	0.52	NA	475	NA	5398.56	108.00	1.51E-03	0-3 sand and silt with clay layers, 3-11 sand and silt, 11-30 sand with some limited clay layers
110DW-02 (Drawdown,Cooper)										5400.00	108	1.51E-03	
110DW-03 (Drawdown,Theis)	30	2	50	4.9	24.5-29.4	0.47	NA	475	NA	2952.00	59.00	7.48E-02	0-6 sand and silt , 6-12 sand, 12-23 sand/clay, 23-30 sand
110DW-03 (Drawdown,Cooper)										3225.60	64	5.85E-02	

T = Transmissivity

K = Hydraulic Conductivity

S = Storage

\* = Pumping well

NA = Not applicable

## TABLE 3

BARONE:8SEP94:CL5-1A1:1/5

MCB-CL5	CTO-232	CL5-1B1.wks		8SEP94	MCB-CAMP LEJEUNE	
STATION	b ft	Q gpm	Sc gpm/ft	T ft-sq/d	T gpd/ft	K ft/d
013RW-01	15	1.0	0.11	7.2	54	0.5
013MW-1	15			106.0	793	7.1
013MW-2	15			82.3	615	5.5
013MW-03	1					0.8
013MW-04	8					0.3
013MW-11	9					0.4
013MW-21	8					0.5
41GW-07						1.2
41GW-08						0.1
41GW-09						3.7
41GW-10						0.9
41GW-12						4.6
69GW-09						1.7
69GW-10						0.2
69GW-12						0.1
69GW-02DW						0.3
69GW-12DW						6.7
74GW-03A						0.6
74GW-06						6.3
74GW-08						3.6
108RW-01	9	0.5	0.08	5.3	40	0.6
108MW-0	9			118.6	887	13.2
108MW-1	9			56.8	425	6.3
108MW-08	9					0.6
108MW-09	8					0.5
108MW-13	8					0.1
108MW-17	8					0.6
109MW-1	15			76.3	570	5.1
109MW-1	15			163.1	1220	10.9
109RW-01	15	3.0	0.48	7.8	58	0.5
109MW-17	15					9.0
109MW-18	15					5.7
110RW-01	50	3.0	0.31	200.0	1496	4.0
110RW-01	50	3.0	0.31	161.9	1211	3.2
110RW-01	50	3.0		106.1	793	2.1
110DW-01	50			7080	52962	142.0
110DW-01	50			7099	53102	142.0
110DW-02	50			5399	40381	108.0
110DW-02	50			5400	40392	108.0
110DW-03	50			2952	22081	59.0
110DW-03	50			3226	24127	64.0
110MW-07	9					0.1
110MW-09	9					0.2
110DW-03	22					5.8

STATION	PUMPING LEVEL	Q gpm	Sc gpm/ft
HP-602	44	154	3.5
HP-603	30	129	4.3
HP-606	38	267	7.0
HP-607	46	246	5.3
HP-608	21	208	9.9
HP-609	45	199	4.4
HP-610	14	214	15.3
HP-613	17	157	9.2
HP-616	15	178	11.9
HP-620	9	224	24.9
HP-622	55	330	6.0
HP-623	30	210	7.0
HP-628	45	172	3.8
HP-629	45	216	4.8
HP-632	21	224	10.7
HP-633	18	205	11.4
HP-634	36	219	6.1
HP-635	33	151	4.6
HP-636	35	149	4.3
HP-637	40	130	3.3
HP-638	84	201	2.4
P-639	52	[--]	0.0
P-640	28	210	7.5
HP-641	44	351	8.0
HP-642	32	[--]	0.0
HP-643	35	269	7.7
HP-644	52	230	4.4
HP-645	40	192	4.8
HP-646	11	154	14.0
HP-647	26	302	11.6
HP-648	84	263	3.1
HP-649	80	100	1.3
HP-650	75	480	6.4
HP-651	69	242	3.5
HP-652	82	216	2.6
HP-653	29	197	6.8
HP-654	30	175	5.8
HP-655		[--]	ERR
HP-660		150	ERR
HP-661	37	275	7.4
HP-662	53	148	2.8
HP-663	23	100	4.3
HP-698	33	216	6.5
HP-699	21	140	6.7

BARONE:8SEP94:CL5-1A1:5/5

STATION	PUMPING LEVEL	Q gpm	Sc gpm/ft
AS-108	8	226	28.3
AS-131	11	310	28.2
AS-190	60	220	3.7
AS-191	16	220	13.8
AS-203	19	220	11.6
AS-4140	6	110	18.3
AS-4150	10	128	12.8
AS-5001	27	185	6.9
AS-5009	53	111	2.1
BA-164	21	214	10.2
BA-190	17	303	17.8

BARONE:8SEP94:CL5-1A1:5/5

**APPENDIX T**  
**SAMPLING STATION CHARACTERIZATION**  
**DATA SHEET**

---

## SAMPLING STATION CHARACTERIZATION DATA SHEET

Station Number: 1-5001 Date: 3-23-94 Time: 1605  
 Samplers: AMB, WJL, MGRM Date: \_\_\_\_\_ Time: \_\_\_\_\_  
 Water Body: Drainage Ditch State: NC County: Overslow  
 Sample Type: Fish Benthic Macroinvertebrate Sediment Surface Water  
 SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other: \_\_\_\_\_

Riparian Zone/Instream Features

Predominant Surrounding Land Use: Forest Urban Industrial Other: \_\_\_\_\_

\* Shore Vegetation: See Habitat at Map

Aquatic Vegetation: N/A

Estimated Stream Width: 2-3 ft Est. Stream Depth: N/A ft Riffle: N/A ft Run: N/A ft Pool: N/A ft

Stream Type: Cold Water Warm Water Velocity: N/A Channelized: Yes  No

Canopy Cover: Open Partly Open Partly Shaded Shaded

Sediment/Substrate:

Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other: \_\_\_\_\_

Sediment Oils: Absent Slight Moderate Profuse N/A

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: N/A Replicate #2: N/A Replicate #3: N/A

Sediment Description: Hard packed, medium grain sand

Water:

Depth	Temp. °C	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
<u>N/A</u>					

Water Odors: Normal Sewage Petroleum Chemical Other: \_\_\_\_\_

Water Surface Oils: Slick Sheen None Secchi: \_\_\_\_\_ ft.

Turbidity: Clear Slightly Turbid Turbid Opaque Water Color: \_\_\_\_\_

Weather Conditions: Sunny ~ 70-75°F Tide: N/A SL Dot

Comments: The station was located in a drainage ditch w/ no running water. Only a few percent erosion. A few days after the sample was collected a petroleum product was discharging into the ditch. Water samples were collected from this discharge.

## SAMPLING STATION CHARACTERIZATION DATA SHEET

Station Number: 1-5002Date: 3-22-94Time: 1600Samplers: AMB, NRM, WST

Date: \_\_\_\_\_

Time: \_\_\_\_\_

Water Body: Drainage DitchState: NCCounty: Cleveland

Sample Type: Fish Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other: \_\_\_\_\_Riparian Zone/Instream FeaturesPredominant Surrounding Land Use: Forest Urban Industrial Other: \_\_\_\_\_\* Shore Vegetation: See Habitat MapAquatic Vegetation: NAEstimated Stream Width: 2-3 ft Est. Stream Depth: NA ft Riffle: NA ft Run: NA ft Pool: NA ftStream Type: Cold Water Warm Water Velocity: NA Channelized: Yes  No Canopy Cover: Open Partly Open Partly Shaded ShadedSediment/Substrate:Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other: \_\_\_\_\_Sediment Oils: Absent Slight Moderate Profuse HNuPonar Grab: Number of Jars Filled with Sediments Replicate #1: NA Replicate #2: NA Replicate #3: NASediment Description: Fine, silty sand, light brownWater:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
<u>NA</u>					

Water Odors: Normal Sewage Petroleum Chemical Other: NA

Water Surface Oils: Slick Sheen None Secchi: \_\_\_\_\_ ft.

Turbidity: Clear Slightly Turbid Turbid Opaque Water Color: \_\_\_\_\_

Weather Conditions: Sunny ~ 70-75°F Tide: In OutComments: There were a few puddles of water in the ditch but it was mostly dry.

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-NR-Sw15D01 Date: March 28, 94 (SD) Time: 1642 (SD)  
 Samplers: Amb, WTT, m6m Date: March 28, 94 (SW) Time: 1315 (SW)  
 Water Body: New River State: NC County: Carson  
 Sample Type: Fish Benthic Macroinvertebrate  Sediment  Surface Water  
 SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer  Sediment Corer Spoon Other: Dip Net

Riparian Zone/Instream Features

Predominant Surrounding Land Use:  Forest  Urban  Industrial Other: \_\_\_\_\_

Shore Vegetation: See Habitat Map

Aquatic Vegetation: None

Estimated Stream Width: NM ft Est. Stream Depth: NM ft Riffle: \_\_\_\_\_ ft Run: 1006 ft Pool: \_\_\_\_\_ ft

Stream Type: Cold Water  Warm Water Velocity: Vegetable Channelized: Yes  No

Canopy Cover:  Open  Partly Open  Partly Shaded  Shaded

Sediment/Substrate:

Sediment Odors:  Normal Sewage Petroleum Chemical Anaerobic Other: \_\_\_\_\_

Sediment Oils:  Absent Slight Moderate Profuse HNU

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: NA Replicate #2: NA Replicate #3: NA

Sediment Description: Cours sand / clay

Water:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	22.5	6.83	11	18000	15

Water Odors:  Normal Sewage Petroleum Chemical Other: \_\_\_\_\_

Water Surface Oils: Slick Sheen None Secchi: NA ft.

Turbidity: Clear Slightly Turbid  Turbid Opaque Water Color: Clear

Weather Conditions: \_\_\_\_\_ Tide: In Out

Comments: NM - not measured due to large size of river

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-NR-SW/SD02 Date: 3-28-94 (SW) Time: 1355 (SW)  
 Samplers: AMBI, WTT, M601 Date: 3-28-94 (SD) Time: 1045 (SD)  
 Water Body: New River State: NC County: Brown  
 Sample Type: Fish Benthic Macroinvertebrate Sediment Surface Water  
 SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other: Dip net/ hand

Riparian Zone/Instream Features

Predominant Surrounding Land Use: Forest Urban Industrial Other:

Shore Vegetation: See Habitat maps

Aquatic Vegetation: NONE

Estimated Stream Width: \*NM ft Est. Stream Depth: NM ft Riffle:    ft Run: 1006 ft Pool:    ft

Stream Type: Cold Water Warm Water Velocity: Negligible Channelized: Yes    No ✓

Canopy Cover: Open Partly Open Partly Shaded Shaded

Sediment/Substrate:

Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other:

Sediment Oils: Absent Slight Moderate Profuse HNA

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: N/A Replicate #2: N/A Replicate #3: N/A

Sediment Description: Sandy soil

Water:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface.	22.5	6.94	10.5	18,000	13.5
bottom					

Water Odors: Normal Sewage Petroleum Chemical Other:

Water Surface Oils: Slick Sheen None Secchi: N/A ft.

Turbidity: Clear Slightly Turbid Turbid Opaque Water Color:

Weather Conditions: Tide: ? In Out

Comments: \*NM - not measured due to large size of river

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-NR-54/SD03

Date: March 25, 94 (SD)

Time: 1719 (SD)

Samplers: AMB, WJJ, MFM

Date: March 28, 94 (SW)

Time: 1710 (SW)

Water Body: New River

State: NC

County: Onslow

Sample Type: Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine  Gill Net  Ponar  Kemmerer  Sediment Corer  Spoon  Other: Dip Method

Riparian Zone/Instream Features

Predominant Surrounding Land Use:  Forest  Urban  Industrial  Other: \_\_\_\_\_

Shore Vegetation: See Habitat Map

Aquatic Vegetation: None

Estimated Stream Width: \*NM ft Est. Stream Depth: \*NM ft Riffle: \_\_\_\_\_ ft Run: 1006 ft Pool: \_\_\_\_\_ ft

Stream Type: Cold Water  Warm Water Velocity: Negligible Channelized: Yes  No

Canopy Cover:  Open  Partly Open  Partly Shaded  Shaded

Sediment/Substrate:

Sediment Odors:  Normal  Sewage  Petroleum  Chemical  Anaerobic  Other: \_\_\_\_\_

Sediment Oils:  Absent  Slight  Moderate  Profuse  HNU

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: NA Replicate #2: NA Replicate #3: NA

Sediment Description: Sandy Silt

Water:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface.	20.0	7.95	11.0	12,000	15

Water Odors:  Normal  Sewage  Petroleum  Chemical  Other: \_\_\_\_\_

Water Surface Oils:  Slick  Sheen  None Secchi: NA ft

Turbidity:  Clear  Slightly Turbid  Turbid Opaque Water Color: Clear

Weather Conditions: \_\_\_\_\_ Tide:  In  Out

Comments: \*NM - Not measured due to large size of river

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-NR-su/SD04 Date: March 25, 84 (SD) Time: 1659 (SD)  
 Samplers: AMB, w/3, w/6w Date: March 29, 84 (su) Time: 1310 (su)  
 Water Body: New River State: NC County: Onslow  
 Sample Type: Fish Benthic Macroinvertebrate  Sediment Surface Water  
 SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer  Sediment Corer Spoon Other:

Riparian Zone/Instream Features

Predominant Surrounding Land Use:  Forest Urban Industrial Other:

Shore Vegetation: See Habitat Map

Aquatic Vegetation: None

Estimated Stream Width: NM ft Est. Stream Depth: NM ft Riffle: — ft Run: 1006 ft Pool: — ft

Stream Type: Cold Water  Warm Water Velocity: Neg/6/0 Channelized: Yes  No

Canopy Cover:  Open Partly Open Partly Shaded Shaded

Sediment/Substrate:

Sediment Odors:  Normal Sewage Petroleum Chemical Anaerobic Other:

Sediment Oils:  Absent Slight Moderate Profuse NH

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: NA Replicate #2: NA Replicate #3: NA

Sediment Description: Course Sand w/ high organic content

Water:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	19.0	9.23	7.4	400	0.25

Water Odors:  Normal Sewage Petroleum Chemical Other:

Water Surface Oils: Slick Sheen  None Secchi: NH ft.

Turbidity: Clear Slightly Turbid  Turbid Opaque Water Color: Clear

Weather Conditions: Tide: ? In Out

Comments: NM - Not measured due to large size of River  
located downstream of Coalfield creek

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-NR-sw/5005 Date: March 25, 94 (SD) Time: 1606 (SD)  
 Samplers: AMB, WJJ, mtm Date: March 29, 94 (SW) Time: 1340 (SW)  
 Water Body: New River State: NC County: Ondow  
 Sample Type: Fish Benthic Macroinvertebrate  Sediment Surface Water  
 SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer  Sediment Corer Spoon Other:

Riparian Zone/Instream Features

Predominant Surrounding Land Use:  Forest Urban Industrial Other:

Shore Vegetation: Soil Habitat n/a

Aquatic Vegetation: N/A

Estimated Stream Width: NA ft Est. Stream Depth: NA ft Riffle: — ft Run: 1006 ft Pool: — ft

Stream Type: Cold Water  Warm Water Velocity: High Channelized: Yes  No

Canopy Cover:  Open Partly Open Partly Shaded Shaded

Sediment/Substrate:

Sediment Odors:  Normal Sewage Petroleum Chemical Anaerobic Other:

Sediment Oils:  Absent Slight Moderate Profuse HNu

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: NA Replicate #2: NA Replicate #3: NA

Sediment Description: Very clayey

Water:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	19	9.76	10	3800	3.5

Water Odors:  Normal Sewage Petroleum Chemical Other:

Water Surface Oils: Slick Sheen  None Secchi: NA ft

Turbidity: Clear Slightly Turbid  Turbid Opaque Water Color:

Weather Conditions: Tide: ? In Out

Comments: \* NA - Not Applicable due to large size of river

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-CC-SW/SD01

Date: March 26, 94 (61)

Time: 0915 (SD)

Samplers: Amb, WTT, WGW

Date: March 29, 94 (SW)

Time: 1250 (SW)

Water Body: Codger's Creek

State: NC

County: Owen

Sample Type: Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other:

Riparian Zone/Instream Features

Predominant Surrounding Land Use: Forest Urban Industrial Other:

Shore Vegetation: See Habitat Map

Aquatic Vegetation: None

Estimated Stream Width: ~30' ft Est. Stream Depth: 3-4 ft Riffle: — ft Run: 100 ft Pool: — ft

Stream Type: Cold Water Warm Water Velocity: Very slow Channelized: Yes — No ✓

Canopy Cover: Open Partly Open

Partly Shaded

Shaded

Sediment/Substrate:

Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other:

Sediment Oils: Absent Slight Moderate Profuse HNU

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: NA Replicate #2: NA Replicate #3: NA

Sediment Description: Organic muck

Water:

Depth	Temp. °C	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	20	8.73	6.0	600	0.25

Water Odors: Normal Sewage Petroleum Chemical Other:

Water Surface Oils: Slick Sheen None Secchi: NA ft.

Turbidity: Clear Slightly Turbid Turbid Opaque Water Color:

Weather Conditions: Tide: ? In Out

Comments: Sample was collected near the staff gauge ~8' from back due to access problems

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-CC-sw/5002

Date: March 27, 94 (sw)

Time: 1017 (sw)

Samplers: AWB, WJJ, MFM

Date: March 27, 94 (SD)

Time: 1036 (SD)

Water Body: Codyels Creek

State: NC

County: O�试

Sample Type: Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other: Dip Method

Riparian Zone/Instream Features

Predominant Surrounding Land Use: Forest Urban Industrial Other:

Shore Vegetation: See Habitat map

Aquatic Vegetation: None

Estimated Stream Width: 25-30' ft Est. Stream Depth: NM ft Riffle: — ft Run: 1006 ft Pool: — ft

Stream Type: Cold Water Warm Water Velocity: Negligible Channelized: Yes    No ✓

Canopy Cover: Open Partly Open Partly Shaded Shaded

Sediment/Substrate:

Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other:

Sediment Oils: Absent Slight Moderate Profuse HNU

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: NA Replicate #2: NA Replicate #3: NA

Sediment Description: Silty mud w/ sand

Water:

Depth	Temp. °C	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	22	6.43	7.74	1400	1.
Bottom	22	6.97	7.74	7500	9

Water Odors: Normal Sewage Petroleum Chemical Other:

Water Surface Oils: Slick Sheen None Secchi: NA ft

Turbidity: Clear Slightly Turbid Turbid Opaque Water Color: Clear

Weather Conditions: Tide: In Out

Comments: NM - Unable to measure due to access -

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-CC-su/SD03 Date: March 27, 94 (su) Time: 1110 (su)  
 Samplers: Amb, WTT, mem Date: March 28, 94 (SD) Time: 1655 (SD)  
 Water Body: Codger's Creek State: NC County: Owlston  
 Sample Type: Fish Benthic Macroinvertebrate Sediment Surface Water

SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other: Dip net/box

Riparian Zone/Instream Features

Predominant Surrounding Land Use:  Forest  Urban  Industrial Other: \_\_\_\_\_

Shore Vegetation: See Habitat Map

Aquatic Vegetation: None

Estimated Stream Width: 20 ft Est. Stream Depth: ~3-4 ft Riffle: — ft Run: 100 ft Pool: — ft

Stream Type: Cold Water  Warm Water Velocity: Negligible Channelized: Yes  No

Canopy Cover: Open  Partly Open  Partly Shaded  Shaded

Sediment/Substrate:

Sediment Odors: Normal Sewage Petroleum Chemical  Anaerobic Other: \_\_\_\_\_

Sediment Oils:  Absent Slight Moderate Profuse  HNU

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: NA Replicate #2: NA Replicate #3: NA

Sediment Description: Silty Sand

Water:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	17	6.82	8.3	350	6.82
Bottom	16	6.79	8.3	9500	6.79

Water Odors:  Normal Sewage Petroleum Chemical Other: \_\_\_\_\_

Water Surface Oils: Slick Sheen  None Secchi: NA ft.

Turbidity: Clear Slightly Turbid  Turbid  Opaque Water Color: Tan

Weather Conditions: \_\_\_\_\_ Tide: In Out

Comments: Just downstream of road by the staff gauge from right back facing down

## SAMPLING STATION CHARACTERIZATION DATA SHEET

Station Number: 28-CC-SW15D04Date: March 27, 94 (SW)Time: 1153 (SW)Samplers: AMB, MCM, WWDDate: March 28, 94 (SD)Time: 1600 (SD)Water Body: Cedars CreekState: NCCounty: O�试

Sample Type: Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine    Gill Net    Ponar — Kemmerer    Sediment Corer    Spoon    Other: Dip net methodRiparian Zone/Instream FeaturesPredominant Surrounding Land Use: Forest    Urban    Industrial    Other: \_\_\_\_\_Shore Vegetation: See Habitat mapAquatic Vegetation: NoneEstimated Stream Width: ~40 ft    Est. Stream Depth: ~4.5 ft    Riffle: — ft    Run: 1006 ft    Pool: — ftStream Type: Cold Water    Warm Water    Velocity: Negligible    Channelized: Yes ✓    No —Canopy Cover: Open    Partly Open    Partly Shaded    ShadedSediment/Substrate:Sediment Odors: Normal    Sewage    Petroleum    Chemical    Anaerobic    Other: \_\_\_\_\_Sediment Oils: Absent    Slight    Moderate    Profuse    HNuPonar Grab: Number of Jars Filled with Sediments    Replicate #1: NA    Replicate #2: NA    Replicate #3: NASediment Description: Black organic material / soilWater:

Depth	Temp. °C	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	20.2	7.01	9.9	6,800	6.9

Water Odors: Normal    Sewage    Petroleum    Chemical    Other: \_\_\_\_\_Water Surface Oils: Slick    Sheen    None    Secchi: NA    ft.Turbidity: Clear    Slightly Turbid    Turbid    Opaque    Water Color: TanWeather Conditions: \_\_\_\_\_ Tide: In    OutComments: On right side of bank (looking down) between two trib.

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-CC-SW15D05

Date: March 27, 94 (sw)

Time: 1246 (sw)

Samplers: AMB, WJJ, M6M

Date: March 27, 94 (SD)

Time: 1259 (SD)

Water Body: Codgol's Creek

State: NC

County: Onslow

Sample Type:

Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine    Gill Net    Ponar    Kemmerer     Sediment Corer    Spoon    Other: Dia Method

Riparian Zone/Instream Features

Predominant Surrounding Land Use:  Forest

Urban

Industrial

Other: \_\_\_\_\_

Shore Vegetation: See Habitat Map

Aquatic Vegetation: None

Estimated Stream Width: 30 ft    Est. Stream Depth: 2-3 ft    Riffle:       ft    Run: 1006 ft    Pool:       ft

Stream Type: Cold Water    Warm Water    Velocity: Very slow    Channelized: Yes       No

Canopy Cover: Open    Partly Open

Partly Shaded

Shaded

Sediment/Substrate:

Sediment Odors: Normal    Sewage    Petroleum    Chemical     Anaerobic    Other: \_\_\_\_\_

Sediment Oils:  Absent    Slight    Moderate    Profuse    NN

Ponar Grab: Number of Jars Filled with Sediments    Replicate #1: NA    Replicate #2: NA    Replicate #3: NA

Sediment Description: Silty soil w/ peat material

Water:

Depth	Temp. °C	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface.	17.5	6.67	5.6	0	272

Water Odors:  Normal    Sewage    Petroleum    Chemical    Other: \_\_\_\_\_

Water Surface Oils: Slick    Sheen     None    Secchi: NA ft.

Turbidity: Clear    Slightly Turbid     Turbid    Opaque    Water Color: Teal

Weather Conditions: \_\_\_\_\_ Tide: In  Out

Comments: ~10' up from mouth on trib (right bank)

# SAMPLING STATION CHARACTERIZATION DATA SHEET

Station Number: 2B-CC-SW/SD 06

Date: March 27, 94 (sw)

Time: 1435 (sw)

Samplers: AMB, ETJ, MLO

Date: March 27, 94 (SD)

Time: 1755 (SD)

Water Body: Cogdol's Creek

State: NC

County: Owlens

Sample Type: Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other: Dip Method

## Riparian Zone/Instream Features

Predominant Surrounding Land Use: Forest

Urban

Industrial

Other:

Shore Vegetation: See Habitat Map

Aquatic Vegetation: None

Estimated Stream Width: 420 ft Est. Stream Depth: 2-2 ft Riffle: 40 ft Run: 100 ft Pool: — ft

Stream Type: Cold Water

Warm Water

Velocity: High

Channelized: Yes —

No ✓

Canopy Cover: Open

Partly Open

Partly Shaded

Shaded

## Sediment/Substrate:

Sediment Odors: Normal

Sewage

Petroleum

Chemical

Anaerobic

Other:

Sediment Oils: Absent

Slight

Moderate

Profuse

HNu

Ponar Grab: Number of Jars Filled with Sediments

Replicate #1: NA

Replicate #2: NA

Replicate #3: NA

Sediment Description: Silty mud

## Water:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface.	(9.5	6.9	4.75	500	0.25

Water Odors: Normal Sewage Petroleum Chemical Other:

Water Surface Oils: Slick Sheen None Secchi: NA ft.

Turbidity: Clear Slightly Turbid Turbid Opaque Water Color: Tan

Weather Conditions: Tide: ? In Out

Comments: n 75' up stream of staff gauge

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-CC-SW/SD07 Date: March 27, 94 (SW) Time: 1327 (SW)  
 Samplers: AM13, WJJ, MGM Date: March 27, 94 (SD) Time: 1358 (SD)  
 Water Body: Codgels Creek State: NC County: Oastow  
 Sample Type: Fish Benthic Macroinvertebrate Sediment Surface Water  
 SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other: Dip Method

Riparian Zone/Instream Features

Predominant Surrounding Land Use: Forest Urban Industrial Other: \_\_\_\_\_

Shore Vegetation: See Habitat Map

Aquatic Vegetation: None

Estimated Stream Width: 20 ft Est. Stream Depth: NM ft Rifle: — ft Run: 1006 ft Pool: — ft

Stream Type: Cold Water Warm Water Velocity: Negligible Channelized: Yes    No ✓

Canopy Cover: Open Partly Open Partly Shaded Shaded

Sediment/Substrate:

Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other: \_\_\_\_\_

Sediment Oils: Absent Slight Moderate Profuse HNU

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: NA Replicate #2: NA Replicate #3: NA

Sediment Description: Silty Muds

Water:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
<u>17.8 Surface</u>	<u>17.5</u>	<u>6.73</u>	<u>4.4</u>	<u>335</u>	<u>0</u>

Water Odors: Normal Sewage Petroleum Chemical Other: \_\_\_\_\_

Water Surface Oils: Slick Sheen None Secchi: NA ft

Turbidity: Clear Slightly Turbid Turbid Opaque Water Color: Tan

Weather Conditions: \_\_\_\_\_ Tide: In Out

Comments: \*NM - Not able to measure.

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 2B-OP-sw/SD 01

Date: March 28, 94 (SD)

Time: 1015

Samplers: Amb, w/31, w/6m

Date: March 28, 94 (SD)

Time: 1215 (SD)

Water Body: Orde Pond

State: NC

County: O�试

Sample Type: Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine  Gill Net  Ponar  Kemmerer  Sediment Corer  Spoon  Other: Dip method

Riparian Zone/Instream Features

Predominant Surrounding Land Use:  Forest  Urban  Industrial

Other: \_\_\_\_\_

Shore Vegetation: See Habitat map

Aquatic Vegetation: NONE

Estimated Stream Width: \_\_\_\_\_ ft Est. Stream Depth: ~8' ft Riffle: \_\_\_\_\_ ft Run: \_\_\_\_\_ ft Pool: 100ft ft

Stream Type: Cold Water  Warm Water Velocity: NA Channelized: Yes  No

Canopy Cover:  Open Partly Open Partly Shaded Shaded

Sediment/Substrate:

Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other: \_\_\_\_\_

Sediment Oils: Absent Slight Moderate Profuse  N/A

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: NA Replicate #2: NA Replicate #3: NA

Sediment Description: clayey silt

Water:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	19.4	6.94	9.0	70	0
Bottom	19.4	7.03	9.0	70	0

Water Odors:  Normal Sewage Petroleum Chemical Other: \_\_\_\_\_

Water Surface Oils: Slick Sheen  None Secchi: NA ft

Turbidity:  Clear Slightly Turbid Turbid Opaque Water Color: Clear

Weather Conditions: \_\_\_\_\_ Tide: NA In Out

Comments: At left side of pond (w/ back to trt plant)

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 2B-OP-Sw/SD02

Date: March 28, 94 (Sw)

Time: 0955 (Sw)

Samplers: Amb, WTT, M6-m1

Date: March 28, 94 (SD)

Time: 1600 (SD)

Water Body: Orde Pond

State: NC

County: Buncombe

Sample Type: Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine  Gill Net  Ponar  Kemmerer  Sediment Corer  Spoon  Other: Dip Method

Riparian Zone/Instream Features

Predominant Surrounding Land Use: Forest  Urban  Industrial  Other: \_\_\_\_\_

Shore Vegetation: See Habitat Map

Aquatic Vegetation: NONE

Estimated Stream Width: \_\_\_\_\_ ft Est. Stream Depth: 8' ft Riffle: \_\_\_\_\_ ft Run: \_\_\_\_\_ ft Pool: 106 ft

Stream Type: Cold Water  Warm Water  Velocity: NA Channelized: Yes  No

Canopy Cover: Open  Partly Open  Partly Shaded  Shaded

Sediment/Substrate:

Sediment Odors: Normal  Sewage  Petroleum  Chemical  Anaerobic  Other: \_\_\_\_\_

Sediment Oils: Absent  Slight  Moderate  Profuse  HNU

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: NA Replicate #2: NA Replicate #3: NA

Sediment Description: \_\_\_\_\_

Water:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	20	6.79	9.0	72	0
Bottom	19.5	6.93	9.0	88	0

Water Odors: Normal  Sewage  Petroleum  Chemical  Other: \_\_\_\_\_

Water Surface Oils: Slick  Sheen  None  Secchi: NA ft.

Turbidity: Clear  Slightly Turbid  Turbid  Opaque  Water Color: Clear (slightly brown)

Weather Conditions: \_\_\_\_\_ Tide: NA In  Out

Comments: At right side of pond (w/ back to trt plant)

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-FS1Bn01

Date: March 23, 94

Time: 0920

Samplers: AMB, WJJ, MGR

Date: March 27, 94 (Bw)

Time: 1415 (Bw)

Water Body: Codgat's Creek

State: NC

County: O�试

Sample Type: Fish Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other: Minnow Traps

Riparian Zone/Instream Features

Predominant Surrounding Land Use: Forest Urban Industrial Other: \_\_\_\_\_

Shore Vegetation: See Habitat Map

Aquatic Vegetation: None

Estimated Stream Width: 30 ft Est. Stream Depth: 1 ft Riffle: — ft Run: 1006 ft Pool: — ft

Stream Type: Cold Water Warm Water Velocity: Vegetable Channelized: Yes — No X

Canopy Cover: Open Partly Open Partly Shaded ~75% Shaded

Sediment/Substrate:

Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other: \_\_\_\_\_

Sediment Oils: Absent Slight Moderate Profuse Hnu

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: 5 Replicate #2: 3 Replicate #3: 3

Sediment Description: Fine silt, dark brown, many sticks, debris

Water:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	13.3	6.8	4.8	440	0

Water Odors: Normal Sewage Petroleum Chemical Other: \_\_\_\_\_

Water Surface Oils: Slick Sheen None Secchi: N/A ft.

Turbidity: Clear Slightly Turbid Turbid Opaque Water Color: Tan

Weather Conditions: Sunny Tide: In Out

Comments: By 28-CC-SW/SD 05

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-FS1BN02

Date: March 23, 94

Time: 1010

Samplers: Amb, WJJ, MGR

Date: March 27, 94 (Bn)

Time: 1449 (BN)

Water Body: Codgills Creek

State: NC

County: O�试

Sample Type:  Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine  Gill Net  Ponar Kemmerer Sediment Corer Spoon ...Other: \_\_\_\_\_

Riparian Zone/Instream Features

Predominant Surrounding Land Use:  Forest Urban Industrial Other: Open meadow

Shore Vegetation: See Habitat Map

Aquatic Vegetation: None

Estimated Stream Width: 40 ft Est. Stream Depth: 1-2 ft Riffle: — ft Run: 1008 ft Pool: — ft

Stream Type: Cold Water  Warm Water Velocity: Negligible Channelized: Yes — No

Canopy Cover: Open  Partly Open 50%  Partly Shaded Shaded

Sediment/Substrate:

Sediment Odors: Normal Sewage Petroleum Chemical  Anaerobic Other: \_\_\_\_\_

Sediment Oils:  Absent Slight Moderate Profuse  HNu

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: 3 Replicate #2: 5 Replicate #3: 6

Sediment Description: Fine silt-sand, 25-35% organic debris (sticks, leaves, needles)

Water:

Depth	Temp. °C	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	14.5	6.8	6.5	351	0
Bn	Surface	17.5	6.92	7.5	320
	Bottom	17.5	6.77	6.0	3,100

Water Odors:  Normal Sewage Petroleum Chemical Other: \_\_\_\_\_

Water Surface Oils: Slick Sheen  None Secchi: 14 ft.

Turbidity: Clear  Slightly Turbid Turbid Opaque Water Color: Toxic

Weather Conditions: Clear sunny Tide: In  Out

Comments: By 28-CC-54/SD04

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 28-FS1B NO 3

Date: 3-24-94

Time: 1015

Samplers: AMB, WTS, mgmt

Date: 3-26-94

Time: 1212

Water Body: Codyels Creek

State: NC

County: Overlow

Sample Type: Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other: \_\_\_\_\_

Riparian Zone/Instream Features

Predominant Surrounding Land Use: Forest Urban Industrial Other: \_\_\_\_\_

Shore Vegetation: Mixed Hardwoods - Seed Habitat Was

Aquatic Vegetation: N/A

Estimated Stream Width: 30-40 ft Est. Stream Depth: 3-4 ft. Riffle: - ft Run: 100 ft Pool: - ft

Stream Type: Cold Water Warm Water Velocity: Very slow Channelized: Yes — No ✓

Canopy Cover: Open Partly Open

Partly Shaded

Shaded

Sediment/Substrate:

Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other: \_\_\_\_\_

Sediment Oils: Absent Slight Moderate Profuse NH4+

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: \_\_\_\_\_ Replicate #2: \_\_\_\_\_ Replicate #3: \_\_\_\_\_

Sediment Description: Silt, (Cherry) w/ organic material, no sewage sand ironstone - shell fragments, no worms observed

Water:

Depth	Temp. °C	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	22.5	3.5	6.8	1150	0
Bottom	22.5	3.5	6.8	1150	0

Water Odors: Normal Sewage Petroleum Chemical Other: \_\_\_\_\_

Water Surface Oils: Slick Sheen None Secchi: N/A ft.

Turbidity: Clear Slightly Turbid Turbid Opaque Water Color: Tan

Weather Conditions: \_\_\_\_\_ Tide: In Out

Comments: Just down from staff gauge

## SAMPLING STATION CHARACTERIZATION DATA SHEET

Station Number: 28-FS/BN-04 Date: 3-23-94 Time: 1300  
 Samplers: AMB, ETJ, m6-m7 Date: 3-26-94 Time: 1315 (PM)  
 Water Body: New River State: NC County: Oastow  
 Sample Type: Fish Benthic Macroinvertebrate Sediment Surface Water  
 SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other:

Riparian Zone/Instream Features

Predominant Surrounding Land Use: Forest Urban Industrial Other:

Shore Vegetation: See Habitat Map

Aquatic Vegetation: None

\* Estimated Stream Width: NA ft Est. Stream Depth: NA ft Riffle: NA ft Run: NA ft Pool: NA ft

Stream Type: Cold Water Warm Water Velocity: Negligible Channelized: Yes        No X

Canopy Cover: Open Partly Open Partly Shaded Shaded

Sediment/Substrate:

Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other:

Sediment Oils: Absent Slight Moderate Profuse None

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: 1 Replicate #2: 1 Replicate #3: 1

Sediment Description: 90% fine sand + 10% gravel Fragments

Water:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	20.2	7.01	9.9	8,000	6.9

Water Odors: Normal Sewage Petroleum Chemical Other:

Water Surface Oils: Slick Sheen None Secchi: NA ft.

Turbidity: Clear Slightly Turbid Turbid Opaque Water Color: Brown

Weather Conditions: Sunny, Warm (60°F) Tide: In ? Out

Comments: \* the sample was collected ~ 3-5' from shore in the new river. The depth was ~ 2ft here. Fish collected over several days via gill nets. Debris on rocks, roots, bricks.

## SAMPLING STATION CHARACTERIZATION DATA SHEET

Station Number: 28-F51BN05Date: 3-24-94Time: 1015Samplers: AMB, WTT, M6MDate: 3-26-94 (pw)Time: 1245Water Body: New RiverState: NCCounty: Owens

Sample Type:

 Fish Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine  Gill Net  Ponar Kemmerer Sediment Corer Spoon Other: \_\_\_\_\_Riparian Zone/Instream FeaturesPredominant Surrounding Land Use:  Forest Urban Industrial Other: \_\_\_\_\_Shore Vegetation: See Habitat MapAquatic Vegetation: None\* Estimated Stream Width: NA ft Est. Stream Depth: NA ft Riffle: NA ft Run: NA ft Pool: NA ftStream Type: Cold Water  Warm Water Velocity: NA, 1.6/0 Channelized: Yes  No Canopy Cover:  Open Partly Open Partly Shaded ShadedSediment/Substrate:Sediment Odors:  Normal Sewage Petroleum Chemical Anaerobic Other: \_\_\_\_\_Sediment Oils:  Absent Slight Moderate Profuse  HNuPonar Grab: Number of Jars Filled with Sediments Replicate #1: 1 Replicate #2: 1 Replicate #3: 1Sediment Description: Silty Clay w/ some sand inclusions, shell fragmentsWater:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	25	NA	NA	14,000	13
BW	21	7.69	10.5	11,800	10.2

Water Odors:  Normal Sewage Petroleum Chemical Other: \_\_\_\_\_Water Surface Oils: Slick Sheen  None Secchi: NA ft.Turbidity: Clear Slightly Turbid  Turbid Opaque Water Color: Brown

Weather Conditions: \_\_\_\_\_ Tide: In Out

Comments: \* The sample was collected ~ 3-5' from shore in the New River.

The depth was ~1ft here: Fish were collected over several days via gill nets

Some further traps in area

**SAMPLING STATION CHARACTERIZATION DATA SHEET**

Station Number: 2B-OP-B5/RN

Date: March 29, 94 (Bn)

Time: 1330 (a.s.)

Samplers: AMPA, M609, WSS

Date: \_\_\_\_\_

Time: \_\_\_\_\_

Water Body: Onde Park

State: North Carolina

County: Ondow

Sample Type: Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other: Electrofish

Riparian Zone/Instream Features

Predominant Surrounding Land Use: Forest Urban Industrial Other: \_\_\_\_\_

Shore Vegetation: See Hub, lot map

Aquatic Vegetation: None

Estimated Stream Width: \_\_\_\_\_ ft Est. Stream Depth: 8' ft Riffle: \_\_\_\_\_ ft Run: \_\_\_\_\_ ft Pool: 1008 ft

Stream Type: Cold Water Warm Water Velocity: NA Channelized: Yes    No ✓

Canopy Cover: Open Partly Open Partly Shaded Shaded

Sediment/Substrate:

Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other: \_\_\_\_\_

Sediment Oils: Absent Slight Moderate Profuse HN

Ponar Grab: Number of Jars Filled with Sediments Replicate #1: \_\_\_\_\_ Replicate #2: \_\_\_\_\_ Replicate #3: \_\_\_\_\_

Sediment Description: Very silty/clay, many red worms

w/ some grass

Water:

Depth	Temp. °C	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)

Water Odors: Normal Sewage Petroleum Chemical Other: \_\_\_\_\_

Water Surface Oils: Slick Sheen None Secchi: NA ft.

Turbidity: Clear Slightly Turbid Turbid Opaque Water Color: clear

Weather Conditions: \_\_\_\_\_ Tide: NA In Out

Comments: \_\_\_\_\_

## SAMPLING STATION CHARACTERIZATION DATA SHEET

Station Number: 30-sw/5001Date: 3-23-94 (sw)Time: 1805 (sw)Samplers: AMB, MGR, WTTDate: 3-25-94 (SD)Time: 1200 (SD)Water Body: Frouch CreekState: NCCounty: Onslow

Sample Type: Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other: Dip for swRiparian Zone/Instream FeaturesPredominant Surrounding Land Use: Forest Urban Industrial Other:Shore Vegetation: See Habitat MapAquatic Vegetation: NoneEstimated Stream Width: 2' ft Est. Stream Depth: 0.5 ft Riffle: — ft Run: 1006 ft Pool: — ftStream Type: Cold Water Warm Water Velocity: Mobil Channelized: Yes   No XCanopy Cover: Open Partly Open Partly Shaded ShadedSediment/Substrate:Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other:Sediment Oils: Absent Slight Moderate Profuse HNUPonar Grab: Number of Jars Filled with Sediments Replicate: #1: NA Replicate #2: NA Replicate #3: NASediment Description: Silty w/o organic matter Silty muckWater:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	13.7	4.1	2.2	16.6	0

Water Odors: Normal Sewage Petroleum Chemical Other: NoneWater Surface Oils: Slick Sheen None Secchi: NA ft.Turbidity: Clear Slightly Turbid Turbid Opaque Water Color: TanWeather Conditions: Tide: NA In Out

Comments:

## SAMPLING STATION CHARACTERIZATION DATA SHEET

Station Number: 30-SW/6002Date: 3-23-94 (sw)Time: 1753 (sw)Samplers: AMB, WJJ, MAMDate: 3-24-94 (SD)Time: 1725 (SD)Water Body: Fresh CreekState: NCCounty: O�试

Sample Type:

Fish

Benthic Macroinvertebrate

Sediment

Surface Water

SAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer Sediment Corer Spoon Other: Dip (sw)Riparian Zone/Instream FeaturesPredominant Surrounding Land Use: Forest Urban Industrial Other: Adjacent to RoadShore Vegetation: See Habitat MapAquatic Vegetation: NoneEstimated Stream Width: 3 ft Est. Stream Depth: 1 ft Riffle: — ft Run: 506 ft Pool: 506 ftStream Type: Cold Water Warm Water Velocity: Mg/10 Channelized: Yes   No XCanopy Cover: Open Partly Open Partly Shaded ShadedSediment/Substrate:Sediment Odors: Normal Sewage Petroleum Chemical Anaerobic Other:  Sediment Oils: Absent Slight Moderate Profuse HnuPonar Grab: Number of Jars Filled with Sediments Replicate #1: NA Replicate #2: NA Replicate #3: NASediment Description: Silty muckWater:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	16.4	5.1	6.2	11.5	0

Water Odors: Normal Sewage Petroleum Chemical Other: NoneWater Surface Oils: Slick Sheen None Secchi: NA ft.Turbidity: Clear Slightly Turbid Turbid Opaque Water Color: TanWeather Conditions: Tide: NA In OutComments: Staff gauge - 1.05

## SAMPLING STATION CHARACTERIZATION DATA SHEET

Station Number: 30-sw/SD 03Date: 3-23-94 (sw)Time: 1718 (sw)Samplers: AMB, WJJ, MGMDate: 3-24-94 (SD)Time: 1635 (SD)Water Body: French CreekState: NCCounty: Oakland

Sample Type: Fish Benthic Macroinvertebrate

 Sediment Surface WaterSAMPLING EQUIPMENT: Seine Gill Net Ponar Kemmerer  Sediment Corer Spoon Other: N/A (sw)Riparian Zone/Instream FeaturesPredominant Surrounding Land Use:  Forest Urban Industrial Other: \_\_\_\_\_Shore Vegetation: Trees (90%), Scrub/Chrub (5%), Herb (5%), (Hardwood veg) Both sides bordered by Coniferous Forest - See Habitat MapAquatic Vegetation: NoneEstimated Stream Width: 4 ft Est. Stream Depth: 0.5 ft Riffle: 106 ft Run: 906 ft Pool: — ftStream Type: Cold Water  Warm Water Velocity: \_\_\_\_\_ Channelized: Yes  No —Canopy Cover: Open Partly Open Partly Shaded  ShadedSediment/Substrate:Sediment Odors: Normal Sewage Petroleum Chemical  Anaerobic Other: \_\_\_\_\_Sediment Oils:  Absent Slight Moderate Profuse HnuPonar Grab: Number of Jars Filled with Sediments Replicate #1: N/A Replicate #2: N/A Replicate #3: N/ASediment Description: Silty / Clayey siltWater:

Depth	Temp. °C.	pH (s.u.)	Dissolved Oxygen (mg/L)	Conductivity (micromhos/cm)	Salinity (ppt)
Surface	15.6	5.5	5.4	286	0

Water Odors: Normal Sewage Petroleum Chemical Other: NoneWater Surface Oils: Slick Sheen  None Secchi: N/A ft.Turbidity: Clear  Slightly Turbid Turbid Opaque Water Color: TanWeather Conditions: Clear, 70°F, Breezy Tide: N/A  9ftComments: STAFF gauge - 1.3

**APPENDIX U**  
**BIOTA POPULATION**

---



SPECIES	COO SAMPLE NO.	DRIDE POND			COODLES CREEK			NEW RIVER						
		DP1 Length (cm)	Mass Weight	Average Weight (g)	FS01 Length (cm)	FS02 Length (cm)	Weight (g)	FS03 Length (cm)	Weight (g)	FS04 Length (cm)	Total Weight	Average Weight (g)	FS05 Length (cm)	Weight (g)
Atlantic Menhaden	2B-NR-FS04										17.2	13		
											18.8	60		
											18.5	60		
											18.5	60		
											17.5	60		
											17	60		
											15.5	40		
											15.5	60		
											17	60		
											17.5	50		
											18	6		
											17.5	55		
											17	50		
											19	60		
											18	55		
											17	50		
											17.5	50		
											16.7	55		
											17.4	50		
											18.5	60		
											17.5	50		
											18.5	80		
											17	50		
											19.5	60		
											15.5	50		
											18.5	50		
											18.5	50		
											17.5	50		
											18.5	80		
											17	50		
											19.5	60		
											15.5	50		
											18.5	50		
											18.5	50		
											17.5	55		
											17.7	65		
											17.5	50		
											18.5	50		
											16.5	50		
											17	47		
											18	60		
											18	60		
											17.5	60		
											16	50		
											16	50		
											17.5	50		
											17.3	60		
											18	70		
											18	70		
											18.5	70		
											17.5	60		
											18.5	60		
											16	50		
											16	50		
											17.5	55		
											16	55		
											19.5	70		
											18	60		
											18.5	60		
											18	70		
											18.5	60		
											18.5	70		
											16.5	60		
											16	50		
											17.5	60		
											16.2	50		
											16.5	50		
											17	60		
											18	60		
											18.5	60		
											16.5	60		
											16.7	60		
											16.5	50		
											17.5	60		
											16.2	50		
											17.5	60		
											19	70		
											18	60		
											17.5	60		
											17.5	60		
											17.5	60		
											16.5	50		
											18.5	70		
											16.7	60		
											15	50		
											16.5	60		
											18	60		
											16.5	60		
											16.5	60		
											18	60		
											16.5	60		
											18.5	60		
											18.5	60		
											18.2	60		
											17	60		
											16.5	50		
											17.5	60		
											17	60		
											18.5	60		

SPECIES	COC SAMPLE NO.	DRIE POND			COGDELS CREEK				NEW RIVER						
		DPI Length (cm)	Mass Weight (g)	Average Weight (g)	FS01 Length (cm)	Weight (g)	FS02 Length (cm)	Weight (g)	FS03 Length (cm)	Weight (g)	FS04 Length (cm)	Total Weight (g)	Average Weight (g)	FS05 Length (cm)	Weight (g)
Atlantic Menhaden												17.2	80		
												17	60		
												17	60		
												19	70		
												17.5	80		
												17.5	60		
												14	40		
												18	60		
												20.5	70		
												16	60		
												17.5	60		
												16.5	60		
												18	60		
												16.3	60		
												15.5	50		
												17.1	60		
												16.5	60		
												18	60		
												17	60		
												18	60		
												15.5	50		
												17.5	80		
												18.5	60		
												18	60		
												17.5	60		
												18.5	60		
												17	60		
												19.5	70		
												16	60		
												17.5	60		
												18	60		
												17.5	60		
												15.5	50		
												17.5	60		
												16.5	60		
												18.5	60		
												18.5	60		
												16.5	50		
												17	60		
												19	70		
												17.5	60		
												18.5	60		
												17	60		
												17.5	60		
												17.5	60		
												17.5	60		
												17	60		
												17	60		
		COUNT										196	196	4	4
		AVERAGE										17.4618182	56.8732289	18.125	56.25
		MAXIMUM										20.5	80	20.5	70
		MINIMUM										14	3.5	16.5	50
Summer Flounder	28-FS04-SMRF01											28.5	225		
	28-FS05-SMRF01													35.5	540
	28-FS05-SMRF02													27.5	215
	28-FS05-SMRF03													31.5	315
	28-FS05-SMRF04													21.5	105
	28-FS05-SMRF05													27.5	210
	28-FS05-SMRF06													37.5	560
	28-FS05-SMRF07													31.5	300
	28-FS04											6	6		
	COUNT											2	2	7	7
	AVERAGE											16.25	116.5	30.3571429	323.571429
	MAXIMUM											26.5	225	37.5	560
	MINIMUM											6	6	21.5	105
Black Drum	28-FS04-BD01											25.5	255		
	COUNT											1	1		
	AVERAGE											25.5	255		
	MAXIMUM											25.5	255		
	MINIMUM											25.5	255		
Spotted Sea Trout	28-FS05-SS01													46.5	960
	COUNT													1	1
	AVERAGE													46.5	960
	MAXIMUM													46.5	960
	MINIMUM													46.5	960









Scientific Name	Orde Pond	Cogdeis Creek			New River	
	28-OP-BN01	28-BN01	28-BN02	28-BN03	28-BN04	28-BN05
NEMERTEA						
Anopla						
Heteronemertea						
Lineidae						
<i>Micrura leidyi</i>				2		
ANNELIDA						
Oligochaeta						
Lumbriculida						
Lumbriculiae						
<i>Eclipidrilus sp.</i>						
Tubificida						
Tubificidae						
<i>Limnodrilus hoffmeisteri</i>	9	7	4	3	8	
<i>Isochaetides freyi</i>						
<i>Spirosperma carolinensis</i>						
Polychaeta						
Ariciida						
Orbiniidae						
<i>Scoloplos fragilis</i>					30	3
Capitellida						
Capitellidae						
<i>Capitella capitata</i>	6	51	28	42	30	
<i>Heteromestus filiformis</i>						
Phyllodocida						
Nereidae						
<i>Nereis succinea</i>	8	12	5	15	3	
Phyllodocidae						
<i>Eteone heteropoda</i>						
Spionida						
Spionidae						
<i>Polydora sp.</i>					1	
<i>Streblospio benedicti</i>					56	3
<i>Scotocolepides viridis</i>						
Terebellida				13		
Ampharetidae						
<i>Hypaniola grayi</i>						
ARTHROPODA						
Crustacea						
Amphipoda						
Corophiidae						
<i>Corophium lacuatre</i>						
Gammaridae						
<i>Crangonyx pseudogracilis</i>						1
<i>Gammarus sp.</i>						
<i>Gammarus tigrinus</i>						
Talitridae					1	
<i>Orchestia grillus</i>						
Tanaidacea						
Tanaidae						
<i>Leptochelia rapox</i>						
Decapoda						
Palaeomonidae						
<i>Palaeomonetes pugio</i>						
Portunidae						
<i>Callinectes sp.</i>					1	
Tanaidacea						
Paratanaidae						
<i>Leptochelia savignyi</i>						1
Insecta						
Coleoptera						
Dytiscidae						
<i>Hydroporus sp.</i>	2					
Elmidae						
<i>Dubiraphis sp.</i>						

Scientific Name	Orde Pond 28-OP-BN01	Cogdels Creek		New River		
		28-BN01	28-BN02	28-BN03	28-BN04	28-BN05
Diptera						
Ceratopogonidae						
<i>Bezzia/Palpomyia</i> sp.	2					
<i>Palpomyia/sphaeromias</i> sp. (biting midges)	2	1				
Chaoboridae						
<i>Chaoborus</i> sp.						
Chironomidae						
<i>Ablabesmyia annulata</i>						
<i>Ablabesmyia ramphus</i> gr.						
<i>Ablabesmyia mallochi</i>						
<i>Chironomus decorus</i> gr.	66	26	1	3		
<i>Chironomus ochreatus</i>						
<i>Cladopelma</i> sp.	1	2				
<i>Cladotanytarsus mancus</i> gr. (midges)	1					
<i>Cinotanypus pinguis</i>			1			
<i>Cricotopus bicinctus</i> gr.						
<i>Cryptochironomus fulvus</i> gr.						
<i>Dicretendipes modestus</i>	8	3				
<i>Dicretendipes nervosus</i>						
<i>Epolcladius</i> sp.						
<i>Glyptotendipes</i> sp.						
<i>Larsia</i> sp.	27					
<i>Macropelopia</i> sp.	1					
<i>Nanocladius bicolor</i> gr.			2			
<i>Nilothauma</i> sp.						
<i>Parauterborniella nigrohalteralis</i>						
<i>Polydellium illinoense</i>						
<i>Polydellium scalaenum</i>						
<i>Procladius</i> sp.	2	7				
<i>Pseudochironomus</i> sp.	1					
<i>Thienemannimyia</i> gr.		19				
Tanytarsus sp.						
<i>Tribeles lucundum</i>						
Tipulidae				1		
<i>Antocha</i> sp.						
<i>Pseudolimnophila</i> sp.						
Ephemeroptera						
Caenidae						
<i>Caenis punctata</i>	1					
Ephemeroptera						
Ephemeridae						
<i>Hexagenia bilineata</i>						
Megaloptera						
Sialidae						
<i>Sialis</i> sp.						
Odonata						
Coenagrionidae						
<i>Argia</i> sp.						
<i>Enallagma signatum/vesperum</i>	2	1				
Libellulidae						
<i>Pechydiplex longipennis</i>						
Trichoptera						
Polycentropodidae						
<i>Phylacentropus</i> sp.						
MOLLUSCA						
Bivalvia						
Mytiloidea						
Mytilidae						
<i>Geukensia demissa</i>					26	1
Veneroida						
Mactridae					30	19
<i>Mulinia lateralis</i>						
Corbiculidae						
<i>Polymesoda caroliniana</i>						
Semelidae					6	
<i>Abra aequalis</i>						
Sphaeriidae						
<i>Pisidium casertanum</i>						
Tellinidae					30	3
<i>Tellina</i> sp.						
<i>Macoma tenta</i>						
Veneridae					5	
<i>Gemma gemma</i>						
Total Taxa	13	11	8	6	13	8
Total Specimens	123	82	85	42	251	63
Species Density	784	523	542	268	1600	402
Shannon-Weiner Diversity	0.651	0.647	0.547	0.493	0.930	0.619
Brillouin's Diversity	0.593	0.765	0.495	0.422	0.890	0.550

## Percent Individuals - Benthic Macroinvertebrate - Site 28

Scientific Name	Orde Pond	28-OP-BN01	28-BN01	28-BN02	28-BN03	28-BN04	28-BN05
NEMERTEA							
Anopla							
Heteronemertea							
Lineidae							
<i>Micrura leidyi</i>							
ANNELEIDA							
Oligochaeta							
Lumbriculida							
Lumbriculiae							
<i>Eclipsidrilus sp.</i>							
Tubificida							
Tubificidae							
<i>Limnodrilus hoffmeisteri</i>	7.32	8.54	4.71	7.14		3.19	
<i>Isochaetides freyi</i>							
<i>Spirosperma carolinensis</i>							
Polychaeta							
Aricida							
Orbiniidae							
<i>Scoloplos fragilis</i>							
Capitellida							
Capitellidae							
<i>Capitella capitata</i>		7.32	60.00	66.67	16.73	47.62	
<i>Heteromestus filiformis</i>							
Phyllodocida							
Nereidae							
<i>Nereis succinea</i>		9.76	14.12	11.90	5.98	4.76	
Phyllodocidae							
<i>Eteone heteropoda</i>							
Spionida							
Spionidae							
<i>Polydora sp.</i>						0.40	
<i>Streblospio benedicti</i>						22.31	
<i>Scolecolepides viridis</i>							4.76
Terebellida					15.29		
Ampharetidae							
<i>Hypaniola grayi</i>							
ARTHROPODA							
Crustacea							
Amphipoda							
Corophiidae							
<i>Corophium lacuatre</i>							
Gammaridae							
<i>Crangonyx pseudogracilis</i>							
<i>Gammarus sp.</i>						0.40	
<i>Gammarus tigrinus</i>							
Talitridae							
<i>Orchestia grillus</i>							
Tanaidacea							
Tanaididae							
<i>Leptochelia rapox</i>							
Decapoda							
Palaemonidae							
<i>Palaemonetes pugic</i>							
Portunidae							
<i>Callinectes sp.</i>							
Tanaidacea							
Paratanaididae							
<i>Leptochelia savignyi</i>							
Insecta							
Coleoptera							
Dytiscidae							
<i>Hydroporus sp.</i>		2.44					
Elmidae							
<i>Dubiraphis sp.</i>							

Scientific Name	Orde Pond	Cogdels Creek			New River	
	28-OP-BN01	28-BN01	28-BN02	28-BN03	28-BN04	28-BN05
Diptera						
Ceratopogonidae						
Bezzla/Palpomyia sp.	1.63					
Palpomyia/sphaeronotus sp.	1.63	1.22				
Chaoboridae						
Chaoborus sp.						
Chironomidae						
Ablabesmyia annulata						
Ablabesmyia ramphus gr.						
Ablebesmyia mallochi						
Chironomus decolor gr.						
Chironomus ochreatus						
Cladopelma sp.						
Cladotanytarsus mancus gr.						
Clinotanypus pinguis						
Cricotopus bicinctus gr.						
Cryptochironomus fulvus gr.						
Dicrotendipes modestus						
Dicrotendipes nervosus						
Epicladius sp.						
Glyptotendipes sp.						
Larsia sp.	21.95					
Macropelopia sp.	0.81					
Nanocladius bicolor gr.						
Nilothauma sp.						
Paraiauterborniella nigrohaineralis						
Polypedilum illinoense						
Polypedilum scalaenum						
Procladius sp.	1.63					
Pseudochironomus sp.	0.81					
Thienemannimyia gr						
Tanytarsus sp.						
Tribolos lucundum						
Tipulidae						
Antocha sp.						
Pseudolimnophila sp.						
Ephemeroptera						
Caenidae						
Caenis punctata						
Ephemeroptera						
Ephemeridae						
Hexagenia bilineata						
Megaloptera						
Sialidae						
Sialis sp.						
Odonata						
Coenagrionidae						
Argia sp.						
Enallagma signatum/vesperum	1.63	1.22				
Libellulidae						
Pechydiplex longipennis						
Trichoptera						
Polycentropodidae						
Phylacentropus sp.						
MOLLUSCA						
Bivalvia						
Mytilidae						
Mytilus demissus						
Venerida						
Mactridae						
Mulinia lateralis						
Corbiculidae						
Polymesoda caroliniana						
Semelidae						
Abra aequalis						
Sphaeriidae						
Pisidium casertanum						
Tellinidae						
Tellina sp.						
Macoma tenta						
Veneridae						
Gemma gemma						
Total Percent	100	100	100	100	100	100