

03.15-04/12/99-0232

FINAL

**AQUATIC ASSESSMENT
FOR STONE BAY**

**MARINE CORPS BASE
CAMP LEJEUNE, NORTH CAROLINA**

CONTRACT TASK ORDER 0100

APRIL 12, 1999

Prepared For:

**DEPARTMENT OF THE NAVY
ATLANTIC DIVISION
NAVAL FACILITIES
ENGINEERING COMMAND
*Norfolk, Virginia***

Under the:

**LANTDIV CLEAN Program
Contract N62470-95-D-6007**

Prepared By:

**BAKER ENVIRONMENTAL, INC.
*Coraopolis, Pennsylvania***

TABLE OF CONTENTS

| | <u>Page</u> |
|--|-------------|
| LIST OF ACRONYMS | iv |
| EXECUTIVE SUMMARY | ES-1 |
| 1.0 INTRODUCTION | 1-1 |
| 1.1 General Description | 1-1 |
| 1.2 Purpose | 1-1 |
| 2.0 SPECIES PROFILES | 2-1 |
| 2.1 Hard Clam (<i>Mercenaria mercenaria</i>) | 2-1 |
| 2.2 Little Black Mussel (<i>Musculus niger</i>) | 2-2 |
| 2.3 American Oyster (<i>Crassostrea virginica</i>) | 2-3 |
| 3.0 INVESTIGATION ACTIVITIES | 3-1 |
| 3.1 Previous Investigation | 3-1 |
| 3.2 Aquatic Assessment Field Investigation | 3-2 |
| 3.2.1 Reference Stations | 3-2 |
| 3.2.2 Study Area Reconnaissance | 3-2 |
| 3.2.3 Water Quality Measurements | 3-3 |
| 3.3 Sediment Sampling | 3-3 |
| 3.4 Biota Sampling | 3-4 |
| 3.5 Sample Station Surveying | 3-5 |
| 3.6 Sample Preparation | 3-5 |
| 3.7 Quality Assurance | 3-6 |
| 3.7.1 Data Quality Objectives | 3-6 |
| 4.0 AQUATIC ASSESSMENT | 4-1 |
| 4.1 Methods | 4-1 |
| 4.1.1 Comparison to Sediment Screening Values | 4-2 |
| 4.1.2 Comparison to Benthic Literature Values | 4-2 |
| 4.1.3 Comparison of Study Area Results to Reference Area Results | 4-3 |
| 4.1.4 Ecological Receptor Models | 4-3 |
| 4.1.5 Uncertainties | 4-5 |
| 4.2 Sediment Evaluation | 4-7 |
| 4.3 Benthic Tissue Evaluation | 4-8 |
| 4.3.1 Clam Tissue | 4-8 |
| 4.3.2 Mussel Tissue | 4-8 |
| 4.3.3 Oyster Tissue | 4-9 |
| 4.4 Qualitative Benthic Evaluation of Copper and Lead Concentrations | 4-9 |
| 4.5 Ecological Receptor Models | 4-9 |
| 5.0 SUMMARY OF FINDINGS | 5-1 |
| 5.1 Sediment Evaluation | 5-1 |
| 5.2 Biota Tissue Evaluation | 5-1 |
| 5.3 Ecological Receptor Models | 5-2 |
| 5.4 Conclusion | 5-2 |
| 6.0 REFERENCES | 6-1 |

TABLE OF CONTENTS
(Continued)

| | |
|------------|---|
| Table 3-1 | CH2M Hill Analytical Data - Surface Water Metal Results |
| Table 3-2 | CH2M Hill Analytical Data - Sediment Metal Results |
| Table 3-3 | CH2M Hill Water Quality Measurements |
| Table 3-4 | Water Quality Measurements |
| Table 3-5 | Positive Detections in Sediment |
| Table 3-6 | Positive Detections in Clam Tissue |
| Table 3-7 | Positive Detections in Mussel Tissue |
| Table 3-8 | Positive Detections in Oyster Tissue |
| | |
| Table 4-1 | Lowest Observed Adverse Effect Levels/No Observed Adverse Effect Levels |
| Table 4-2 | Frequency and Range of Sediment Data Compared to Sediment Screening Levels |
| Table 4-3 | Frequency and Range of Clam Tissue Data Compared to Reference Areas |
| Table 4-4 | Frequency and Range of Mussel Tissue Data Compared to Reference Areas |
| Table 4-5 | Frequency and Range of Oyster Tissue Data Compared to Reference Areas |
| Table 4-6 | Qualitative Comparison of Copper Concentrations Detected in Benthic Samples |
| Table 4-7 | Qualitative Comparison of Lead Concentrations Detected in Benthic Samples |
| Table 4-8 | Summary of Ecological Contaminants of Concern Per Media |
| Table 4-9 | Aquatic Species - Conservative Models - Study Area |
| Table 4-10 | Aquatic Species - Less Conservative Models - Study Area |
| Table 4-11 | Aquatic Species - Less Conservative Models - Reference Areas |

LIST OF FIGURES

| | |
|------------|-------------------------|
| Figure 1-1 | Site Location Map |
| Figure 3-1 | Sample Location Map |
| Figure 3-2 | Study Area Location Map |

LIST OF ATTACHMENTS

| | |
|--------------|------------------------------|
| Attachment A | Site Photographs |
| Attachment B | Analytical Results |
| Attachment C | Form I's |
| Attachment D | Analytical Quality Assurance |
| Attachment E | Ecological Information |

LIST OF ACRONYMS AND ABBREVIATIONS

| | |
|------------------|---|
| $\mu\text{g/kg}$ | microgram per kilogram |
| $\mu\text{g/L}$ | microgram per liter |
| AET | apparent effects threshold |
| Baker | Baker Environmental, Inc. |
| BSAF | biota to soil/water/sediment accumulation factor |
| cm | centimeter |
| CLP | contract laboratory program |
| DO | dissolved oxygen |
| DQO | data quality objective |
| ECOC | ecological contaminant of concern |
| EMD | Environmental Management Division |
| ER-L | effects range - low |
| ER-M | effects range - median |
| GPS | global positioning system |
| HQ | hazard quotient |
| LANTDIV | Naval Facilities Engineering Command, Atlantic Division |
| LCS | laboratory control sample |
| LD ₅₀ | mean lethal dose |
| LOAEL | lowest-observed-adverse -effect level |
| MCB | Marine Corps Base, Camp Lejeune, North Carolina |
| mg/kg | milligram per kilogram |
| mg/L | milligram per liter |
| mm | millimeter |
| NOAA | National Oceanographic and Atmospheric Administration |
| NOAEL | no-observed-adverse-effect level |
| ppm | parts per million |
| ppt | parts per thousand |
| QA | quality assurance |
| SSV | sediment screening value |
| SU | standard unit |
| TAL | target analyte list |
| TOC | total organic carbon |
| USEPA | United States Environmental Protection Agency |

EXECUTIVE SUMMARY

The New River system is an important resource to Marine Corps Base (MCB), Camp Lejeune and the general public. The estuary is used for many purposes including recreational activities and commercial fishing. MCB, Camp Lejeune takes a pro-active stance to ensure that none of the activities related to base operations cause an adverse impact to the surrounding environment.

In the interest of maintaining a healthy ecosystem, the base has completed an ecological assessment in a section of the river known as Stone Bay. The investigation was completed to address potential impact to the bay from ongoing operations at the Stone Bay Rifle Range. In particular, concerns were expressed as to the effect of spent ammunition on the aquatic environment. This study has evaluated the potential impact of copper and lead from spent ammunition at the rifle range.

Prior to initiating this detailed study, a previous investigation was completed by CH2M Hill during the summer of 1998. The CH2M Hill study entailed the collection of surface water and sediment samples which were analyzed for metals. The results of this study indicated that metal concentrations were below screening criteria established for the protection of aquatic species, suggesting that operations at the rifle range have not had an adverse impact upon ecological receptors in Stone Bay. Although the 1998 CH2M Hill study suggested that range operations had not negatively impacted the bay, it was felt that a more rigorous study be undertaken in Stone Bay to ensure that the range is not posing a risk to the aquatic environment.

Therefore, as a follow-up to the investigation completed by CH2M Hill, Baker Environmental completed the aquatic assessment presented herein. This assessment was performed using guidance established by the United States Environmental Protection Agency (USEPA) and included the collection of sediment and shellfish species. The shellfish for this study included: clams, mussels, and oysters. Sediment and shellfish were collected from locations within the safety fan portion of Stone Bay (referred to as the study area). The sediment and shellfish tissue were analyzed in the laboratory for copper and lead concentrations.

In addition to the sediment and shellfish samples collected within the study area, sediment and shellfish were collected from two areas within the bay outside the rifle range safety fan. These reference areas were selected to represent background conditions within the New River. The reference area information was used for comparative purposes to distinguish between baseline conditions within the New River system and the study area.

Sediment collected in the study area was evaluated in three ways: 1) detected copper and lead concentrations were compared with USEPA Region IV sediment screening values for the protection of benthic species, 2) detected copper and lead concentrations were compared to reference area sediment concentrations, and 3) detected copper and lead concentrations were evaluated in aquatic receptor models.

Shellfish tissue collected from the study area also was evaluated in three ways: 1) detected copper and lead concentrations were compared with concentrations detected in nationwide National Oceanic and Atmospheric Administration (NOAA) Status and Trend studies, 2) detected copper and lead concentrations were compared to reference area benthic tissue concentrations, and 3) detected copper and lead concentrations were evaluated in aquatic receptor models.

Aquatic receptor models were used in this study as another tool to assess the detected concentrations of copper and lead in the sediment and shellfish samples. These aquatic receptor models are USEPA-accepted, mathematically-generated models that represent different species of animals potentially inhabiting the area that may ingest sediment and shellfish within the bay. The great blue heron and the mink were the receptor models selected for this assessment. The models use species-specific ingestion rates in conjunction with site-specific sediment and shellfish copper and lead concentrations to determine potential risks to the specified animal.

Results from the Stone Bay aquatic assessment are summarized below. These conclusions are based upon previous data collected, site-specific data obtained in January 1999, USEPA-accepted methods, and relevant scientific literature.

- The contaminants of concern from the rifle range, copper and lead, were detected below conservative sediment screening values, indicating that the shellfish communities within the study area are not adversely impacted by rifle range activities.
- Both copper and lead were detected below reference concentrations among oyster tissue samples and only slightly greater than reference concentrations in the clam and mussel tissue samples.
- A qualitative comparison of copper and lead concentrations to literature values suggests that there is no difference in copper and lead concentrations between the study area and literature values.
- The results of the heron and mink models indicate that there are no risks present to these animals greater than the risks present in reference areas.

Results from this aquatic assessment indicate no significant differences between study area and reference area ecological conditions. An evaluation of sediment samples, biota tissue samples, and results of the ecological receptor models indicate no harmful effects from copper and lead concentrations to shellfish inhabiting Stone Bay. Models of animals eating the shellfish demonstrated a slight risk to the great blue heron in the study area as well as in the reference areas. Based upon an evaluation of all information, we cannot identify any differences between areas assumed to be unaffected by range operations and the study area. The results of this aquatic assessment indicate that no further actions are required to address the potential effects of spent ammunition on the aquatic environment of Stone Bay.

1.0 INTRODUCTION

This document presents the analytical results and findings from an aquatic assessment of Stone Bay at Marine Corps Base (MCB) Camp Lejeune, North Carolina. The report has been prepared by Baker Environmental, Inc. (Baker) for the Atlantic Division, Naval Facilities Engineering Command (LANTDIV) and MCB, Camp Lejeune.

1.1 General Description

Located in Onslow County, North Carolina, MCB Camp Lejeune is the host to six Marine Corps commands and two Navy commands. The entire facility includes approximately 236 square miles and is located within the generally flat, Atlantic Coastal Plain. As shown in Figure 1-1, MCB, Camp Lejeune is bisected by the New River which flows in a southeasterly direction and forms a large estuary before entering the Atlantic Ocean. Stone Bay is part of the large estuary formed by the New River.

1.2 Purpose

This aquatic assessment of Stone Bay has been performed to address concerns regarding the possible impact of ongoing operations at the Stone Bay Rifle Range. Specifically, that copper and lead present in spent ammunition from the rifle range, may be impacting the aquatic environment of Stone Bay. Two factors have brought about the concerns: (1) the rifle range fan, which delineates the area of projectile impact, extends into a portion of Stone Bay, and (2) the rifle range does not employ target backstops or projectile recycling; therefore, a portion of the total number of projectiles enter the bay during range operations.

The purpose of this investigation is to determine the potential impact of copper and lead in spent ammunition, on the aquatic environment. Analytical results from the sampling effort have been used to conduct a semi-quantitative ecological risk assessment on the aquatic habitat of Stone Bay. The portion of Stone Bay located within the rifle range fan, which is referred to as the "study area" throughout this report, was of primary concern during the project planning phase and during preparation of this report.

This aquatic assessment focuses upon the potential of copper and lead, to impact benthic organisms (i.e., organisms living in or on the bottom of a water body) within the study area. The target benthic organisms for this study are oysters, clams, and mussels. The target organisms were selected because of the following characteristics:

- The organisms are sedentary; therefore, they are constantly exposed to the surface water and sediment within the study area.
- The organisms represent an important intermediate trophic level in the aquatic food chain.
- Each of the target organisms are known to inhabit Stone Bay. Several planting sites managed by the state of North Carolina are located adjacent to the study area. The areas are used by commercial fisherman to harvest oysters and clams.
- Oyster, clams, and mussels are known to be sensitive to contamination.

Note: Further details of the selected species are presented in Section 2.0.

The aquatic assessment contained in this report consists of sediment and biota tissue analyses of target organisms obtained from five sampling locations within the study area and from two reference sampling locations. The reference locations are not expected to be impacted by rifle range activities. Copper and lead concentrations detected in the sediment have been compared to sediment screening values developed for the protection of benthic species. The concentration of copper and lead in sediment samples collected from the study area have also been compared with concentrations in sediment obtained from the reference stations. The biota tissue samples collected from the study area have been compared to the biota tissue samples collected from the reference stations. In addition, copper and lead tissue samples have been compared to literature residue values for concentrations that are typical nationwide.

2.0 SPECIES PROFILES

The following sections present a brief description and profile of each target organism used in the aquatic assessment of Stone Bay.

2.1 Hard Clam (*Mercenaria mercenaria*)

The hard clam, or Quohog is found from the Gulf of St. Lawrence in Canada throughout the Gulf of Mexico to Texas. Hard clams are abundant from Virginia to Massachusetts and support isolated breeding communities above Cape Cod. They occur throughout the South Atlantic in estuaries from the intertidal zone to a water depth of 15 meters or more (NCDENR, 1997).

Hard clams support an important commercial fishery along the Atlantic coast. Among the species of clams harvested in the United States, hard clams yield the highest dollar value, and are exceeded only by surf clams and ocean quohogs, in kilograms of meats harvested. The harvest value of hard clams in North Carolina increased significantly from 1971 to 1995. The sustained increase may be attributed to a rise in both price and landings over that time frame. Annual dockside value reached an all time peak in 1987, with a nominal value of approximately \$8.4 million. Expressed in constant dollars (i.e., removing the effects of inflation by using consumer price index values from 1982 - 1984), the value of hard clams rose 1,789% from 1971 to the peak in 1987, then declined 44% between 1987 and 1995. Most of the decline can be attributed to a decrease in the number of mechanical harvest fisheries and closure of many harvest areas due to red tide in 1988. With respect to gear used by clambers for harvesting, during the period 1979 - 1993, hand harvesting accounted for 69% of the total production. Prices received by fisherman vary by different sizes, or grades, of hard clams. In general, the average price for hard clams has increased from 1 cent per clam in 1971 to 13 cents per clam in 1995 (NCDENR, 1997).

Hard clams live in the substrate with the long shell axis 25°-45° from vertical. The average depth at which clams live is 2 centimeter (cm) in sand and 1 cm in mud; smaller clams burrow deeper than large clams. Horizontal movement of adult clams is limited and the distance traveled is generally correlated with clam size, smaller clams being more active (Eversole, 1987).

Suspension feeding bivalves, such as the hard clam, obtain food by filtering suspended particulate matter and absorbing dissolved organics from the water. Water enters through the ventral inhalant siphon, passes through the gills to an exhalent cavity and out the dorsal exhalent siphon. Food particles suspended on the inhalant surface of the gills are sorted and passed to the gill edges and moved to the anterior of the labial palps (Eversole, 1987).

Crabs appear to be the major predators of the hard clam in the South Atlantic region. The blue crab is probably the most destructive predator among crabs; mud crabs and stone crabs prey less on hard clams (Eversole, 1987).

Temperature has been considered the most important environmental requirement in determining the time of spawning, because a certain degree of gonad ripeness or maturation must be attained before hard clams can respond to specific spawning stimuli. The hard clam has been found growing in waters of 4 parts per thousand (ppt) to over 35 ppt salinity, but growth is optimal at 24 - 28 ppt. Native clam beds are known to occur at salinities of 10 - 28 ppt in North Carolina. Dissolved oxygen concentrations of 6.8 - 7.4 milligrams per liter (mg/L) are recommended for successful culture of the hard clam and are critical to the larval life stages. Adult hard clams encounter a wide range of dissolved oxygen (DO) concentrations and have evolved several metabolic mechanisms to handle such conditions. The hard clam usually lives in well-buffered areas; however, pH may decrease below 7.0 Standard Units (S.U.) in tide pools and estuaries with poor circulation, heavy siltation, pollution, and hydrogen sulfide production. Substrate type and the degree of sorting appears to be an important factor influencing the setting of hard clam larvae. It has been observed in the laboratory that hard clams prefer to set in sand rather than in mud. Adult hard clams occur most frequently in sandy bottoms with shells (Eversole, 1987).

2.2 Little Black Mussel (*Musculus niger*)

Mussels have a wide range of habitat and are found from the Arctic Ocean to North Carolina. They are most diverse in eastern North America. They spend their entire life partially or wholly buried in mud, sand, or gravel in permanent bodies of water. Mussels prefer salinity ranging from 0 ppt to 35 ppt. The vast majority of species are found in streams, but a few are present in ponds or lakes. Although they can be found in almost any type of stream bottom, mussels are usually absent from, or rare in, areas of shifting sand or deep silt (FMM, 1999). The shells of mussels are thin and oval

shaped. The beak of the mussel is located close to the front end. They have rather prominent radiating lines at both ends with a relatively smooth area at the center of each valve. Mussels typically are a deep brownish black color with a rusty brown peristracum and a pearly white interior. The mussel moves from place to place using its foot as a prehensile organ and spinning a new byssus (mass of filament used for attachment) when a satisfactory situation has been found (Morris, 1973). Mussels attach to plant stems, rhizomes, stones, or shells by means of their byssus (White C.P., 1997).

Mussels continuously pump water through their bodies filtering food from the incoming tide. The food consists of detritus, which is organic matter found on the stream or lake bottom, and plankton, composed of microscopic plants and animals suspended in the water. Water enters via the incurrent or branchial siphon and exits via the excurrent or anal siphon (FMM, 1999).

Mussels are long-lived, with many species living more than 10 years and some reported to live more than 100 years. In many species, the surface of the shell has distinct black lines or ridges, which are believed to represent winter rest periods. The rest periods, or growth rings, are often used to estimate the age of a mussel. Mussels are an important food source for many animals, including muskrats, minks, otters, fishes, and some birds (FMM, 1999).

Mussels are one of the most endangered groups of animals in North America. Surveys conducted over the past few decades have documented significant declines in mussel populations across North America. Among the factors thought to be responsible for the declines are over harvest; siltation of habitat from agriculture, poor land management, channelization, and impoundments; competition from exotic species such as the zebra mussel; and pollution by herbicides, pesticides, and other chemicals (FMM, 1999).

2.3 American Oyster (*Crassostrea virginica*)

The American oyster, also referred to as the eastern oyster or the common oyster, plays a valuable role in the estuaries of North Carolina because its colonization of bottom lands creates a productive habitat, and the animal itself is harvested as a food item. The commercial oyster fishery is one of the most valuable seafood industries in the nation. Oyster production in 1991 was valued at \$98 million, which represents about 3% of the \$3.3 billion dockside value of the U.S. commercial

seafood industry. Oyster production in North Carolina during 1991 (\$1.2 million) comprised 1.8% of the total state commercial edible seafood production of \$66.8 million. The average price per pound of dockside oyster meat harvested in North Carolina during 1991 was \$3.35; above the national average of \$3.08. Nationally, oysters are among the top ten species in annual harvest value, as well as in price per pound (NCDENR, 1995).

The American oyster may be found in coastal areas from the Gulf of St. Lawrence in Canada throughout along the eastern seaboard. The American oyster may also be found in the Gulf of Mexico, the Bay of Campeche, Mexico and throughout the West Indies. Optimum salinity range for the species falls between 10 and 28 ppt, although oysters may be found in salinities as low as 5 ppt and as high as 32 ppt. Salinities of less than 10 to 12 ppt can prevent larval setting even though adult oysters may continue to exist. Low levels of DO may also cause mortality of set oysters. Adult oysters can survive for several days when DO concentrations are less than 1.0 mg/L, but survival times vary inversely with temperature. Although water temperature may affect larval development and is important in the annual growth and development of parasites, it only directly affects oyster stocks in extreme cases. Oysters can tolerate ambient water temperatures from 1° to 36° C (NCDENR, 1997).

Oysters are dioecious, (having male reproductive organs in one individual and female in another) but have the ability to change sexes once each year. Formation of eggs and sperm is stimulated by increasing water temperatures during spring. Fertilized eggs develop through trochophore and veliger larval stages over a period of two to three weeks. Larvae can migrate vertically in the water column and may be able to maintain their position in the estuary by avoiding certain temperature or salinity changes. Oyster larvae have been known to travel at least 30 miles. Dispersion of the larvae is largely dependent upon prevailing currents and flushing rates of estuaries. As the larval stage ends, oysters must locate a suitable attachment point or perish. Oyster growth is highest during the first six months after setting and gradually declines throughout the life of the oyster (NCDENR, 1997).

Gastropods, primarily oyster drills, are among the most destructive oyster predators. Another predator, blue crabs can readily consume up to 19 oysters per day. Of the fish that are known to feed on oysters, perhaps the most impressive is the black drum. Oysters up to 112 millimeters (mm) in length have been consumed by large drum (i.e., drum over 90+ cm in length). Other fish that

consume oysters, include Atlantic croaker, spot, toad fish, and sheepshead. In addition the cownose ray has been found to prey on oysters as well (NCDENR, 1997).

Petroleum products, heavy metals, pesticides, chlorine, and detergents can negatively impact oyster populations. The increased use of these organic compounds and metals in and around suitable estuaries has been shown to adversely impact oysters (NCDENR, 1997).

The most critical habitat areas for oyster populations are the oyster beds or rocks which form by the accumulation of oyster shells over the course of many years. Significant concentrations of oysters can also be found on outcropping of fossil shell beds, hard clam and bay scallop shells also on exposed roots. Rock jetties, sea walls, and pilings also contribute to oyster habitat (NCDENR, 1995).

3.0 INVESTIGATION ACTIVITIES

This section presents analytical results and findings from investigations conducted at Stone Bay by CH2M Hill during 1998 and Baker 1999. Findings from this aquatic assessment are based upon the field investigation conducted by Baker in January 1999. The section that follows defines reference stations used for comparison in the study and identifies the individual tasks completed as part of the January 1999 field investigation at Stone Bay. In general, the study utilizes biota (i.e., benthic species) and sediment samples collected from within the rifle range fan area of Stone Bay. Biota samples were collected, identified, measured, weighed, and recorded during the field investigation. Field photographs are included in Attachment A as additional site information. Each sample station was field surveyed using Global Positioning System (GPS) technology. Sediment samples were collected from approximately the same depths and locations as the benthic samples. Sediment analytical results were used to correlate the concentrations of metals in the sediment to corresponding tissue analytical results.

The subsections that follow provide details regarding the sampling strategy, established criteria, and quality control procedures.

3.1 Previous Investigation

During the summer of 1998, surface water and sediment samples were collected in the Stone Bay area of the New River at MCB, Camp Lejeune (CH2M Hill, 1998). This investigation was conducted as part of a baseline evaluation of the New River in support of the construction of the new advanced wastewater treatment facilities at the base. Surface water and sediment samples from five locations in Stone Bay were collected and analyzed for metals. Surface water samples were analyzed for total and dissolved metals and sediment samples were analyzed for total metals, total organic carbon (TOC), percent solids, acid volatile sulfide, and grain size. The analytical results of copper and lead detected in the surface water are presented in Table 3-1 and sediment is presented in Table 3-2. A complete report of the analytical results are presented in the CH2M Hill letter report contained within in Attachment B. Water quality parameters, which are provided in Table 3-3, were also collected at the time of this study. The locations of the sampling points are depicted in Figure 3-1.

The results of the initial study were compared to sediment screening values developed for the protection of aquatic species. As shown on Table 3-2, in each case, concentrations observed in the sediment were below the established screening criteria, suggesting that the operations at the rifle range have not had a significant impact upon ecological receptors in Stone Bay. As a follow-up to the preliminary sediment and surface water investigation, this investigation includes a tissue study of benthic species to provide a more thorough evaluation of the potential impact of copper and lead from the rifle range on aquatic receptors and the habitat of Stone Bay.

3.2 Aquatic Assessment Field Investigation

The subsections that follow describe the field investigation activities conducted during January 1999 as part of the Aquatic Assessment of Stone Bay.

3.2.1 Reference Stations

As a part of the January 1999 investigation, two areas were used as reference stations. The reference stations are areas that are ecologically similar to the study area (i.e., habitat, species potentially present, salinity, substrate type), but that are most likely not impacted by rifle range activities. The reference stations provide information regarding naturally occurring metals and the existence of any regional metal contamination, independent of the rifle range. The locations of the reference samples are shown in Figure 3-1.

Samples obtained from the reference stations were used for a qualitative comparison of the analytical data obtained from the study area to determine significant differences in the sediment and biota tissue between the study area and the ecologically similar reference area.

3.2.2 Study Area Reconnaissance

Prior to commencement of sampling activities in January 1999, the study area was reviewed with range personnel and Environmental Management Division (EMD) personnel to discuss general operations and proposed sampling methods and locations. Additional topics of discussion included, time of work on-site, site access points, verification that the reference stations selected were appropriate for this study, and formulation of a general overview of the surrounding habitat. During

the preliminary activities, the exact benthic species to be collected were determined. The target species were oysters, clams, and mussels. Several organisms were collected and examined to determine the size range available.

3.2.3 Water Quality Measurements

Prior to the collection of biota and sediment samples, depth of water and water quality parameters were measured. At each sampling station, surface water was measured for pH, specific conductance, temperature, salinity, turbidity and dissolved oxygen. All readings were measured in-situ by submerging a probe to the appropriate depth. The measurements were recorded on field data sheets during site operations and later tabulated. The results of these measurements are provided in Table 3-4.

3.3 Sediment Sampling

Sediment samples were collected from approximately the same depths and locations as biota samples. In general, the sediment samples were collected from a depth of approximately zero to six inches below the surface of the sediment.

A total of seven sediment samples was collected during the investigation. Five of the samples were collected within the study area of Stone Bay and two samples were collected from the reference locations. Figure 3-1 depicts the locations of each of the sediment samples. Figure 3-2 provides a detailed illustration of the study area and sampling locations. Each sediment sample was visually classified in the field to determine general soil type. Each of the sediment samples was analyzed for Target Analyte List (TAL) metals. Sediment characteristic analyses (i.e., grain size and TOC) were not performed on the samples obtained within the study area because similar information was obtained during the 1998 sampling event (CH2M Hill, 1998). Sediment obtained from the two reference stations were analyzed for grain size, Atterberg limits, and TOC to ensure that similar substrate conditions had been utilized. Substrate conditions at the reference stations are similar to what was observed in the study area, however the sediment sample collected from reference station RF-SD02 exhibited some clay. Results of the copper and lead sediment analyses are provided on Table 3-5. In general, the substrate material within the study area is comprised of varying amounts of silt and sand. The bottom material is comprised of mostly sand in the central portion of the study

area and in the area near sediment sample SB-SD01. More fine material was observed near station SB-SD02 near the entrance of Stone Creek.

Each sediment sample was collected using a stainless steel sediment corer with a dedicated acetate sleeve. The samples were collected by manually pushing the sediment corer into the river bed and extracting an appropriate volume of sediment. The sample was transferred directly from the acetate sleeve of the sediment corer to a laboratory-prepared glass container.

3.4 **Biota Sampling**

The locations of the biota sampling stations were based upon the availability of benthic species in the area. Because the benthic species collected were shellfish (oysters, clams, and mussels), the sampling stations coincided with locations of shellfish beds to ensure an adequate sample volume was obtained.

A total of 14 biota composite samples was collected during this investigation, comprised of different biota species from seven sample locations. Five of the samples were collected from within the study area of Stone Bay and two samples were obtained at the reference locations. Each biota composite sample consisted of several individual organisms to meet the weight requirements of the laboratory analytical procedures. Figures 3-1 and 3-2 depict the locations of each biota sample. As shown in Figure 3-2, shellfish were not present in the central portion of the study area. The substrate material observed in this portion of the study area was entirely comprised of a thick layer of small sticks, branches, and roots. It is assumed that this layer of organic debris may inhibit the species from populating this area. It was noted by the commercial fisherman hired to assist during the biota collection, that the debris observed in the central portion of Stone Bay may be due to the fact that fishermen typically do not harvest shellfish in this area due to range operations. During normal shellfish harvesting, the methods used to collect species frequently remove debris from the bottom.

The collection of biota samples was conducted with the assistance of a commercial shell fisherman who is familiar with the New River. Biota samples were obtained using boat-mounted rakes, tongs, and grab samplers. The biota samples were collected in accordance with Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories Volume I Fish and Sampling Analyses (USEPA, 1993).

Each of the biota samples was analyzed for TAL metals and percent lipids. Table 3-6 presents the results of clam tissue copper and lead analyses and Tables 3-7 and 3-8 provide a summary of copper and lead analytical results for mussel and oyster tissue samples, respectively. Results are presented for both the study area and the designated reference stations. A complete listing of the analytical results is provided in Attachment B. Form I's associated with the analytical results are contained in Attachment C.

3.5 Sample Station Surveying

Each sampling station was surveyed using a global positioning system. Spatial data were collected using code signals from satellites and then were differentially corrected with exact time interval data from a known base station. The resulting data yields point accuracies within the submeter (i.e., less than 40 to 75 centimeters) range. Upon differential correction, spatial data were exported into existing data files to produce the appropriate figures.

3.6 Sample Preparation

Sediment samples were taken directly from the sediment corer device and placed into laboratory prepared sample jars. Each jar was properly labeled and sealed and the samples were kept on wet ice prior to and during shipment to the analytical laboratory.

Biota samples were analyzed via a tissue composite method. Individual organisms were composited to acquire 20 to 30 grams of tissue sample for metal analyses. Each benthic organism collected was measured and weighed individually. The exterior shells of each benthic organism were scrubbed and rinsed with deionized water to remove the sediment and prevent possible cross contamination. The organisms intended for each composite sample were placed in a labeled, reclosable, freezer bags with the shells left intact. The samples were shipped on ice to the analytical laboratory. Upon receipt, the benthic samples were shucked and composite samples were formulated prior to chemical analysis of the tissue samples.

3.7 Quality Assurance

The purpose of Quality Assurance (QA) is to establish internal means for data generation and review to ensure that the work performed is completed at the highest professional standard. The objectives of the QA program include the following items:

- To generate data in accordance with procedures appropriate for the intended data use.
- To obtain data of sufficient quality to meet reasonable scientific scrutiny.
- To obtain data of acceptable precision, accuracy, completeness, representativeness, and comparability as required by the project.

The fundamental mechanisms that were employed to achieve the quality goals can be categorized as prevention, assessment, and correction where:

- Prevention of errors occurs through planning, following documented instructions and procedures, and careful selection of trained personnel.
- Assessment of all QA sampling reports furnished by the laboratory.
- Correction of noted conditions adverse to data quality.

3.7.1 Data Quality Objectives

Data quality objectives (DQOs) are qualitative or quantitative statements developed by the data users to specify the quality of data needed from a particular data collection activity. The DQOs are expressed in terms of precision, accuracy, representativeness, completeness, comparability and uncertainty; which are defined as follows:

- Precision - A measure of mutual agreement among individual measurements of the same property, usually prescribed similar conditions. Precision is usually expressed in terms of

the standard deviation, however, various measures of precision exist depending upon the prescribed conditions.

- Accuracy - The degree of agreement of a measurement or an average of measurements, X , with an accepted reference or true value, T , expressed as the difference between the two values, $X-T$. Accuracy is a measure of the bias in a system.
- Representativeness - Expresses the degree to which data accurately and precisely represent a characteristic of a population, parameter variations at a sampling point, a process condition, or an environmental concern.
- Completeness - A measure of the amount of the valid data obtained from the measurement system compared to the amount that was expected under normal conditions.
- Comparability - Expresses the confidence with which one data set can be compared with another.
- Uncertainty - The likelihood of all types of errors associated with a particular decision.

DQOs are intended to help develop sampling and analytical strategies designed to support the objectives of this assessment. DQOs define the level of certainty in the data that is acceptable for this assessment. The variables associated with sampling and analysis contribute to some level of uncertainty in any data generated. The objectives of this study included keeping the total uncertainty within an acceptable range. To achieve this objective, specific data quality requirements such as detection limits have to be specified. The expected detection limits of media were provided to the laboratory to ensure this requirement was met.

The data collected during this assessment was used to assess the following items:

- Identify the presence or absence of metals based upon the samples collected.
- Assess potential bioaccumulation of metals in aquatic receptors.

- Determine the potential impacts, if any, to the aquatic environment from metals that may be the result of ongoing range operations.

The DQOs for the aquatic assessment of Stone Bay have been met through several methods. Sediment samples collected within the study area and reference stations were analyzed using standard Contract Laboratory Procedures (CLP) typically used for environmental samples collected at MCB, Camp Lejeune. The quality control standards for trace metals undergoing CLP analyses are provided in Attachment D. Biota sample analyses employed the use of a Standard Reference Material (SRM), a method commonly used by marine scientists in the analysis of tissue samples. This is a proven method prepared by the National Institute of Standards and Technology (NIST) and is described in the Certificate of Analysis (SRM 1566a for Oyster Tissue) provided in Attachment D. The SRM is used for calibrating instrumentation and validating methods for the chemical analysis of marine bivalve tissue. The SRM gives acceptance ranges of elemental concentrations. In some cases, these acceptance ranges may not be met using the standard CLP type digestion and analysis. In such cases, the analysis includes a Laboratory Control Sample (LCS) to establish the remaining acceptance ranges and percent recoveries. The SRM and LCS will be used to ensure proper digestion procedures, analyses, and reporting of the tissue sample results.

4.0 AQUATIC ASSESSMENT

The section that follows provides a qualitative and quantitative evaluation of aquatic samples collected from within the study area. This assessment provides risk-based conclusions that address whether ecological risks to the aquatic environment are the result of copper and lead detected in the sample media. The methodology used in the assessment is provided first (Section 4.1), followed by the actual results of the applied methods (Sections 4.2 through 4.5).

4.1 Methods

The methodologies used in this evaluation mirror the procedures outlined in the Ecological Risk Assessment Guidance for Superfund (USEPA, 1997) and the Guidelines for Ecological Risk Assessment (USEPA, 1998). This aquatic assessment was conducted using a qualitative and quantitative analysis of sediment, clam tissue, mussel tissue, and oyster tissue collected from Stone Bay within the study area. The aquatic environment was evaluated using the following methods:

- Comparison of the study area sediment concentrations to Region IV sediment screening values (SSVs).
- Comparison of study area sediment concentrations to reference area sediment concentrations.
- Comparison of study area tissue concentrations to literature values for nationwide concentrations detected in shellfish tissue.
- Comparison of study area tissue concentrations to reference area tissue concentrations.
- Calculation of aquatic species receptor models for species potentially inhabiting Stone Bay that may ingest surface water, sediment and shellfish from the study area.

The following information provides a detailed description of the methods used to assess the aquatic environment.

4.1.1 Comparison to Sediment Screening Values

Sediment values were used to select ecological contaminants of concern (ECOC) in samples collected from the study area. Concentrations detected above an SSV were retained as ECOCs in this assessment. The SSVs used in this assessment were obtained from the Supplemental Guidance to RAGS: Region 4 Bulletins - Ecological Risk Assessment (USEPA, 1995). The sediment values presented in this document were derived from statistical interpretation of effects reported in literature for direct toxicity. The values were derived from marine environment studies; however, freshwater environment studies may also have been used. In addition to Region IV sediment values, effects-range low (ER-L) and Effects-range median (ER-M) values (Long et al., 1995) were used to assess the sediment collected from Stone Bay.

Concentrations detected below the ER-L/SSV represent a minimal effects range (i.e., effects that would rarely be observed). Concentrations above the ER-L/SSV, but below the ER-M represent a possible effects range (i.e., effects that would occasionally be observed). Concentrations detected above the ER-M present a probable-effects range (i.e., effects that would frequently be observed). Hazard quotients (HQs) were calculated for each detected inorganic. The HQs represent the magnitude by which a contaminant exceeds an SSV. The HQs for this assessment were calculated by dividing the maximum detected sediment concentration by the lowest SSV available for that metal. An HQ calculated above one represents a potential risk to the aquatic environment from concentrations of that contaminant.

4.1.2 Comparison to Benthic Literature Values

Literature values established for copper and lead concentrations in benthic tissue were used to qualitatively assess the concentrations detected in the benthic tissue obtained from the study area. The literature values represent body burden residues detected in benthic species that have been demonstrated to impact the health of the organism itself (Irwin, 1997a/1997b). Maximum and mean concentrations found in biota tissue collected as a part of the National Oceanic and Atmospheric Administration (NOAA) Status and Trends Studies (1990) were used for comparative purposes.

4.1.3 Comparison of Study Area Results to Reference Area Results

The ranges of detected concentrations in the sediment, clam, mussel, and oyster samples collected from the study area were directly compared to the range of detected concentrations in corresponding samples collected from the reference areas. Two reference locations were sampled during this investigation; however, not every shellfish species was found at each proposed location (See Section 3.4).

A comparison of copper and lead concentrations found within the study area to reference concentrations was used to select the ECOCs for the benthic organisms. Reference concentrations were used as selection criteria because there are no specific screening values established for the protection of the target organisms.

4.1.4 Ecological Receptor Models

Ecological receptor models were used to evaluate potential risks to higher trophic levels in the aquatic food chain. Potential risks posed to prey species from ingestion of surface water, sediment, and benthic species from within the study area were evaluated in the models. Sediment and benthic analytical results obtained during the field investigation and surface water analytical results from the previous CH2M Hill study (see Section 3.1) were used as input values for the receptor models. Two species were selected for modeling: the great blue heron and the mink. A summary of life history information for the modeled species is presented in Attachment E.

Several different versions of the receptor models were calculated. The differences in the versions reflect the conservatism incorporated within the models. The most conservative models used the maximum detected concentrations compared with toxicity dose concentrations found to have no adverse effects. The comparative toxicity dose concentrations are referred to as no-observed adverse-effects levels (NOAELs). The least conservative models used arithmetic means of the detected concentrations compared to toxicity dose concentrations found to have the least observed effects to the species or a similar species. The comparative least effects toxicity doses are referred to as lowest-observed-adverse effects levels (LOAELs). The comparative NOAEL and LOAEL for copper and lead concentrations used in the receptor models calculated for this assessment are

presented on Table 4-1. A summary of the studies from which these numbers are based on is presented in Attachment E.

The ecological receptor models and the assumptions made within the models are presented in the sections that follow.

4.1.4.1 Receptor Model Hazard Quotients

The HQ method was used to estimate potential risks to ecological receptors within the study area. This method compares exposure concentrations with ecological endpoints such as reproductive failure or reduced growth. The following equation was used to calculate HQs:

$$\text{Hazard Quotient} = \frac{\text{Maximum Exposure} / \text{Mean Exposure Concentration}}{\text{NOAEL} / \text{LOAEL}}$$

Where:

| | | |
|--------------------------------|---|--|
| Mean Exposure Concentration | = | Arithmetic Mean Concentration Calculated |
| Maximum Exposure Concentration | = | Maximum Concentration Detected |
| NOAEL | = | No Observed Adverse Effect Level |
| LOAEL | = | Lowest Observed Adverse Effect Level |

An HQ equal to or greater than one indicates that exposure to the particular metal has the potential to cause adverse effects to the species. An HQ less than one indicates that the metal is not expected to cause adverse effects to the species. The greater the HQ, the greater the magnitude of potential risk to the species; however, for this assessment, any HQ greater than one was evaluated as a potential risk.

4.1.4.2 Receptor Model Assumptions

This aquatic assessment evaluates exposure to contaminants through food, water, and incidental ingestion of sediment. The following assumptions were made during preparation of the aquatic models calculated for this study:

- Maximum concentrations and arithmetic mean concentrations were used to represent site-wide concentrations in the receptor model calculations.
- A biota to soil/water/sediment accumulation factor (BSAF) of 1 was assumed for the vegetation, invertebrates, fish, and small mammals.
- Copper and lead were assumed to be 100 percent bioavailable.
- Because toxicity values could not be found for the specific receptor species, values reported for closely related species were used.
- If chronic NOAEL values were not available for copper and lead, LOAEL values were used. A factor of 10 was used to convert reported LOAEL values to NOAEL values. If several toxicity values were reported for a receptor species, the most conservative value was used in the risk calculations regardless of the toxic mechanism. Toxicity values obtained from long-term feeding studies were preferable to those obtained from single dose oral studies.
- Some doses were originally reported as part per million contaminants in a diet. These were converted to daily intakes (in units of mg/kg-day) by using the following formula:

$$\text{Daily Intake (mg/kg-day)} = \text{ECOC Dose (mg/kg diet)} \times \text{Ingestion Rate (kg/day)} \times 1/\text{Body Weight (kg)}$$

Dietary toxicity levels for species were converted to a daily dose based on body weight. For the ecological assessment, incidental sediment ingestion was also included in the calculation to determine the total daily intake for the receptor species. This daily dose was then used to evaluate the risk to other species if no specific toxicity data were available for a target receptor.

4.1.5 Uncertainties

As with any such ecological assessment, this investigation of Stone Bay is subject to uncertainties. Uncertainty exists in several steps of the process including: correlation of tissue concentrations to

adverse effects to species, study of lead in shellfish, use of screening levels, and the use of ecological receptor models.

4.1.5.1 Correlation of Tissue Concentrations to Adverse Effects in Species

There is uncertainty associated with correlating tissue concentrations to adverse effects on benthic species. Tissue concentrations do not infer adverse effects; however, tissue data has been used for qualitative evaluation of copper and lead.

4.1.5.2 Study of Lead in Shellfish

Uncertainty is associated with the study of rifle range contaminants of concern, specifically lead. Lead, when taken into living organisms, reacts similar to calcium and will most likely mineralize in bones, or in this case, the shells of the organism. However, it is recognized that organisms from polluted areas can build up substantial concentrations of lead in muscle tissue.

4.1.5.3 Screening Levels

Potential adverse impacts to aquatic receptors from contaminants in the sediments were evaluated by comparing sediment concentrations to SSVs. These SSVs have uncertainty associated with them because the procedures for developing them are not as established as those used in developing water screening values. In addition, sediment type (pH, acid volatile sulfide, total organic carbon) also has a significant impact on the bioavailability and toxicity of contaminants. The SSVs were developed using data obtained from freshwater, tidal freshwater, and marine environments. Therefore, their applicability in evaluating potential effects to aquatic organisms from contaminants in marine habitats introduces uncertainty due to differences in the toxicity of individual contaminants to freshwater and saltwater organisms and the bioavailability of contaminants in the two aquatic systems.

4.1.5.4 Ecological Receptor Models

There are some differences of opinion found in the literature as to the effectiveness of using ecological receptor models to predict concentrations of contaminants found in ecological species.

The food chain models currently used incorporate simplistic assumptions that may not represent conditions at the site, bioavailability of contaminants, or site-specific behavior of the receptors.

In some instances, NOAEL values were not found in the literature. If NOAEL values were not reported, then LOAEL values were used to calculate a NOAEL. A LOAEL was divided by a factor of ten to obtain NOAEL values. There is uncertainty in this calculation of NOAELs; however, the uncertainty most likely errs on the conservative side.

Doses in toxicological studies are typically reported in units of mg of contaminant/kg diet, or in units of mg contaminant/kg body weight/day. All doses reported as mg/kg in diet were converted to units of mg/kg-body weight/day. If body weights were reported for the test animals in a given study, these values were used for making this conversion. Otherwise, the body weight and ingestion rate for the species reported in other literature sources were used.

There is uncertainty associated with some of the toxicity values derived from a single species. Prediction of ecosystem effects from laboratory studies is difficult. Laboratory studies cannot take into account the effects of environmental factors which may add to the effects of contaminant stress. NOAELs were generally selected from studies using single contaminant exposure scenarios.

There is uncertainty in the total daily intake models used to evaluate a reduction of receptor populations or sub-populations. Many input parameters are based on default values (i.e., ingestion rates) that may or may not adequately represent the actual values of the parameters. In addition, there is uncertainty in the level to which the indicator species will represent other species potentially exposed to copper and lead concentrations at the site.

4.2 Sediment Evaluation

As presented in Table 4-2, sediment ECOCs within the study area were identified by a comparison of detected concentrations to SSVs. If a concentration exceeded an SSV, the inorganic was retained as a sediment ECOC. Reference area sediment concentrations are also presented in Table 4-2 for comparative purposes.

Study area sediment concentrations of copper and lead were detected below SSVs, indicating no potential risks to aquatic receptors from sediment concentrations. Therefore, no ECOCs were identified in the sediment collected from the study area. Sediment concentrations of copper and lead detected in the CH2M Hill study (see Section 3.1) were similar to concentrations detected during this investigation.

4.3 Benthic Tissue Evaluation

Three species were evaluated for this aquatic assessment: the hard clam, little black mussel, and the American oyster. Life history information for these species is provided in the profiles presented in Section 2.0. As discussed in Section 3.4, two species per sampling station were proposed for analysis. However, due to conditions in the field, two of the same species were not available from every location within the study and reference areas. Two species were collected from every sampling station; however, the two species are not the same at every station. Benthic tissue from the study area was analyzed by comparison to reference area tissue concentrations.

Tissue concentrations detected among clam, mussel, and oyster samples were compared to reference concentrations. The following sections present the ECOCs identified in each of the species based upon the reference comparison.

4.3.1 Clam Tissue

Table 4-3 presents copper and lead detected in the clam tissue collected from the study area and the ECOCs selected. Copper and lead were detected slightly above reference area tissue concentrations and retained as clam ECOCs. It is noted that the clam tissue evaluation is based on one composite sample collected from the study area. A clam sample was only collected from one station in the study area, most likely due to the low salinity in this portion of Stone Bay (see Section 3.2.3).

4.3.2 Mussel Tissue

Table 4-4 presents copper and lead detected in the mussel tissue collected from the study area and the ECOCs selected. Copper and lead concentrations in mussel tissue were above reference station

concentrations and retained as mussel ECOCs. The mussel comparison of study area tissue to reference area tissue is based on one composite sample collected in the reference areas.

4.3.3 Oyster Tissue

Table 4-5 presents the concentrations of copper and lead detected in oyster tissue collected from the study area compared to concentrations detected in the reference areas. The concentrations from the study area were below reference concentrations; therefore, no ECOCs were identified for the oyster species in this assessment.

4.4 Qualitative Benthic Evaluation of Copper and Lead Concentrations

Tables 4-6 and 4-7 present qualitative comparisons of the copper and lead concentrations (respectively) detected in the oyster and mussel tissue versus literature values. No literature values for the clam were available. The literature values are NOAA Status and Trends Studies (1990) (Irwin, 1997a and Irwin, 1997b). Study area oyster concentrations were below literature copper values. Study area mussel concentrations only slightly exceed literature copper values.

The study area oyster and mussel tissue lead concentrations only slightly exceed the maximum literature values. Reference oyster tissue concentrations also exceed the literature value for lead. This qualitative comparison does not show a significant difference between study area copper and lead concentrations and nationwide NOAA Status and Trends concentrations.

4.5 Ecological Receptor Models

Ecological receptor models for the heron and mink were calculated with site-specific concentrations from the study area. As discussed in Section 4.1.4, receptor models were calculated for ECOCs identified in sediment, oyster, clams, and mussels. Table 4-8 presents a summary of the ECOCs identified per sample media. Data input into the receptor models included surface water, sediment, and biota tissue. It is noted that although no ECOCs were identified in the sediment and oyster samples collected from the study area, copper and lead concentrations from these media were evaluated in the receptor models. Only the concentrations detected in one of the biota species could be used in the model. Therefore, to remain conservative, the highest ECOC concentration for the

three different benthic species was used to represent biota. The input values and receptor models are presented in Attachment E. The following paragraphs present the results of both the conservative and less conservative versions of the receptor models.

The most conservative receptor models, maximum concentrations and NOAEL values are presented in the NOAEL columns in Table 4-9. Sediment and biota concentrations of copper and lead resulted in HQs greater than one in the two receptor species. Surface water concentrations were also incorporated into the models; however, due to the low ingestion rates, surface water does not provide a significant effect to receptor risk. The highest risks (HQs greater than 10) to the receptor species were identified in the heron due to copper concentrations (HQ = 44) and in the mink due to copper (HQ = 13) and lead concentrations (HQ = 20).

The least conservative receptor models, mean concentrations and LOAEL values, are presented in the LOAEL columns in Table 4-10. The only HQ over one was calculated for copper in the heron model (HQ = 3), indicating only a slight potential for risk to the heron.

Table 4-11 presents results from the least conservative receptor models calculated using reference area sediment and tissue concentrations. Risks due to copper concentrations were higher in the reference area receptor models than the risks identified in the study area. The reference receptor models were calculated to demonstrate that areas considered to be unimpacted by the rifle range produced greater risks to the receptor species from sediment and tissue concentrations of copper and lead, indicating that the rifle range is not posing any adverse risk to the existing aquatic habitat within the bay.

5.0 SUMMARY OF FINDINGS

The purpose of this assessment was to determine the potential impact of copper and lead from the spent ammunition, upon the aquatic environment of Stone Bay. A summary of the results from the aquatic assessment presented in Section 4.0 is provided below.

5.1 Sediment Evaluation

The contaminants of concern from the rifle range, copper and lead, were detected below conservative sediment screening values, indicating that the benthic macroinvertebrate communities within the study area are not adversely impacted by rifle range activities. The detections and comparison to screening values are presented in Table 4-2 and discussed in detail in Section 4.2. These data indicate no adverse impact due to exposure to copper and lead in the study area.

5.2 Biota Tissue Evaluation

As presented on Tables 4-3 through 4-5 and discussed in Section 4.3, study area tissue concentrations versus reference area tissue concentrations suggest the following inorganics were elevated for each of the identified species:

- Hard Clam: copper and lead
- Little Black Mussel: copper and lead
- American Oyster: none

The contaminants of concern from rifle range activities were identified as copper and lead. Both copper and lead were detected below reference concentrations among oyster tissue samples and not significantly greater than reference concentrations in the clam and mussel tissue samples. In addition, the qualitative comparison of copper and lead concentrations to literature values suggests that there is no significant difference in copper and lead concentrations between the study area and literature values. A majority of the highest inorganic concentrations among mussel and clam tissue samples were detected in the sample obtained from the edge of the rifle range fan, SB-MU04-99A.

5.3 Ecological Receptor Models

Mean inorganic concentrations and LOAEL values were used to calculate the following HQs used to assess potential risks to receptor species from consuming sediment and biota within the study area. An HQ greater than one indicates a potential risk to the receptor species.

- Great Blue Heron - copper resulted in an HQ slightly above one. Reference concentrations demonstrated similar risks to the heron from concentrations of copper. The results of the heron model indicate that there are no significant risks above the risks present in reference areas.
- Mink - No risk to the mink model was demonstrated in the least conservative receptor models. It is noted that a slight risk from copper was calculated in the reference mink model.

5.4 Conclusion

This assessment was conducted to determine whether rifle range contaminants of concern (copper and lead) are adversely impacting the aquatic habitat within the identified study area within Stone Bay. This assessment was conducted in accordance with the methodologies presented in Ecological Risk Assessment Guidance for Superfund (USEPA, 1997) and the Guidelines for Ecological Risk Assessment (USEPA, 1998). Data used to assess the study area included sediment, clam tissue, mussel tissue, and oyster tissue analyzed for copper and lead concentrations. The assessment used data collected from reference areas within Stone Bay identified to be unimpacted by rifle range activities. The reference area data were used for comparative purposes to determine whether potential contamination is site-related or the result of regional conditions within the bay.

Results from this assessment indicate no significant differences between study area and reference area ecological conditions. An evaluation of sediment samples, biota tissue samples, and results of the ecological receptor models indicate no deleterious effects from the potential contaminants of concern, copper and lead, to benthic organisms inhabiting Stone Bay.

6.0 REFERENCES

CH2M Hill, October 28, 1998. Stones Bay Sediment and Water Quality Sampling Results.

CMEE. Canadian Ministry of Environment and Energy. 1993. Guidelines for the Protection and Management of Aquatic Sediment Quality in Ontario. August 1993.

Eversole, A.G., 1987. Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (South Atlantic)--Hard Clam. U.S. Fish Wildl. Serv. Biol. Rep. 82(11.75). U.S. Army Corps of Engineers, TR EL-82-4.

Fitchko, J. 1989. Criteria for Contaminated Soil/Sediment Cleanup. Pudvan Publishing Company, orthbrook, Illinois.

FMM. 1999. Freshwater Mussels of the Midwest. Accessed from Worldwide Web 2/22/99.

Irwin, Roy J. 1997a. Environmental Contaminants Encyclopedia Copper Entry. National Park Service Water Resources Divisions, Water Operations Branch. July 1, 1997.

Irwin, Roy J. 1997b. Environmental Contaminants Encyclopedia Lead Entry. National Park Service Water Resources Divisions, Water Operations Branch. July 1, 1997.

Long, Edward R., Donald D. Mac Donald, Sherri L. Smith, and Fred Calder. 1995. "Incidence of Adverse Biological Effects Within Ranges of Chemical Concentrations in Marine and Estuarine Sediment." Environmental Management. Volume 19.

Morris, P.A., A Field Guide to Shells of the Atlantic and Gulf Coasts and the West Indies. Houghton Mifflin Company Boston, 1973.

NCDENR. 1997. North Carolina Department of Environment, and Natural Resources Division of Marine Fisheries. North Carolina Fishery Management Plan Hard Clam. February, 1997.

NCDENR. 1995. North Carolina Department of Environment and Natural Resources. North Carolina Oyster Restoration and Fishery Management Plan. Compiled by Michael D. Marshall, Division of Marine Fisheries and North Carolina Blue Ribbon Advisory Council on Oysters. October, 1995.

Sullivan, J.J. Ball, E. Brick, S. Hausman, G. Pilarski, and D. Sopcich. 1985. Report for the Technical Subcommittee on Determination of Dredge Material Suitability for In-Water Disposal. Wisconsin Department of Natural Resources. Cited in: Fitchko. 1989.

Tetra Tech, Inc. 1986. Development of Sediment Quality Values for Puget Sound, Volume I. Puget Sound Dredged Disposal Analysis Report. Cited in: Fitchko. 1989.

USEPA. 1998. United States Environmental Protection Agency. Guidelines for Ecological Risk Assessment. Risk Assessment Forum, EPA/630/R-95/002B. August 1998.

USEPA. 1997. United States Environmental Protection Agency. Ecological Risk Assessment Guidance for Superfund: Process for Designing and Conducting Ecological Risk Assessments (Interim Final). Environmental Response Team, EPA/540/R-97-006. June 5, 1997.

USEPA. 1995. United States Environmental Protection Agency. Supplemental Guidance to RAGS: Region 4 Bulletins - Ecological Risk Assessment. Waste Management Division. November 1995.

USEPA. 1993. United States Environmental Protection Agency. Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories Volume I: Fish and Sampling Analyses. Office of Water.

White, C.P., Chesapeake Bay, Nature of the Estuary, A Field Guide. Tidewater Publishers. Centerville, Maryland, 1997.

Baker

Baker Environmental, Inc.

TABLES

TABLE 3-1

CH2M HILL ANALYTICAL DATA - SURFACE WATER METAL RESULTS
 STONE BAY AQUATIC ASSESSMENT, CTO-0100
 MCB, CAMP LEJEUNE, NORTH CAROLINA

| Analyte | Units | Trip Blank | | Sampling Stations | | | | | | | | | | W.Q. Stds Aquatic Life |
|---------|-------|------------|-------|-------------------|-------|------------|-------|------------|-------|------------|-------|------------|-------|---------------------------|
| | | | | CLMSWSB101 | | CLMSWSB201 | | CLMSWSB301 | | CLMSWSB401 | | CLMSWSB501 | | |
| | | Total | Diss. | Total | Diss. | Total | Diss. | Total | Diss. | Total | Diss. | Total | Diss. | |
| Copper | µg/L | 0.08 | 0.03 | 0.71 | 0.62 | 0.88 | 0.61 | 0.91 | 0.61 | 1.53 | 0.86 | 0.95 | 0.58 | 3 (AL) |
| Lead | µg/L | ND | ND | 0.22 | ND | 0.415 | ND | 0.437 | ND | 0.926 | ND | 0.641 | ND | 25 (N) |

Notes:

ND = Not Detected

AL = Values represent action levels as specified in 15A NCAC 2B.0220

N = See 15A NCAC 2B.0220 for narrative description of limits.

µg/L = microgram per liter

W.Q. Stds = Water Quality Standards

TABLE 3-2

CH2M HILL ANALYTICAL DATA - SEDIMENT METAL RESULTS
 STONE BAY AQUATIC ASSESSMENT, CTO-0100
 MCB, CAMP LEJEUNE, NORTH CAROLINA

| Analyte | Units | Sampling Stations | | | | | NOAA Guidelines | |
|---------|-------|-------------------|------------|------------|------------|------------|-----------------|------|
| | | CLMSDSB101 | CLMSDSB201 | CLMSDSB301 | CLMSDSB401 | CLMSDSB501 | ER-L | ER-M |
| Copper | mg/kg | ND | 18 | 5.7 | 15 | ND | 70 | 390 |
| Lead | mg/kg | 1.5 | 12 | 4 | 3.5 | 3.5 | 35 | 110 |

Notes:

mg/kg = milligram per kilogram

NOAA = National Oceanic and Atmospheric Administration

ER-L = Effects range - low

ER-M = Effects Range - median

ND = Not Detected

TABLE 3-3

CH2M HILL WATER QUALITY MEASUREMENTS
 STONE BAY AQUATIC ASSESSMENT, CTO-0100
 MCB, CAMP LEJEUNE, NORTH CAROLINA

| Sample ID | Depth (feet) | Position | Latitude (N) | Longitude (W) | Sample Depth | Temperature (C°) | SPC (ms/cm) | Cond (ms/cm) | Salinity (ppm) | DO (%) | DO (mg/L) | Depth (feet) | pH (S.U.) | ORP (mV) |
|-----------|--------------|----------|-----------------|-----------------|--------------|------------------|-------------|--------------|----------------|--------|-----------|--------------|-----------|----------|
| SB-101 | Total: 4'9" | Initial: | 34° 36' 49.576" | 77° 26' 18.704" | Bottom: | 31.57 | 16.98 | 19.02 | 9.86 | 123.1 | 8.58 | 3.78 | 8.42 | 183.9 |
| | Secchi: 1'8" | Final: | 34° 36' 49.705" | 77° 26' 19.168" | Middle: | 31.64 | 16.79 | 18.87 | 9.77 | 127.1 | 8.86 | 2.30 | 8.48 | 186.5 |
| | | | | | Surface: | 31.67 | 16.45 | 18.52 | 9.55 | 127.9 | 8.93 | 1.22 | 8.50 | 188.6 |
| SB-201 | Total: 4'4" | Initial: | 34° 36' 42.566" | 77° 26' 44.151" | Bottom: | 31.47 | 17.24 | 19.41 | 10.08 | 121.0 | 8.45 | 3.46 | 8.44 | 232.1 |
| | Secchi: 1'6" | Final: | 34° 36' 42.539" | 77° 26' 44.164" | Middle: | 31.54 | 17.05 | 19.20 | 9.96 | 122.5 | 8.54 | 2.10 | 8.48 | 233.5 |
| | | | | | Surface: | 31.55 | 17.21 | 19.37 | 10.00 | 122.6 | 8.54 | 0.93 | 8.49 | 234.7 |
| SB-301 | Total: 3'6" | Initial: | 34° 36' 39.766" | 77° 26' 42.329" | Bottom: | 31.35 | 17.07 | 19.14 | 9.96 | 120.4 | 8.40 | 2.53 | 8.43 | 208.6 |
| | Secchi: 1'2" | Final: | 34° 36' 39.050" | 77° 26' 43.261" | Surface: | 31.49 | 17.00 | 19.06 | 9.91 | 123.4 | 8.62 | 0.98 | 8.47 | 210.1 |
| SB-401 | Total: 2' | Initial: | 34° 36' 33.365" | 77° 26' 37.203" | Surface: | 31.95 | 16.26 | 18.42 | 116.10 | 8.07 | 9.45 | 1.017 | 8.47 | 234.3 |
| | Secchi: 1' | Final: | 34° 36' 33.336" | 77° 26' 36.763" | | | | | | | | | | |
| SB-501 | Total: 2'4" | Initial: | 34° 36' 19.066" | 77° 26' 20.169" | Surface: | 32.35 | 16.08 | 18.32 | 9.31 | 110.6 | 7.62 | 0.99 | 8.41 | 234.7 |
| | Secchi: 1' | Final: | 34° 36' 19.240" | 77° 26' 20.141" | | | | | | | | | | |

Notes:

ID = Sample Identification

N = North

W = West

Secchi = Secchi disk measurement

SPC = Specific Conductance

ms/cm = milliohms per centimeter

Cond = Conductivity

ppm = parts per million

mg/L = milligram per liter

S.U. = Standard Units

DO = Dissolved Oxygen

ORP = Oxidation-Reduction Potential

mV = millivolts

TABLE 3-4

WATER QUALITY MEASUREMENTS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| Sample ID | Depth (feet) | Temperature (C°) | Cond (ms/cm) | Salinity (ppt) | DO (mg/L) | pH (S.U.) |
|-----------|--------------|------------------|--------------|----------------|-----------|-----------|
| SB-SD01 | 1.5 | 9.5 | 35.9 | 22.6 | 11.32 | 8.36 |
| SB-SD02 | 5.0 | 12.2 | 32.4 | 19.9 | 9.85 | 8.30 |
| SB-SD03 | 2.0 | 12.9 | 33.2 | 20.7 | 10.33 | 8.27 |
| SB-SD04 | 6.0 | 11.70 | 34.8 | 22.0 | 10.80 | 8.38 |
| SB-SD05 | 3.0 | 9.3 | 33.5 | 20.6 | 11.30 | 8.54 |
| RF-SD01 | 4.0 | 9.8 | 39.3 | 24.5 | 10.43 | 8.45 |
| RF-SD02 | 4.0 | 9.7 | 36.0 | 22.6 | 11.26 | 8.53 |

Notes:

ID = Sample Identification
ms/cm = milliohms per centimeter
Cond = Conductivity
ppt = parts per thousand
mg/L = milligram per liter
S.U. = Standard Units
DO = Dissolved Oxygen

TABLE 3-5

**POSITIVE DETECTIONS IN SEDIMENT
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID | SB-SD01-99A | SB-SD02-99A | SB-SD03-99A | SB-SD04-99A | SB-SD05-99A |
|---------------------------|-------------|-------------|-------------|-------------|-------------|
| SAMPLE DATE | 1/14/99 | 1/15/99 | 1/15/99 | 1/15/99 | 1/16/99 |
| INORGANICS (mg/kg) | | | | | |
| Copper | 2.67 U | 16.6 | 2.4 J | 8.7 J | 2.35 U |
| Lead | 6 | 5.7 | 2.5 | 20.7 | 3.4 |

TABLE 3-5 (continued)

**POSITIVE DETECTIONS IN SEDIMENT
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Copper | 2.35 U | 2.67 U | 2.4 J | 16.6 | SB-SD02-99A | 3/5 | 9.23 | 8.7 |
| Lead | ND | ND | 2.5 | 20.7 | SB-SD04-99A | 5/5 | 7.66 | 5.7 |

TABLE 3-5 (continued)

POSITIVE DETECTIONS IN SEDIMENT
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| SAMPLE ID | RF-SD01-99A | RF-SD02-99A |
|--------------------|-------------|-------------|
| SAMPLE DATE | 1/16/99 | 1/16/99 |
| INORGANICS (mg/kg) | | |
| Copper | 2.4 U | 4.6 J |
| Lead | 4.6 | 10.5 |

TABLE 3-5 (continued)

POSITIVE DETECTIONS IN SEDIMENT
 REFERENCE AREAS
 STONE BAY AQUATIC ASSESSMENT, CTO-0100
 MCB, CAMP LEJEUNE, NORTH CAROLINA

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Copper | 2.4 U | 2.4 U | 4.6 J | 4.6 J | RF-SD02-99A | 1/2 | 4.6 | 4.6 |
| Lead | ND | ND | 4.6 | 10.5 | RF-SD02-99A | 2/2 | 7.55 | 7.55 |

TABLE 3-5 (continued)

POSITIVE DETECTIONS IN SEDIMENT
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| SAMPLE ID | RF-SD01-99A | RF-SD02-99A |
|----------------------|-------------|-------------|
| SAMPLE DATE | 1/16/99 | 1/16/99 |
| TOC (mg/kg) | | |
| Total Organic Carbon | 4930 | 12400 |

TABLE 3-5 (continued)

**POSITIVE DETECTIONS IN SEDIMENT
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|--|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| TOC (mg/kg) Total Organic Carbon | ND | ND | 4930 | 12400 | RF-SD02-99A | 2/2 | 8665 | 8665 |

TABLE 3-6

**POSITIVE DETECTIONS IN CLAM TISSUE
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| | |
|---------------------------|-------------|
| SAMPLE ID | SB-CL04-99A |
| SAMPLE DATE | 1/15/99 |
| INORGANICS (mg/kg) | |
| Copper | 12.6 |
| Lead | 1.8 |
| WET WEIGHT BASIS | |
| Percent Lipids (%) | 0.2 |
| Moisture (%) | 89 |

TABLE 3-6 (continued)

**POSITIVE DETECTIONS IN CLAM TISSUE
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Copper | ND | ND | 12.6 | 12.6 | SB-CL04-99A | 1/1 | 12.6 | 12.6 |
| Lead | ND | ND | 1.8 | 1.8 | SB-CL04-99A | 1/1 | 1.8 | 1.8 |
| WET WEIGHT BASIS | | | | | | | | |
| Percent Lipids (%) | ND | ND | 0.2 | 0.2 | SB-CL04-99A | 1/1 | 0.2 | 0.2 |
| Moisture (%) | ND | ND | 89 | 89 | SB-CL04-99A | 1/1 | 89 | 89 |

TABLE 3-6 (continued)

POSITIVE DETECTIONS IN CLAM TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| SAMPLE ID | RF-CL01-99A | RF-CL02-99A |
|---------------------------|-------------|-------------|
| SAMPLE DATE | 1/16/99 | 1/16/99 |
| INORGANICS (mg/kg) | | |
| Copper | 9.5 | 10.1 |
| Lead | 1 | 0.7 |
| WET WEIGHT BASIS | | |
| Percent Lipids (%) | 0.2 | 0.1 |
| Moisture (%) | 85 | 88 |

TABLE 3-6 (continued)

**POSITIVE DETECTIONS IN CLAM TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|----------------------------------|-------------------------------|-------------------------------|-----------------------------|-----------------------------|---------------------------------------|-----------------------------------|---|------------------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Copper | ND | ND | 9.5 | 10.1 | RF-CL02-99A | 2/2 | 9.8 | 9.8 |
| Lead | ND | ND | 0.7 | 1 | RF-CL01-99A | 2/2 | 0.85 | 0.85 |
| WET WEIGHT BASIS | | | | | | | | |
| Percent Lipids (%) | ND | ND | 0.1 | 0.2 | RF-CL01-99A | 2/2 | 0.15 | 0.15 |
| Moisture (%) | ND | ND | 85 | 88 | RF-CL02-99A | 2/2 | 86.5 | 86.5 |

TABLE 3-7

**POSITIVE DETECTIONS IN MUSSEL TISSUE
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID | SB-MU01/02-99A | SB-MU04-99A | SB-MU05-99A |
|---------------------------|-----------------------|--------------------|--------------------|
| SAMPLE DATE | 1/16/99 | 1/16/99 | 1/15/99 |
| INORGANICS (mg/kg) | | | |
| Copper | 12.5 | 12.9 | 4.1 |
| Lead | 4.8 | 1.6 | 1.3 |
| WET WEIGHT BASIS | | | |
| Percent Lipids (%) | 1.6 | 1 | 2.1 |
| Moisture (%) | 85 | 93 | 81 |

TABLE 3-7 (continued)

**POSITIVE DETECTIONS IN MUSSEL TISSUE
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Copper | ND | ND | 4.1 | 12.9 | SB-MU04-99A | 3/3 | 9.83 | 12.5 |
| Lead | ND | ND | 1.3 | 4.8 | SB-MU01/02-99A | 3/3 | 2.57 | 1.6 |
| WET WEIGHT BASIS | | | | | | | | |
| Percent Lipids (%) | ND | ND | 1 | 2.1 | SB-MU05-99A | 3/3 | 1.56667 | 1.6 |
| Moisture (%) | ND | ND | 81 | 93 | SB-MU04-99A | 3/3 | 86.33333 | 85 |

TABLE 3-7 (continued)

**POSITIVE DETECTIONS IN MUSSEL TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| | |
|---------------------------|-------------|
| SAMPLE ID | RF-MU02-99A |
| SAMPLE DATE | 1/16/99 |
| INORGANICS (mg/kg) | |
| Copper | 4.1 |
| Lead | 1 |
| WET WEIGHT BASIS | |
| Percent Lipids (%) | 1.1 |
| Moisture (%) | 82 |

TABLE 3-7 (continued)

**POSITIVE DETECTIONS IN MUSSEL TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Copper | ND | ND | 4.1 | 4.1 | RF-MU02-99A | 1/1 | 4.1 | 4.1 |
| Lead | ND | ND | 1 | 1 | RF-MU02-99A | 1/1 | 1 | 1 |
| WET WEIGHT BASIS | | | | | | | | |
| Percent Lipids (%) | ND | ND | 1.1 | 1.1 | RF-MU02-99A | 1/1 | 1.1 | 1.1 |
| Moisture (%) | ND | ND | 82 | 82 | RF-MU02-99A | 1/1 | 82 | 82 |

TABLE 3-8

**POSITIVE DETECTIONS IN OYSTER TISSUE
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID | SB-OY01-99A | SB-OY02-99A | SB-OY03-99A | SB-OY04-99A | SB-OY05-99A |
|---------------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| SAMPLE DATE | 01/14/99 | 01/15/99 | 01/15/99 | 01/15/99 | 01/15/99 |
| INORGANICS (mg/kg) | | | | | |
| Copper | 50.3 | 45.1 | 46 | 21.4 | 17.1 |
| Lead | 1.6 | 0.69 | 1 | 0.82 | 0.4 |
| WET WEIGHT BASIS | | | | | |
| Percent Lipids (%) | 0.2 | 0.3 | 0.6 | 0.6 | 0.4 |
| Moisture (%) | 85 | 84 | 88 | 89 | 80 |

TABLE 3-8 (continued)

POSITIVE DETECTIONS IN OYSTER TISSUE
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Copper | ND | ND | 17.1 | 50.3 | SB-OY01-99A | 5/5 | 35.98 | 45.1 |
| Lead | ND | ND | 0.4 | 1.6 | SB-OY01-99A | 5/5 | 0.9 | 0.82 |
| WET WEIGHT BASIS | | | | | | | | |
| Percent Lipids (%) | ND | ND | 0.2 | 0.6 | SB-OY03-99A,SB-OY04-99A | 5/5 | 0.42 | 0.4 |
| Moisture (%) | ND | ND | 80 | 89 | SB-OY04-99A | 5/5 | 85.2 | 85 |

TABLE 3-8 (continued)

**POSITIVE DETECTIONS IN OYSTER TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID | RF-OY01-99A | RF-OY02-99A |
|---------------------------|-------------|-------------|
| SAMPLE DATE | 1/16/99 | 1/16/99 |
| INORGANICS (mg/kg) | | |
| Copper | 8.2 | 88.2 |
| Lead | 2 | 1.2 |
| WET WEIGHT BASIS | | |
| Percent Lipids (%) | 0.1 | 0.6 |
| Moisture (%) | | |

TABLE 3-8 (continued)

**POSITIVE DETECTIONS IN OYSTER TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Copper | ND | ND | 8.2 | 88.2 | RF-OY02-99A | 2/2 | 48.2 | 48.2 |
| Lead | ND | ND | 1.2 | 2 | RF-OY01-99A | 2/2 | 1.6 | 1.6 |
| WET WEIGHT BASIS | | | | | | | | |
| Percent Lipids (%) | ND | ND | 0.1 | 0.6 | RF-OY02-99A | 2/2 | 0.35 | 0.35 |
| Moisture (%) | ND | ND | ND | ND | | 0/0 | ND | ND |

TABLE 4-1

**LOWEST OBSERVED ADVERSE EFFECT LEVELS /
NO OBSERVED ADVERSE EFFECT LEVELS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| Ecological Contaminant of Concern | Heron | | Mink | |
|--------------------------------------|-------|-------|-------|-------|
| | LOAEL | NOAEL | LOAEL | NOAEL |
| Copper | 2.35 | 0.235 | 10 | 1 |
| Lead | 3 | 0.3 | 1.5 | 0.15 |

Notes:

The studies from which these toxicity numbers are based can be found in Appendix F.

LOAELs and NOAELS are reported in mg/kg/day

LOAEL - Lowest Observed Adverse Effects Level

NOAEL - No Observed Adverse Effects Level

TABLE 4-2

FREQUENCY AND RANGE OF SEDIMENT DATA COMPARED TO SEDIMENT SCREENING LEVELS
 STUDY AREA
 STONE BAY AQUATIC ASSESSMENT, CTO-0100
 MCB, CAMP LEJEUNE, NORTH CAROLINA

| Analyte | Sediment Screening Values (SSVs) | | Contaminant Frequency/Range | | No. of Positive Detects Above SSV | | Max. HQ | Reference Areas | | | Ecological Contaminant of Concern? | Comments |
|---------|----------------------------------|---------------------|--|------------------------------|-----------------------------------|------|---------|-----------------|------|---|------------------------------------|-----------|
| | | | No. of Positive Detects/No. of Samples | Range of Positive Detections | ER-L | ER-M | | Range | Mean | No. of Positive Detects Above Reference | | |
| | SSV/ER-L ⁽¹⁾ | ER-M ⁽²⁾ | | | | | | | | | | |
| Copper | 18.7 | 270 | 3/5 | 2.4J - 16.6 | 0 | 0 | 0.89 | 4.6J | 4.6J | 2 | No | Below SSV |
| Lead | 30.2 | 218 | 5/5 | 2.5 - 20.7 | 0 | 0 | 0.69 | 4.6 - 10.5 | 7.55 | 1 | No | Below SSV |

Notes:

- J - value reported is estimated
- mg/kg - milligram per kilogram
- SSV - Sediment Screening Value
- HQ - Hazard Quotient (maximum detected value divided by the lowest screening value)

(1) Region IV Sediment Screening Value (USEPA, 1995), unless otherwise noted
 (2) Long et al. (1995) value, unless otherwise noted

TABLE 4-3

FREQUENCY AND RANGE OF CLAM TISSUE DATA COMPARED TO REFERENCE AREAS
 STUDY AREA
 STONE BAY AQUATIC ASSESSMENT, CTO-0100
 MCB, CAMP LEJEUNE, NORTH CAROLINA

| Analyte (mg/kg) | Contaminant Frequency/Range | | Reference Areas | | | Ecological Contaminant of Concern? | Comments |
|-----------------|--|------------------------------|-----------------|------|---|------------------------------------|----------|
| | No. of Positive Detects/No. of Samples | Range of Positive Detections | Range | Mean | No. of Positive Detects Above Reference | | |
| | | | | | | | |
| Copper | 1/1 | 12.6 | 9.5 - 10.1 | 9.8 | 1 | Yes | |
| Lead | 1/1 | 1.8 | 0.7 - 1 | 0.85 | 1 | Yes | |

Notes:

Shaded area represents selected ecological contaminants of concern

Clams were only found at one station in the study area

mg/kg - milligram per kilogram

TABLE 4-4

FREQUENCY AND RANGE OF MUSSEL TISSUE DATA COMPARED TO REFERENCE AREAS
 STUDY AREA
 STONE BAY AQUATIC ASSESSMENT, CTO-0100
 MCB, CAMP LEJEUNE, NORTH CAROLINA

| Analyte (mg/kg) | Contaminant Frequency/Range | | Reference Areas | | Ecological Contaminant of Concern? | Comments |
|-----------------|--|------------------------------|-----------------|---|------------------------------------|----------|
| | No. of Positive Detects/No. of Samples | Range of Positive Detections | Detection | No. of Positive Detects Above Reference | | |
| | | | | | | |
| Lead | 3/3 | 1.3 - 4.8 | 1 | 3 | Yes | |

Notes:

Shaded area represents selected ecological contaminants of concern

Reference is based on one sample: therefore, the mean value is not calculated.

mg/kg - milligram per kilogram

TABLE 4-5

FREQUENCY AND RANGE OF OYSTER TISSUE DATA COMPARED TO REFERENCE AREAS
 STUDY AREA
 STONE BAY AQUATIC ASSESSMENT, CTO-0100
 MCB, CAMP LEJEUNE, NORTH CAROLINA

| Analyte (mg/kg) | Contaminant Frequency/Range | | Reference Areas | | | Ecological Contaminant of Concern? | Comments |
|-----------------|--|------------------------------|-----------------|------|---|------------------------------------|-----------------|
| | No. of Positive Detects/No. of Samples | Range of Positive Detections | Range | Mean | No. of Positive Detects Above Reference | | |
| | | | | | | | |
| Copper | 5/5 | 17.1 - 50.3 | 8.2 - 88.2 | 48.2 | 0 | No | Below Reference |
| Lead | 5/5 | 0.4 - 1.6 | 1.2 - 2 | 1.6 | 0 | No | Below Reference |

Notes:

mg/kg - milligram per kilogram

TABLE 4-6

QUALITATIVE COMPARISON OF COPPER CONCENTRATIONS DETECTED IN BENTHIC SAMPLES
 STUDY AREA
 STONE BAY AQUATIC ASSESSMENT, CTO-0100
 MCB, CAMP LEJEUNE, NORTH CAROLINA

| Species | Range of Detections (mg/kg) | Mean Concentration (mg/kg) | Literature Values | | No. of Detects Above Literature Values | Reference Areas | |
|---------|-----------------------------------|----------------------------------|--------------------|-----------------|--|------------------|-----------------|
| | | | Maximum (mg/kg) | Mean (mg/kg) | | Range (mg/kg) | Mean (mg/kg) |
| Oyster | 17.1 - 50.3 | 35.98 | 360 | 150 | 0 | 8.2 - 88.2 | 48.2 |
| Clam | 12.6 | 12.6 | NF | NF | NA | 9.5 - 10.1 | 9.8 |
| Mussel | 4.1 - 12.9 | 9.83 | 11 | 8.9 | 2 | 4.1 | 4.1 |

Notes:

mg/kg - milligram per kilogram

NF - Not Found

NA - Not applicable

Source: Irwin, 1997a

TABLE 4-7

QUALITATIVE COMPARISON OF LEAD CONCENTRATIONS DETECTED IN BENTHIC SAMPLES
STUDY AREA

STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| Species | Range of Detections (mg/kg) | Mean Concentration (mg/kg) | Literature Values | | No. of Detects Above Literature Values | Reference Areas | |
|---------|-----------------------------------|----------------------------------|--------------------|-----------------|---|------------------|-----------------|
| | | | Maximum (mg/kg) | Mean (mg/kg) | | Range (mg/kg) | Mean (mg/kg) |
| Oyster | 0.44 - 1.6 | 0.9 | 0.94 | 0.52 | 2 | 1.2 - 2 | 1.6 |
| Clam | 1.8 | 1.8 | NF | NF | NA | 0.7 - 1 | 0.85 |
| Mussel | 1.3 - 4.8 | 2.57 | 4.3 | 1.8 | 1 | 1 | 1 |

Notes:

mg/kg - milligram per kilogram

NF - Not Found

NA - Not applicable

Source: Irwin, 1997b

TABLE 4-8

SUMMARY OF ECOLOGICAL CONTAMINANTS OF CONCERN PER MEDIA
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| Analyte | Sediment | Clam | Mussel | Oyster |
|---------|----------|------|--------|--------|
| Copper | | X | X | |
| Lead | | X | X | |

TABLE 4-9

**AQUATIC SPECIES - CONSERVATIVE MODELS
MAXIMUM CONCENTRATION HAZARD QUOTIENT VALUES
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| Ecological Contaminants of Concern | Heron | | Mink | |
|---------------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| | NOAEL HQ _n | LOAEL HQ ₁ | NOAEL HQ _n | LOAEL HQ ₁ |
| Copper | 44.0 | 4.0 | 13.0 | 1.0 |
| Lead | 4.0 | 0.4 | 20.0 | 2.0 |

Highlighted values represent Hazard Quotients (HQs) greater than 1.0

HQ_n. Hazard Quotient based on the NOAEL

HQ₁. Hazard Quotient based on the LOAEL

TABLE 4-10

**AQUATIC SPECIES - LESS CONSERVATIVE MODELS
MEAN CONCENTRATION HAZARD QUOTIENT VALUES
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| Ecological Contaminants of Concern | Heron | | Mink | |
|---------------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| | NOAEL HQ _n | LOAEL HQ _l | NOAEL HQ _n | LOAEL HQ _l |
| Copper | 31.0 | 3.0 | 9.0 | 0.9 |
| Lead | 2.0 | 0.2 | 9.0 | 0.9 |

Highlighted values represent Hazard Quotients (HQs) greater than 1.0

HQ_n - Hazard Quotient based on the NOAEL

HQ_l - Hazard Quotient based on the LOAEL

TABLE 4-11

AQUATIC SPECIES - LESS CONSERVATIVE MODELS
MEAN CONCENTRATION HAZARD QUOTIENT VALUES
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| Ecological Contaminants of Concern | Heron | | Mink | |
|---------------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| | NOAEL HQ _n | LOAEL HQ _l | NOAEL HQ _n | LOAEL HQ _l |
| Copper | 41.0 | 4.0 | 12.0 | 1.0 |
| Lead | 2.0 | 0.2 | 7.0 | 0.7 |

Highlighted values represent Hazard Quotients (HQs) greater than 1.0


HQ_n. Hazard Quotient based on the NOAEL

HQ_l. Hazard Quotient based on the LOAEL

Baker

Baker Environmental, Inc.


FIGURES



2100505W

FIGURE 1-1
 SITE LOCATION MAP
 STONE BAY AQUATIC ASSESSMENT
 CTO-100
 MARINE CORPS BASE, CAMP LEJEUNE
 NORTH CAROLINA

02323AABIY



LEGEND


- SEDIMENT/BIOTA SAMPLE STATION
- SEDIMENT/BIOTA REFERENCE STATION
- CH2M HILL SEDIMENT SAMPLE STATION
- APPROXIMATE LOCATION OF SHELLFISH PLANTING SITES
- RIFLE RANGE FAN
- STONE BAY RIFLE RANGE FAN
- SPOIL AREA

FIGURE 3-1
SAMPLE LOCATION MAP
STONE BAY AQUATIC ASSESSMENT
CTO-100
MARINE CORPS BASE, CAMP LEJEUNE
NORTH CAROLINA


1200 0 600 1200
 1 inch = 1200 ft.



SOURCE OF BASE MAPPING: MCB CAMP LEJEUNE, 1998.



SOURCE OF BASE MAPPING: MCB CAMP LEJEUNE, 1998.



LEGEND

- SEDIMENT/BIOTA SAMPLE STATION
- CH2M HILL SEDIMENT SAMPLE STATION
- ATTEMPTED BIOTA SAMPLE STATION (NO SPECIES PRESENT)
- APPROXIMATE LOCATION OF SHELLFISH PLANTING SITES
- RIFLE RANGE FAN
- STONE BAY RIFLE RANGE FAN

FIGURE 3-2
STUDY AREA LOCATION MAP
STONE BAY AQUATIC ASSESSMENT
CTO-100
MARINE CORPS BASE, CAMP LEJEUNE
NORTH CAROLINA

Baker

Baker Environmental, Inc.
ATTACHMENT A

Site Photographs

Rifle Range



1999.01

This photograph of Bravo Range was taken facing down range (north). The target backstop seen in the background is approximately 500 meters away.



1999.02

Bravo Range. The asphalt road leads to the target area, which is commonly referred to as the "Butts".

02323AAB2Y

Rifle Range

1999.03

This photograph was taken behind the target back-stop area or "Butts". Personnel protected by the backstop raise and lower targets for the shooters.



1999.04

This photograph was taken from the top of the range backstop, looking north in the direction of Stone Bay. Note that the tree line height is diminished, due to the number of rounds traveling over the area.



Stone Bay



1999.05

Sampling in deep water was accomplished using "shellfish tongs" to retrieve clams, oysters, and mussels from the bottom of the bay.



1999.06

Some shellfish samples were collected by hand, as samplers waded in shallow water. This photograph was taken at the end of the rifle range near Stone Creek.

Stone Bay

1999.07

This photograph was taken from Stone Bay facing toward the edge of the rifle range. The flag pole seen in the center of the photograph is used to warn civilian personnel when the rifle range is in operation.



1999.08

This photograph was taken from reference station number one. Sneads Ferry Bridge can be seen in the background.



Baker

Baker Environmental, Inc.
ATTACHMENT B

Analytical Results

B.1

CH2M Hill Letter Report

TECHNICAL MEMORANDUM

CH2MHILL

Stones Bay Sediment and Water Quality Sampling Results

PREPARED FOR: Mr. D.J. Cotnoir/Commander Atlantic Division

PREPARED BY: Bill Kreuzberger/CH2M HILL-CLT

COPIES: Mr. Dave Marasco/Camp Lejeune
Mike Mischuk/CH2M HILL-MKE
Tom Dupuis/CH2M HILL-MKE
Stewart Barnes/CH2M HILL-HRO
Steven Lehmann/CH2M HILL-CLT

DATE: October 28, 1998

Introduction and Background

Marine Corp Base (MCB), Camp Lejeune provides specialized training to prepare troops for amphibious and land combat operations. The buildings and facilities onsite support 144,000 marines, sailors and their families. A new 15-million gallon per day (mgd) advanced wastewater treatment facility is being constructed; and due to strong public reaction, the North Carolina Department of Environment and Natural Resources (NCDENR), Division of Coastal Management, has required Camp Lejeune to conduct in-stream sediment and water quality monitoring.

A monitoring program was initiated in June 1998 to quantify the discharge's impact, if any, on the estuarine environment. Ten transects were sampled at locations between Wilson Bay and Courthouse Bay. A modification to the monitoring program was requested by LANTDIV and required additional sediment and water quality sampling at five stations in an area of the New River Estuary known as Stones Bay.

Field Sampling

On September 16, 1998, CH2M HILL collected sediment and water quality samples from five stations in Stones Bay. Bob Deppen navigated the boat provided by Camp Lejeune. Dave Marasco, Camp Lejeune contact, was also in attendance. The sampling plan, Attachment 1, identifies the sampling locations, the sample matrix, chemical analysis that was performed, and the sampling methods used. The plan was followed as described with the exception of the following deviations.

- SB-3 could not be reached due to a water depth of less than two feet. SB-3 was sampled at an alternate location that was at the mouth of the stream. The final locations of all sites are noted on the attached map.
- As noted in Exhibit 3 of the attached sampling plan, a new polyethylene pail and spoon, each of which had been decontaminated previously, were supplied and used at each site instead of completing the decontamination process on a single pail and spoon between sites.
- The metals samples were filtered between 26 and 29 hours after the samples were first taken.
- Sampling was completed two weeks after Hurricane Bonnie, and the river was still turbid.

Post-It™ brand fax transmittal memo 7671 # of pages ▶ 4

| | | | |
|-------|--------------|---------|--------------|
| To | JEFF TERSIC | From | KATE LANDMAN |
| Co. | | Co. | |
| Dept. | | Phone # | |
| Fax # | 412 269 2002 | Fax # | |

STONES BAY SEDIMENT AND WATER QUALITY SAMPLING RESULTS

Results

The tables below summarize the sediment and water quality results from the attached lab reports from En Chem and Frontier Geosciences. Table 1, Sediment Inorganic Results, includes the metals, solids percent, acid volatile sulfide (AVS) and total organic carbon (TOC) results from the sediment samples, as well as the ER-L and ER-M levels established by the National Oceanic and Atmospheric Administration (NOAA). Graph 1, Grain Size Distribution, represents the grain size distribution from the same sediment samples, and the graph reveals that more fines are present at stations SB-3 and SB-4, while more sand is present at stations SB-1 and SB-5. Table 2, Water Quality Metals Results, summarizes the water quality results, as well as includes the NCDENR Water Quality Standards for tidal saltwaters with a classification of SA. There are no sediment standards to report in Table 1.

TABLE 1: SEDIMENT INORGANIC RESULTS

| Analyte | Units | Sampling Stations | | | | | NOAA Guidelines | |
|-------------|-------|-------------------|------------|------------|------------|------------|-----------------|------|
| | | CLMSDSB101 | CLMSDSB201 | CLMSDSB301 | CLMSDSB401 | CLMSDSB501 | ER-L | ER-M |
| As | mg/kg | < 1.3 | 3.6 | 2.2 | < 1.4 | < 1.4 | 33 | 85 |
| Cd | mg/kg | < 0.13 | < 0.21 | < 0.15 | < 0.14 | < 0.14 | 5 | 9 |
| Cr | mg/kg | 2.4 | 15 | 4.6 | 3.6 | 4.7 | 80 | 145 |
| Cu | mg/kg | < 1.3 | 18 | 5.7 | 15 | < 1.4 | 70 | 390 |
| Pb | mg/kg | 1.5 | 12 | 4 | 3.5 | 3.5 | 35 | 110 |
| Hg | mg/kg | < 0.13 | < 0.21 | < 0.15 | < 0.14 | < 0.14 | 0.15 | 1.30 |
| Ni | mg/kg | < 0.67 | 3.10 | < 0.77 | < 0.70 | 0.98 | 30 | 50 |
| Se | mg/kg | < 1.3 | < 2.1 | < 1.5 | < 1.4 | < 1.4 | NA | NA |
| Ag | mg/kg | < 0.67 | < 1.1 | < 0.77 | < 0.70 | < 0.70 | 1 | 2.2 |
| Zn | mg/kg | 3.7 | 19 | 5.5 | 4.2 | 4.0 | 120 | 270 |
| Solids | % | 74.4 | 47 | 65.3 | 71.8 | 71.9 | NA | NA |
| TOC as NPOC | mg/kg | 2000 | 11000 | 3300 | 2600 | 2000 | NA | NA |
| AVS | mg/kg | 130 | 270 | < 61 | 51 | < 56 | NA | NA |

TOC = Total Organic Carbon

NPOC = Non-purgeable Organic Carbon

AVS = Acid Volatile Sulfide

STONES BAY SEDIMENT AND WATER QUALITY SAMPLING RESULTS





TABLE 3: WATER QUALITY METALS RESULTS

| Metal | Units | Trip Blank | | CLMSWSB101 | | CLMSWSB201 | | CLMSWSB301 | | CLMSWSB401 | | CLMSWSB501 | | W. Q. Stds. Aquatic Life |
|-------|-------|------------|-------|------------|-------|------------|-------|------------|-------|------------|-------|------------|-------|-----------------------------|
| | | Total | Diss. | |
| As | µg/L | ND | ND | 1.06 | 0.943 | 1.16 | 1.03 | 1.27 | 1.02 | 1.44 | 1.04 | 1.37 | 1.05 | 50 |
| Cd | µg/L | 0.001 | 0 | 0.011 | 0.004 | 0.011 | 0.004 | 0.012 | 0.003 | 0.017 | 0.004 | 0.014 | 0.003 | 5.0 |
| Cr | µg/L | ND | ND | 0.29 | 0.13 | 0.56 | 0.10 | 0.72 | 0.06 | 1.37 | 0.06 | 1.30 | 0.07 | 20 |
| Cu | µg/L | 0.08 | 0.03 | 0.71 | 0.62 | 0.88 | 0.61 | 0.91 | 0.61 | 1.53 | 0.86 | 0.95 | 0.58 | 3 (AL) |
| Pb | µg/L | ND | ND | 0.220 | ND | 0.415 | ND | 0.437 | ND | 0.926 | ND | 0.641 | ND | 25 (N) |
| Hg | ng/L | 0.04 | 0.39 | 1.81 | 0.90 | 2.28 | 0.89 | 2.21 | 0.85 | 2.29 | 1.02 | 2.59 | 1.00 | 25 |
| Ni | µg/L | 0.02 | 0.02 | 0.24 | 0.24 | 0.30 | 0.25 | 0.31 | 0.26 | 0.48 | 0.25 | 0.44 | 0.29 | 8.3 |
| Se | µg/L | ND | ND | 0.120 | 0.095 | 0.126 | 0.099 | 0.122 | 0.103 | 0.157 | 0.102 | 0.153 | 0.102 | 71 |
| Ag | µg/L | ND | ND | 0.001 | ND | 0.004 | ND | 0.004 | ND | 0.005 | ND | 0.004 | ND | 0.1 (AL) |
| Zn | µg/L | ND | ND | 0.84 | ND | 1.33 | ND | 1.28 | ND | 2.29 | 0.24 | 1.75 | 0.15 | 86 (AL) |

ND = Analyte not detected above the estimated method detection limit (MDL)

AL = Values represent action levels as specified in 15A NCAC 2B .0220

N = See 15A NCAC 2B.0220 for narrative description of limits



Legend

- Final
- ~ Hydrography
- Transects

1000 0 1000 2000 Feet

B.2
Aquatic Assessment Data

**FREQUENCY OF DETECTIONS IN SEDIMENT
STONE BAY
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID | SB-SD01-99A | SB-SD02-99A | SB-SD03-99A | SB-SD04-99A | SB-SD05-99A |
|---------------------------|-------------|-------------|-------------|-------------|-------------|
| SAMPLE DATE | 1/14/99 | 1/15/99 | 1/15/99 | 1/15/99 | 1/16/99 |
| INORGANICS (mg/kg) | | | | | |
| Aluminum | 1750 | 1780 | 1580 | 14800 | 1580 |
| Antimony | 0.66 U | 0.61 U | 0.57 U | 1.36 U | 0.58 U |
| Arsenic | 1.1 J | 0.85 J | 0.68 U | 6.8 | 0.84 J |
| Barium | 2.9 J | 2.1 J | 2.2 J | 14.4 J | 2.7 J |
| Beryllium | 0.25 U | 0.23 U | 0.22 U | 0.52 U | 0.22 U |
| Cadmium | 0.1 U | 0.09 U | 0.09 U | 0.21 U | 0.09 U |
| Calcium | 484 J | 161 J | 323 J | 4930 | 1320 |
| Chromium | 5.6 | 4.2 | 3.2 | 29.8 | 3.9 |
| Cobalt | 1.37 U | 1.26 U | 1.18 U | 2.82 U | 1.21 U |
| Copper | 2.67 U | 16.6 | 2.4 J | 8.7 J | 2.35 U |
| Iron | 2380 | 2120 | 1480 | 17400 | 1660 |
| Lead | 6 | 5.7 | 2.5 | 20.7 | 3.4 |
| Magnesium | 843 J | 592 J | 511 J | 5280 | 475 J |
| Manganese | 8.9 | 6.3 | 4.2 | 88 | 8.7 |
| Mercury | 0.05 U | 0.06 U | 0.06 U | 0.13 U | 0.06 U |
| Nickel | 2.54 U | 2.33 U | 2.19 U | 5.8 J | 2.24 U |
| Potassium | 430 J | 341 J | 285 J | 2510 J | 273 J |
| Selenium | 0.81 U | 0.75 U | 0.7 U | 1.67 U | 0.72 U |
| Silver | 0.75 J | 0.9 J | 0.46 J | 4.6 J | 0.6 J |
| Sodium | 3980 | 2530 | 2080 | 15000 | 1860 |
| Thallium | 0.51 U | 0.47 U | 0.44 U | 1.3 J | 0.45 U |
| Vanadium | 5.3 J | 4.5 J | 3.7 J | 25.7 J | 3.6 J |
| Zinc | 5.8 | 5.6 | 2.9 J | 38.8 | 3.3 J |

**FREQUENCY OF DETECTIONS IN SEDIMENT
STONE BAY
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Aluminum | ND | ND | 1580 | 14800 | SB-SD04-99A | 5/5 | 4298 | 1750 |
| Antimony | 0.57 U | 1.36 U | ND | ND | | 0/5 | ND | ND |
| Arsenic | 0.68 U | 0.68 U | 0.84 J | 6.8 | SB-SD04-99A | 4/5 | 2.4 | 0.98 |
| Barium | ND | ND | 2.1 J | 14.4 J | SB-SD04-99A | 5/5 | 4.86 | 2.7 |
| Beryllium | 0.22 U | 0.52 U | ND | ND | | 0/5 | ND | ND |
| Cadmium | 0.09 U | 0.21 U | ND | ND | | 0/5 | ND | ND |
| Calcium | ND | ND | 161 J | 4930 | SB-SD04-99A | 5/5 | 1443.6 | 484 |
| Chromium | ND | ND | 3.2 | 29.8 | SB-SD04-99A | 5/5 | 9.34 | 4.2 |
| Cobalt | 1.18 U | 2.82 U | ND | ND | | 0/5 | ND | ND |
| Copper | 2.35 U | 2.67 U | 2.4 J | 16.6 | SB-SD02-99A | 3/5 | 9.23 | 8.7 |
| Iron | ND | ND | 1480 | 17400 | SB-SD04-99A | 5/5 | 5008 | 2120 |
| Lead | ND | ND | 2.5 | 20.7 | SB-SD04-99A | 5/5 | 7.66 | 5.7 |
| Magnesium | ND | ND | 475 J | 5280 | SB-SD04-99A | 5/5 | 1540.2 | 592 |
| Manganese | ND | ND | 4.2 | 88 | SB-SD04-99A | 5/5 | 23.22 | 8.7 |
| Mercury | 0.05 U | 0.13 U | ND | ND | | 0/5 | ND | ND |
| Nickel | 2.19 U | 2.54 U | 5.8 J | 5.8 J | SB-SD04-99A | 1/5 | 5.8 | 5.8 |
| Potassium | ND | ND | 273 J | 2510 J | SB-SD04-99A | 5/5 | 767.8 | 341 |
| Selenium | 0.7 U | 1.67 U | ND | ND | | 0/5 | ND | ND |
| Silver | ND | ND | 0.46 J | 4.6 J | SB-SD04-99A | 5/5 | 1.46 | 0.75 |
| Sodium | ND | ND | 1860 | 15000 | SB-SD04-99A | 5/5 | 5090 | 2530 |
| Thallium | 0.44 U | 0.51 U | 1.3 J | 1.3 J | SB-SD04-99A | 1/5 | 1.3 | 1.3 |
| Vanadium | ND | ND | 3.6 J | 25.7 J | SB-SD04-99A | 5/5 | 8.56 | 4.5 |
| Zinc | ND | ND | 2.9 J | 38.8 | SB-SD04-99A | 5/5 | 11.28 | 5.6 |

**FREQUENCY OF DETECTIONS IN CLAM TISSUE
STONE BAY
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

SAMPLE ID SB-CL04-99A
SAMPLE DATE 1/15/99

INORGANICS (mg/kg)

| | |
|-----------|--------|
| Aluminum | 275 |
| Antimony | 1.85 U |
| Arsenic | 12.4 |
| Barium | 1.7 J |
| Beryllium | 0.71 U |
| Cadmium | 0.47 |
| Calcium | 12400 |
| Chromium | 1.2 |
| Cobalt | 2.3 J |
| Copper | 12.6 |
| Iron | 442 |
| Lead | 1.8 |
| Magnesium | 8370 |
| Manganese | 33.5 |
| Mercury | 0.048 |
| Nickel | 4.1 |
| Potassium | 11600 |
| Selenium | 2.8 |
| Silver | 0.57 U |
| Sodium | 71900 |
| Thallium | 1.42 U |
| Vanadium | 1.9 J |
| Zinc | 117 |

**FREQUENCY OF DETECTIONS IN CLAM TISSUE
STONE BAY
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Aluminum | ND | ND | 275 | 275 | SB-CL04-99A | 1/1 | 275 | 275 |
| Antimony | 1.85 U | 1.85 U | ND | ND | | 0/1 | ND | ND |
| Arsenic | ND | ND | 12.4 | 12.4 | SB-CL04-99A | 1/1 | 12.4 | 12.4 |
| Barium | ND | ND | 1.7 J | 1.7 J | SB-CL04-99A | 1/1 | 1.7 | 1.7 |
| Beryllium | 0.71 U | 0.71 U | ND | ND | | 0/1 | ND | ND |
| Cadmium | ND | ND | 0.47 | 0.47 | SB-CL04-99A | 1/1 | 0.47 | 0.47 |
| Calcium | ND | ND | 12400 | 12400 | SB-CL04-99A | 1/1 | 12400 | 12400 |
| Chromium | ND | ND | 1.2 | 1.2 | SB-CL04-99A | 1/1 | 1.2 | 1.2 |
| Cobalt | ND | ND | 2.3 J | 2.3 J | SB-CL04-99A | 1/1 | 2.3 | 2.3 |
| Copper | ND | ND | 12.6 | 12.6 | SB-CL04-99A | 1/1 | 12.6 | 12.6 |
| Iron | ND | ND | 442 | 442 | SB-CL04-99A | 1/1 | 442 | 442 |
| Lead | ND | ND | 1.8 | 1.8 | SB-CL04-99A | 1/1 | 1.8 | 1.8 |
| Magnesium | ND | ND | 8370 | 8370 | SB-CL04-99A | 1/1 | 8370 | 8370 |
| Manganese | ND | ND | 33.5 | 33.5 | SB-CL04-99A | 1/1 | 33.5 | 33.5 |
| Mercury | ND | ND | 0.048 | 0.048 | SB-CL04-99A | 1/1 | 0.05 | 0.05 |
| Nickel | ND | ND | 4.1 | 4.1 | SB-CL04-99A | 1/1 | 4.1 | 4.1 |
| Potassium | ND | ND | 11600 | 11600 | SB-CL04-99A | 1/1 | 11600 | 11600 |
| Selenium | ND | ND | 2.8 | 2.8 | SB-CL04-99A | 1/1 | 2.8 | 2.8 |
| Silver | 0.57 U | 0.57 U | ND | ND | | 0/1 | ND | ND |
| Sodium | ND | ND | 71900 | 71900 | SB-CL04-99A | 1/1 | 71900 | 71900 |
| Thallium | 1.42 U | 1.42 U | ND | ND | | 0/1 | ND | ND |
| Vanadium | ND | ND | 1.9 J | 1.9 J | SB-CL04-99A | 1/1 | 1.9 | 1.9 |
| Zinc | ND | ND | 117 | 117 | SB-CL04-99A | 1/1 | 117 | 117 |

**FREQUENCY OF DETECTIONS IN CLAM TISSUE
STONE BAY
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| | |
|-------------|-------------|
| SAMPLE ID | SB-CL04-99A |
| SAMPLE DATE | 1/15/99 |

WET WEIGHT BASIS

| | |
|--------------------|-----|
| Percent Lipids (%) | 0.2 |
| Moisture (%) | 89 |

**FREQUENCY OF DETECTIONS IN MUSSEL TISSUE
STONE BAY
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID | SB-MU01/02-99A | SB-MU04-99A | SB-MU05-99A |
|---------------------------|----------------|-------------|-------------|
| SAMPLE DATE | 1/16/99 | 1/16/99 | 1/15/99 |
| INORGANICS (mg/kg) | | | |
| Aluminum | 1010 | 455 | 317 |
| Antimony | 1.11 U | 38.81 U | 0.09 U |
| Arsenic | 11 | 37.4 | 11.3 |
| Barium | 1.5 J | 2.3 J | 0.83 J |
| Beryllium | 0.05 J | 14.93 U | 0.03 U |
| Cadmium | 0.24 | 0.52 | 0.19 |
| Calcium | 19500 | 24100 | 8470 |
| Chromium | 1.9 | 2 | 0.82 |
| Cobalt | 0.55 J | 0.64 J | 0.35 J |
| Copper | 12.5 | 12.9 | 4.1 |
| Iron | 1100 | 643 | 366 |
| Lead | 4.8 | 1.6 | 1.3 |
| Magnesium | 3080 | 7950 | 3010 |
| Manganese | 26.3 | 91.9 | 18.1 |
| Mercury | 0.088 | 0.157 | 0.086 |
| Nickel | 3 | 3 J | 0.73 J |
| Potassium | 7130 | 18900 | 6080 |
| Selenium | 3.4 | 9.4 | 3.3 |
| Silver | 0.34 U | 11.94 U | 0.03 U |
| Sodium | 24100 | 69800 | 27600 |
| Thallium | 0.86 U | 29.85 U | 0.07 U |
| Vanadium | 3.3 | 4.9 | 1.9 |
| Zinc | 45 | 101 | 34.1 |

**FREQUENCY OF DETECTIONS IN MUSSEL TISSUE
STONE BAY
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Aluminum | ND | ND | 317 | 1010 | SB-MU01/02-99A | 3/3 | 594 | 455 |
| Antimony | 0.09 U | 38.81 U | ND | ND | | 0/3 | ND | ND |
| Arsenic | ND | ND | 11 | 37.4 | SB-MU04-99A | 3/3 | 19.9 | 11.3 |
| Barium | ND | ND | 0.83 J | 2.3 J | SB-MU04-99A | 3/3 | 1.54 | 1.5 |
| Beryllium | 0.03 U | 14.93 U | 0.05 J | 0.05 J | SB-MU01/02-99A | 1/3 | 0.05 | 0.05 |
| Cadmium | ND | ND | 0.19 | 0.52 | SB-MU04-99A | 3/3 | 0.32 | 0.24 |
| Calcium | ND | ND | 8470 | 24100 | SB-MU04-99A | 3/3 | 17356.67 | 19500 |
| Chromium | ND | ND | 0.82 | 2 | SB-MU04-99A | 3/3 | 1.57 | 1.9 |
| Cobalt | ND | ND | 0.35 J | 0.64 J | SB-MU04-99A | 3/3 | 0.51 | 0.55 |
| Copper | ND | ND | 4.1 | 12.9 | SB-MU04-99A | 3/3 | 9.83 | 12.5 |
| Iron | ND | ND | 366 | 1100 | SB-MU01/02-99A | 3/3 | 703 | 643 |
| Lead | ND | ND | 1.3 | 4.8 | SB-MU01/02-99A | 3/3 | 2.57 | 1.6 |
| Magnesium | ND | ND | 3010 | 7950 | SB-MU04-99A | 3/3 | 4680 | 3080 |
| Manganese | ND | ND | 18.1 | 91.9 | SB-MU04-99A | 3/3 | 45.43 | 26.3 |
| Mercury | ND | ND | 0.086 | 0.157 | SB-MU04-99A | 3/3 | 0.11 | 0.09 |
| Nickel | ND | ND | 0.73 J | 3 J | SB-MU01/02-99A,SB-MU04-99A | 3/3 | 2.24 | 3 |
| Potassium | ND | ND | 6080 | 18900 | SB-MU04-99A | 3/3 | 10703.33 | 7130 |
| Selenium | ND | ND | 3.3 | 9.4 | SB-MU04-99A | 3/3 | 5.37 | 3.4 |
| Silver | 0.03 U | 11.94 U | ND | ND | | 0/3 | ND | ND |
| Sodium | ND | ND | 24100 | 69800 | SB-MU04-99A | 3/3 | 40500 | 27600 |
| Thallium | 0.07 U | 29.85 U | ND | ND | | 0/3 | ND | ND |
| Vanadium | ND | ND | 1.9 | 4.9 | SB-MU04-99A | 3/3 | 3.37 | 3.3 |
| Zinc | ND | ND | 34.1 | 101 | SB-MU04-99A | 3/3 | 60.03 | 45 |

**FREQUENCY OF DETECTIONS IN MUSSEL TISSUE
STONE BAY
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID | SB-MU01/02-99A | SB-MU04-99A | SB-MU05-99A |
|-------------------------|----------------|-------------|-------------|
| SAMPLE DATE | 1/16/99 | 1/16/99 | 1/15/99 |
| WET WEIGHT BASIS | | | |
| Percent Lipids (%) | 1.6 | 1 | 2.1 |
| Moisture (%) | 85 | 93 | 81 |

**FREQUENCY OF DETECTIONS IN OYSTER TISSUE
STONE BAY
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID | SB-OY01-99A | SB-OY02-99A | SB-OY03-99A | SB-OY04-99A | SB-OY05-99A |
|---------------------------|-------------|-------------|-------------|-------------|-------------|
| SAMPLE DATE | 01/14/99 | 01/15/99 | 01/15/99 | 01/15/99 | 01/15/99 |
| INORGANICS (mg/kg) | | | | | |
| Aluminum | 111 | 49.3 | 90.6 | 69.5 | 78.9 |
| Antimony | 9.39 U | 12.42 U | 1.69 U | 2.65 U | 3.9 U |
| Arsenic | 12.5 | 7.4 | 10 | 11.9 | 4.1 |
| Barium | 0.83 J | 0.35 J | 0.62 J | 0.68 J | 0.3 J |
| Beryllium | 3.61 U | 4.78 U | 0.65 U | 1.02 U | 1.5 U |
| Cadmium | 1.6 | 1.1 | 1.7 | 1.3 | 0.6 |
| Calcium | 2330 | 2830 | 4040 | 7840 | 5960 |
| Chromium | 0.65 | 0.46 | 0.66 | 0.72 | 0.43 |
| Cobalt | 0.37 J | 0.34 J | 3.51 U | 5.51 U | 0.21 J |
| Copper | 50.3 | 45.1 | 46 | 21.4 | 17.1 |
| Iron | 225 | 123 | 199 | 228 | 137 |
| Lead | 1.6 | 0.69 | 1 | 0.82 | 0.4 |
| Magnesium | 2740 | 2450 | 3960 | 5110 | 2620 |
| Manganese | 9.8 | 6.4 | 11.8 | 7 | 5.7 |
| Mercury | 0.063 | 0.069 | 0.252 | 0.07 | 0.053 |
| Nickel | 2.4 | 1.1 J | 2.3 | 1.8 J | 1.3 |
| Potassium | 8940 | 8780 | 9090 | 11200 | 5000 |
| Selenium | 3.2 | 2.8 | 3.1 | 2.8 | 1 |
| Silver - | 0.34 J | 3.82 U | 0.52 U | 0.82 U | 1.2 U |
| Sodium | 19700 | 18600 | 31700 | 43800 | 22400 |
| Thallium | 7.22 U | 9.55 U | 1.3 U | 2.04 U | 3 U |
| Vanadium | 1.4 J | 0.68 J | 0.56 J | 1.7 J | 0.35 J |
| Zinc | 1280 | 793 | 722 | 728 | 894 |

**FREQUENCY OF DETECTIONS IN OYSTER TISSUE
STONE BAY
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Aluminum | ND | ND | 49.3 | 111 | SB-OY01-99A | 5/5 | 79.86 | 78.9 |
| Antimony | 1.69 U | 12.42 U | ND | ND | | 0/5 | ND | ND |
| Arsenic | ND | ND | 4.1 | 12.5 | SB-OY01-99A | 5/5 | 9.18 | 10 |
| Barium | ND | ND | 0.3 J | 0.83 J | SB-OY01-99A | 5/5 | 0.56 | 0.62 |
| Beryllium | 0.65 U | 4.78 U | ND | ND | | 0/5 | ND | ND |
| Cadmium | ND | ND | 0.6 | 1.7 | SB-OY03-99A | 5/5 | 1.26 | 1.3 |
| Calcium | ND | ND | 2330 | 7840 | SB-OY04-99A | 5/5 | 4600 | 4040 |
| Chromium | ND | ND | 0.43 | 0.72 | SB-OY04-99A | 5/5 | 0.58 | 0.65 |
| Cobalt | 3.51 U | 5.51 U | 0.21 J | 0.37 J | SB-OY01-99A | 3/5 | 0.31 | 0.34 |
| Copper | ND | ND | 17.1 | 50.3 | SB-OY01-99A | 5/5 | 35.98 | 45.1 |
| Iron | ND | ND | 123 | 228 | SB-OY04-99A | 5/5 | 182.4 | 199 |
| Lead | ND | ND | 0.4 | 1.6 | SB-OY01-99A | 5/5 | 0.9 | 0.82 |
| Magnesium | ND | ND | 2450 | 5110 | SB-OY04-99A | 5/5 | 3376 | 2740 |
| Manganese | ND | ND | 5.7 | 11.8 | SB-OY03-99A | 5/5 | 8.14 | 7 |
| Mercury | ND | ND | 0.053 | 0.252 | SB-OY03-99A | 5/5 | 0.1 | 0.07 |
| Nickel | ND | ND | 1.1 J | 2.4 | SB-OY01-99A | 5/5 | 1.78 | 1.8 |
| Potassium | ND | ND | 5000 | 11200 | SB-OY04-99A | 5/5 | 8602 | 8940 |
| Selenium | ND | ND | 1 | 3.2 | SB-OY01-99A | 5/5 | 2.58 | 2.8 |
| Silver | 0.52 U | 3.82 U | 0.34 J | 0.34 J | SB-OY01-99A | 1/5 | 0.34 | 0.34 |
| Sodium | ND | ND | 18600 | 43800 | SB-OY04-99A | 5/5 | 27240 | 22400 |
| Thallium | 1.3 U | 9.55 U | ND | ND | | 0/5 | ND | ND |
| Vanadium | ND | ND | 0.35 J | 1.7 J | SB-OY04-99A | 5/5 | 0.94 | 0.68 |
| Zinc | ND | ND | 722 | 1280 | SB-OY01-99A | 5/5 | 883.4 | 793 |

FREQUENCY OF DETECTIONS IN OYSTER TISSUE
STONE BAY
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| SAMPLE ID | SB-OY01-99A | SB-OY02-99A | SB-OY03-99A | SB-OY04-99A | SB-OY05-99A |
|-------------------------|-------------|-------------|-------------|-------------|-------------|
| SAMPLE DATE | 01/14/99 | 01/15/99 | 01/15/99 | 01/15/99 | 01/15/99 |
| WET WEIGHT BASIS | | | | | |
| Percent Lipids (%) | 0.2 | 0.3 | 0.6 | 0.6 | 0.4 |
| Moisture (%) | 85 | 84 | 88 | 89 | 80 |

**FREQUENCY OF DETECTIONS IN SEDIMENT
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID | RF-SD01-99A | RF-SD02-99A |
|---------------------------|-------------|-------------|
| SAMPLE DATE | 1/16/99 | 1/16/99 |
| INORGANICS (mg/kg) | | |
| Aluminum | 3000 | 8160 |
| Antimony | 0.59 U | 1.01 U |
| Arsenic | 2.4 | 6 |
| Barium | 3.5 J | 8.9 J |
| Beryllium | 0.23 U | 0.39 U |
| Cadmium | 0.09 U | 0.15 U |
| Calcium | 5320 | 26900 |
| Chromium | 7.8 | 19.2 |
| Cobalt | 1.23 U | 2.1 J |
| Copper | 2.4 U | 4.6 J |
| Iron | 3620 | 10100 |
| Lead | 4.6 | 10.5 |
| Magnesium | 1420 | 4200 |
| Manganese | 17.7 | 58.7 |
| Mercury | 0.05 U | 0.09 U |
| Nickel | 2.28 U | 4.7 J |
| Potassium | 605 J | 1500 J |
| Selenium | 0.73 U | 1.6 J |
| Silver | 1.5 J | 3.2 J |
| Sodium | 3470 | 9470 |
| Thallium | 0.46 U | 0.77 U |
| Vanadium | 7.7 J | 19.2 J |
| Zinc | 7.9 | 23.8 |

**FREQUENCY OF DETECTIONS IN SEDIMENT
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Aluminum | ND | ND | 3000 | 8160 | RF-SD02-99A | 2/2 | 5580 | 5580 |
| Antimony | 0.59 U | 1.01 U | ND | ND | | 0/2 | ND | ND |
| Arsenic | ND | ND | 2.4 | 6 | RF-SD02-99A | 2/2 | 4.2 | 4.2 |
| Barium | ND | ND | 3.5 J | 8.9 J | RF-SD02-99A | 2/2 | 6.2 | 6.2 |
| Beryllium | 0.23 U | 0.39 U | ND | ND | | 0/2 | ND | ND |
| Cadmium | 0.09 U | 0.15 U | ND | ND | | 0/2 | ND | ND |
| Calcium | ND | ND | 5320 | 26900 | RF-SD02-99A | 2/2 | 16110 | 16110 |
| Chromium | ND | ND | 7.8 | 19.2 | RF-SD02-99A | 2/2 | 13.5 | 13.5 |
| Cobalt | 1.23 U | 1.23 U | 2.1 J | 2.1 J | RF-SD02-99A | 1/2 | 2.1 | 2.1 |
| Copper | 2.4 U | 2.4 U | 4.6 J | 4.6 J | RF-SD02-99A | 1/2 | 4.6 | 4.6 |
| Iron | ND | ND | 3620 | 10100 | RF-SD02-99A | 2/2 | 6860 | 6860 |
| Lead | ND | ND | 4.6 | 10.5 | RF-SD02-99A | 2/2 | 7.55 | 7.55 |
| Magnesium | ND | ND | 1420 | 4200 | RF-SD02-99A | 2/2 | 2810 | 2810 |
| Manganese | ND | ND | 17.7 | 58.7 | RF-SD02-99A | 2/2 | 38.2 | 38.2 |
| Mercury | 0.05 U | 0.09 U | ND | ND | | 0/2 | ND | ND |
| Nickel | 2.28 U | 2.28 U | 4.7 J | 4.7 J | RF-SD02-99A | 1/2 | 4.7 | 4.7 |
| Potassium | ND | ND | 605 J | 1500 J | RF-SD02-99A | 2/2 | 1052.5 | 1052.5 |
| Selenium | 0.73 U | 0.73 U | 1.6 J | 1.6 J | RF-SD02-99A | 1/2 | 1.6 | 1.6 |
| Silver | ND | ND | 1.5 J | 3.2 J | RF-SD02-99A | 2/2 | 2.35 | 2.35 |
| Sodium | ND | ND | 3470 | 9470 | RF-SD02-99A | 2/2 | 6470 | 6470 |
| Thallium | 0.46 U | 0.77 U | ND | ND | | 0/2 | ND | ND |
| Vanadium | ND | ND | 7.7 J | 19.2 J | RF-SD02-99A | 2/2 | 13.45 | 13.45 |
| Zinc | ND | ND | 7.9 | 23.8 | RF-SD02-99A | 2/2 | 15.85 | 15.85 |

**FREQUENCY OF DETECTIONS IN SEDIMENT
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| | | |
|----------------------|-------------|-------------|
| SAMPLE ID | RF-SD01-99A | RF-SD02-99A |
| SAMPLE DATE | 1/16/99 | 1/16/99 |
| TOC (mg/kg) | | |
| Total Organic Carbon | 4930 | 12400 |

**FREQUENCY OF DETECTIONS IN SEDIMENT
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID | Minimum | Maximum | Minimum | Maximum | Location of | Frequency | Arithmetic Mean | Median |
|----------------------|------------|------------|----------|----------|----------------|--------------|------------------|------------------|
| SAMPLE DATE | Non-Detect | Non-Detect | Detected | Detected | Maximum Detect | of Detection | Positive Detects | Positive Detects |
| TOC (mg/kg) | | | | | | | | |
| Total Organic Carbon | ND | ND | 4930 | 12400 | RF-SD02-99A | 2/2 | 8665 | 8665 |

**FREQUENCY OF DETECTIONS IN CLAM TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID | RF-CL01-99A | RF-CL02-99A |
|---------------------------|-------------|-------------|
| SAMPLE DATE | 1/16/99 | 1/16/99 |
| INORGANICS (mg/kg) | | |
| Aluminum | 64.2 | 249 |
| Antimony | 5.63 U | 0.14 U |
| Arsenic | 11.1 | 16.2 |
| Barium | 0.99 J | 1.7 J |
| Beryllium | 2.16 U | 0.05 U |
| Cadmium | 0.21 | 0.23 J |
| Calcium | 3840 | 5720 |
| Chromium | 0.62 | 1.2 |
| Cobalt | 1.4 J | 2.1 J |
| Copper | 9.5 | 10.1 |
| Iron | 158 | 370 |
| Lead | 1 | 0.7 |
| Magnesium | 4950 | 6020 |
| Manganese | 9 | 9 |
| Mercury | 0.05 | 0.054 |
| Nickel | 3.6 | 4 |
| Potassium | 6240 | 9740 |
| Selenium | 2 | 3.1 |
| Silver | 1.73 U | 0.04 U |
| Sodium | 39900 | 48900 |
| Thallium | 4.33 U | 0.11 U |
| Vanadium | 2.5 | 3.1 |
| Zinc | 86.1 | 125 |

**FREQUENCY OF DETECTIONS IN CLAM TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Aluminum | ND | ND | 64.2 | 249 | RF-CL02-99A | 2/2 | 156.6 | 156.6 |
| Antimony | 0.14 U | 5.63 U | ND | ND | | 0/2 | ND | ND |
| Arsenic | ND | ND | 11.1 | 16.2 | RF-CL02-99A | 2/2 | 13.65 | 13.65 |
| Barium | ND | ND | 0.99 J | 1.7 J | RF-CL02-99A | 2/2 | 1.35 | 1.35 |
| Beryllium | 0.05 U | 2.16 U | ND | ND | | 0/2 | ND | ND |
| Cadmium | ND | ND | 0.21 | 0.23 J | RF-CL02-99A | 2/2 | 0.22 | 0.22 |
| Calcium | ND | ND | 3840 | 5720 | RF-CL02-99A | 2/2 | 4780 | 4780 |
| Chromium | ND | ND | 0.62 | 1.2 | RF-CL02-99A | 2/2 | 0.91 | 0.91 |
| Cobalt | ND | ND | 1.4 J | 2.1 J | RF-CL02-99A | 2/2 | 1.75 | 1.75 |
| Copper | ND | ND | 9.5 | 10.1 | RF-CL02-99A | 2/2 | 9.8 | 9.8 |
| Iron | ND | ND | 158 | 370 | RF-CL02-99A | 2/2 | 264 | 264 |
| Lead | ND | ND | 0.7 | 1 | RF-CL01-99A | 2/2 | 0.85 | 0.85 |
| Magnesium | ND | ND | 4950 | 6020 | RF-CL02-99A | 2/2 | 5485 | 5485 |
| Manganese | ND | ND | 9 | 9 | RF-CL01-99A,RF-CL02-99A | 2/2 | 9 | 9 |
| Mercury | ND | ND | 0.05 | 0.054 | RF-CL02-99A | 2/2 | 0.05 | 0.05 |
| Nickel | ND | ND | 3.6 | 4 | RF-CL02-99A | 2/2 | 3.8 | 3.8 |
| Potassium | ND | ND | 6240 | 9740 | RF-CL02-99A | 2/2 | 7990 | 7990 |
| Selenium | ND | ND | 2 | 3.1 | RF-CL02-99A | 2/2 | 2.55 | 2.55 |
| Silver | 0.04 U | 1.73 U | ND | ND | | 0/2 | ND | ND |
| Sodium | ND | ND | 39900 | 48900 | RF-CL02-99A | 2/2 | 44400 | 44400 |
| Thallium | 0.11 U | 4.33 U | ND | ND | | 0/2 | ND | ND |
| Vanadium | ND | ND | 2.5 | 3.1 | RF-CL02-99A | 2/2 | 2.8 | 2.8 |
| Zinc | ND | ND | 86.1 | 125 | RF-CL02-99A | 2/2 | 105.55 | 105.55 |

**FREQUENCY OF DETECTIONS IN CLAM TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| | | |
|-------------|-------------|-------------|
| SAMPLE ID | RF-CL01-99A | RF-CL02-99A |
| SAMPLE DATE | 1/16/99 | 1/16/99 |

WET WEIGHT BASIS

| | | |
|--------------------|-----|-----|
| Percent Lipids (%) | 0.2 | 0.1 |
| Moisture (%) | 85 | 88 |

**FREQUENCY OF DETECTIONS IN MUSSEL TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

SAMPLE ID RF-MU02-99A
SAMPLE DATE 1/16/99

INORGANICS (mg/kg)

| | |
|-----------|---------|
| Aluminum | 319 |
| Antimony | 17.43 U |
| Arsenic | 15.2 |
| Barium | 1.2 J |
| Beryllium | 6.7 U |
| Cadmium | 0.21 |
| Calcium | 19800 |
| Chromium | 1.1 |
| Cobalt | 0.34 J |
| Copper | 4.1 |
| Iron | 473 |
| Lead | 1 |
| Magnesium | 3410 |
| Mangancsc | 22.8 |
| Mercury | 0.082 |
| Nickel | 1.1 J |
| Potassium | 6390 |
| Selenium | 3.5 |
| Silver | 5.36 U |
| Sodium | 27200 |
| Thallium | 13.41 U |
| Vanadium | 3.1 |
| Zinc | 38.3 |

**FREQUENCY OF DETECTIONS IN MUSSEL TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| | |
|-------------|-------------|
| SAMPLE ID | RF-MU02-99A |
| SAMPLE DATE | 1/16/99 |

WET WEIGHT BASIS

| | |
|--------------------|-----|
| Percent Lipids (%) | 1.1 |
| Moisture (%) | 32 |

**FREQUENCY OF DETECTIONS IN OYSTER TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID | RF-OY01-99A | RF-OY02-99A |
|---------------------------|-------------|-------------|
| SAMPLE DATE | 1/16/99 | 1/16/99 |
| INORGANICS (mg/kg) | | |
| Aluminum | 104 | 320 |
| Antimony | 0.13 U | 3.88 U |
| Arsenic | 18.8 | 32.2 |
| Barium | 1.2 J | 1.2 J |
| Beryllium | 0.05 U | 1.49 U |
| Cadmium | 0.2 J | 1.8 |
| Calcium | 4390 | 18200 |
| Chromium | 0.85 | 1.5 |
| Cobalt | 2.6 | 0.84 J |
| Copper | 8.2 | 88.2 |
| Iron | 243 | 581 |
| Lead | 2 | 1.2 |
| Magnesium | 5900 | 6150 |
| Manganese | 10.3 | 19.2 |
| Mercury | 0.094 | 0.156 |
| Nickel | 4.5 | 2.1 J |
| Potassium | 7500 | 15200 |
| Selenium | 2.3 | 5.6 |
| Silver | 0.04 U | 1.4 |
| Sodium | 46800 | 51000 |
| Thallium | 0.1 U | 2.99 U |
| Vanadium | 3.3 | 4.6 |
| Zinc | 88.9 | 2230 |

**FREQUENCY OF DETECTIONS IN OYSTER TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| SAMPLE ID SAMPLE DATE | Minimum Non-Detect | Maximum Non-Detect | Minimum Detected | Maximum Detected | Location of Maximum Detect | Frequency of Detection | Arithmetic Mean Positive Detects | Median Positive Detects |
|---------------------------|-----------------------|-----------------------|---------------------|---------------------|-------------------------------|---------------------------|-------------------------------------|----------------------------|
| INORGANICS (mg/kg) | | | | | | | | |
| Aluminum | ND | ND | 104 | 320 | RF-OY02-99A | 2/2 | 212 | 212 |
| Antimony | 0.13 U | 3.88 U | ND | ND | | 0/2 | ND | ND |
| Arsenic | ND | ND | 18.8 | 32.2 | RF-OY02-99A | 2/2 | 25.5 | 25.5 |
| Barium | ND | ND | 1.2 J | 1.2 J | RF-OY01-99A,RF-OY02-99A | 2/2 | 1.2 | 1.2 |
| Beryllium | 0.05 U | 1.49 U | ND | ND | | 0/2 | ND | ND |
| Cadmium | ND | ND | 0.2 J | 1.8 | RF-OY02-99A | 2/2 | 1 | 1 |
| Calcium | ND | ND | 4390 | 18200 | RF-OY02-99A | 2/2 | 11295 | 11295 |
| Chromium | ND | ND | 0.85 | 1.5 | RF-OY02-99A | 2/2 | 1.18 | 1.18 |
| Cobalt | ND | ND | 0.84 J | 2.6 | RF-OY01-99A | 2/2 | 1.72 | 1.72 |
| Copper | ND | ND | 8.2 | 88.2 | RF-OY02-99A | 2/2 | 48.2 | 48.2 |
| Iron | ND | ND | 243 | 581 | RF-OY02-99A | 2/2 | 412 | 412 |
| Lead | ND | ND | 1.2 | 2 | RF-OY01-99A | 2/2 | 1.6 | 1.6 |
| Magnesium | ND | ND | 5900 | 6150 | RF-OY02-99A | 2/2 | 6025 | 6025 |
| Manganese | ND | ND | 10.3 | 19.2 | RF-OY02-99A | 2/2 | 14.75 | 14.75 |
| Mercury | ND | ND | 0.094 | 0.156 | RF-OY02-99A | 2/2 | 0.13 | 0.13 |
| Nickel | ND | ND | 2.1 J | 4.5 | RF-OY01-99A | 2/2 | 3.3 | 3.3 |
| Potassium | ND | ND | 7500 | 15200 | RF-OY02-99A | 2/2 | 11350 | 11350 |
| Selenium | ND | ND | 2.3 | 5.6 | RF-OY02-99A | 2/2 | 3.95 | 3.95 |
| Silver | 0.04 U | 0.04 U | 1.4 | 1.4 | RF-OY02-99A | 1/2 | 1.4 | 1.4 |
| Sodium | ND | ND | 46800 | 51000 | RF-OY02-99A | 2/2 | 48900 | 48900 |
| Thallium | 0.1 U | 2.99 U | ND | ND | | 0/2 | ND | ND |
| Vanadium | ND | ND | 3.3 | 4.6 | RF-OY02-99A | 2/2 | 3.95 | 3.95 |
| Zinc | ND | ND | 88.9 | 2230 | RF-OY02-99A | 2/2 | 1159.45 | 1159.45 |

**FREQUENCY OF DETECTIONS IN OYSTER TISSUE
REFERENCE AREAS
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA**

| | | |
|-------------------------|-------------|-------------|
| SAMPLE ID | RF-OY01-99A | RF-OY02-99A |
| SAMPLE DATE | 1/16/99 | 1/16/99 |
| WET WEIGHT BASIS | | |
| Percent Lipids (%) | 0.1 | 0.6 |
| Moisture (%) | | |

FREQUENCY AND RANGE OF SEDIMENT DATA COMPARED TO SEDIMENT SCREENING LEVELS AND REFERENCE AREAS
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| Analyte | Sediment Screening Values (SSVs) | | Contaminant Frequency/Range | | No. of Positive Detects Above SSV | | Reference Areas | |
|---------------------------|----------------------------------|----------------------|--|------------------------------|-----------------------------------|------|-----------------|--------|
| | SSV/ ER-L ⁽¹⁾ | ER-M ⁽²⁾ | No. of Positive Detects/No. of Samples | Range of Positive Detections | ER-L | ER-M | Range | Mean |
| | | | | | | | | |
| Inorganics (mg/kg) | | | | | | | | |
| Aluminum | NE | NE | 5/5 | 1,580 - 14,800 | NA | NA | 3,000 - 8,160 | 5,580 |
| Arsenic | 7.24 | 70 | 4/5 | 0.84J - 6.8 | 0 | 0 | 2.4 - 6 | 4.2 |
| Barium | 500 ⁽³⁾ | NE | 5/5 | 2.1J - 14.4J | 0 | NA | 3.5J - 8.9J | 6.2 |
| Calcium | NE | NE | 5/5 | 161J - 4,930 | NA | NA | 5,320 - 26,900 | 16,110 |
| Chromium | 52.3 | 370 | 5/5 | 3.2 - 29.8 | 0 | 0 | 7.8 - 19.2 | 13.5 |
| Copper | 18.7 | 270 | 3/5 | 2.4J - 16.6 | 0 | 0 | 4.6J | 4.6J |
| Iron | NE | 27000 ⁽⁴⁾ | 5/5 | 1,480 - 17,400 | NA | 0 | 3,620 - 10,100 | 6,860 |
| Lead | 30.2 | 218 | 5/5 | 2.5 - 20.7 | 0 | 0 | 4.6 - 10.5 | 7.55 |
| Magnesium | NE | NE | 5/5 | 475J - 5,280 | NA | NA | 1,420 - 4,200 | 2,810 |
| Manganese | 460 ⁽⁵⁾ | 1100 ⁽⁵⁾ | 5/5 | 4.2 - 88 | 0 | 0 | 17.7 - 58.7 | 38.2 |
| Nickel | 15.9 | 51.6 | 1/5 | 5.8J | 0 | 0 | 4.7J | 4.7 |
| Potassium | NE | NE | 5/5 | 273J - 2,510J | NA | NA | 605J - 1,500J | 1,053 |
| Silver | 1 | 3.7 | 5/5 | 0.46J - 4.6J | 1 | 1 | 1.5J - 3.2J | 2.35 |
| Sodium | NE | NE | 5/5 | 1,860 - 15,000 | NA | NA | 3,470 - 9,470 | 6,470 |
| Thallium | NE | NE | 1/5 | 1.3J | NA | NA | ND | NA |
| Vanadium | NE | NE | 5/5 | 3.6J - 25.7J | NA | NA | 7.7J - 19.2J | 13.45 |
| Zinc | 124 | 410 | 5/5 | 2.9J - 38.8 | 0 | 0 | 7.9 - 23.8 | 15.85 |

Notes:

J - value reported is estimated
mg/kg - milligrams per kilogram
SSV - Sediment Screening Value
NE - Not Established
ND - Not Detected
NA - Not Applicable

- (1) Region IV Sediment Screening Value (USEPA, 1995), unless otherwise noted
- (2) Long et al. (1995) value, unless otherwise noted
- (3) Sullivan, et al., 1985
- (4) Tetra Tech, 1986 (apparent effects threshold)
- (5) Canadian Screening Value (CMEE, 1993)

FREQUENCY AND RANGE OF MUSSEL TISSUE DATA COMPARED TO REFERENCE AREAS
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| Analyte | Contaminant Frequency/Range | | Reference Area Detection |
|---------------------------|--|------------------------------|--------------------------|
| | No. of Positive Detects/No. of Samples | Range of Positive Detections | |
| Inorganics (mg/kg) | | | |
| Aluminum | 3/3 | 317 - 1,010 | 319 |
| Arsenic | 3/3 | 11 - 37.4 | 15.2 |
| Barium | 3/3 | 0.83J - 2.3J | 1.2J |
| Beryllium | 1/3 | 0.05J | ND |
| Cadmium | 3/3 | 0.19 - 0.52 | 0.21 |
| Calcium | 3/3 | 8,470 - 24,100 | 19,800 |
| Chromium | 3/3 | 0.82 - 2 | 1.1 |
| Cobalt | 3/3 | 0.35J - 0.64J | 0.34J |
| Copper | 3/3 | 4.1 - 12.9 | 4.1 |
| Iron | 3/3 | 366 - 1100 | 473 |
| Lead | 3/3 | 1.3 - 4.8 | 1 |
| Magnesium | 3/3 | 3,010 - 7,950 | 3,410 |
| Manganese | 3/3 | 18.1 - 91.9 | 22.8 |
| Mercury | 3/3 | 0.086 - 0.157 | 0.082 |
| Nickel | 3/3 | 0.73J - 3J | 1.1J |
| Potassium | 3/3 | 6,080 - 18,900 | 6,390 |
| Selenium | 3/3 | 3.3 - 9.4 | 3.5 |
| Sodium | 3/3 | 24,100 - 69,800 | 27,200 |
| Vanadium | 3/3 | 1.9 - 4.9 | 3.1 |
| Zinc | 3/3 | 34.1 - 101 | 38.3 |

Notes:

J - value reported is estimated

mg/kg - milligrams per kilogram

Reference is based on one sample: therefore, the mean value is not calculated.

FREQUENCY AND RANGE OF OYSTER TISSUE DATA COMPARED TO REFERENCE AREAS
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| Analyte | Contaminant Frequency/Range | | Reference Areas | |
|---------------------------|--|------------------------------|-----------------|----------|
| | No. of Positive Detects/No. of Samples | Range of Positive Detections | Reference Areas | |
| | | | Range | Mean |
| Inorganics (mg/kg) | | | | |
| Aluminum | 5/5 | 49.3 - 111 | 104 - 320 | 212 |
| Arsenic | 5/5 | 4.1 - 12.5 | 18.8 - 32.2 | 25.5 |
| Barium | 5/5 | 0.3J - 0.83J | 1.2J | 1.2 |
| Cadmium | 5/5 | 0.6 - 1.7 | 0.2J - 1.8 | 1 |
| Calcium | 5/5 | 2,330 - 7,840 | 4,390 - 18,200 | 11,295 |
| Chromium | 5/5 | 0.43 - 0.72 | 0.85 - 1.5 | 1.18 |
| Cobalt | 3/5 | 0.21J - 0.37J | 0.84J - 2.6 | 1.72 |
| Copper | 5/5 | 17.1 - 50.3 | 8.2 - 88.2 | 48.2 |
| Iron | 5/5 | 123 - 228 | 243 - 581 | 412 |
| Lead | 5/5 | 0.4 - 1.6 | 1.2 - 2 | 1.6 |
| Magnesium | 5/5 | 2,450 - 5,110 | 5,900 - 6,150 | 6,025 |
| Manganese | 5/5 | 5.7 - 11.8 | 10.3 - 19.2 | 14.75 |
| Mercury | 5/5 | 0.053 - 0.252 | 0.094 - 0.156 | 0.13 |
| Nickel | 5/5 | 1.1J - 2.4 | 2.1 - 4.5 | 3.3 |
| Potassium | 5/5 | 5,000 - 11,200 | 7,500 - 15,200 | 11,350 |
| Selenium | 5/5 | 1 - 3.2 | 2.3 - 5.6 | 4 |
| Silver | 1/5 | 0.34J | 1.4 | 1.4 |
| Sodium | 5/5 | 18,600 - 43,800 | 46,800 - 51,000 | 48,900 |
| Vanadium | 5/5 | 0.35J - 1.7J | 3.3 - 4.6 | 3.95 |
| Zinc | 5/5 | 722 - 1,280 | 88.9 - 2,230 | 1,159.50 |

Notes:

J - value reported is estimated
mg/kg - milligrams per kilogram

FREQUENCY AND RANGE OF CLAM TISSUE DATA COMPARED TO REFERENCE AREAS
STUDY AREA
STONE BAY AQUATIC ASSESSMENT, CTO-0100
MCB, CAMP LEJEUNE, NORTH CAROLINA

| Analyte | Contaminant Frequency/Range | | Reference Areas | |
|---------------------------|--|------------------------------|-----------------|--------|
| | No. of Positive Detects/No. of Samples | Range of Positive Detections | Range | Mean |
| Inorganics (mg/kg) | | | | |
| Aluminum | 1/1 | 275 | 64.2 - 249 | 157 |
| Arsenic | 1/1 | 12.4 | 11.1 - 16.2 | 13.65 |
| Barium | 1/1 | 1.7J | 0.99J - 1.7J | 1.35 |
| Cadmium | 1/1 | 0.47 | 0.21 - 0.23J | 0.22 |
| Calcium | 1/1 | 12,400 | 3,840 - 5,720 | 4,780 |
| Chromium | 1/1 | 1.2 | 0.62 - 1.2 | 0.91 |
| Cobalt | 1/1 | 2.3J | 1.4J - 2.1J | 1.75 |
| Copper | 1/1 | 12.6 | 9.5 - 10.1 | 9.8 |
| Iron | 1/1 | 442 | 158 - 370 | 264 |
| Lead | 1/1 | 1.8 | 0.7 - 1 | 0.85 |
| Magnesium | 1/1 | 8,370 | 4,950 - 6,020 | 5,485 |
| Manganese | 1/1 | 33.5 | 9 | 9 |
| Mercury | 1/1 | 0.05 | 0.05 - 0.054 | 0.05 |
| Nickel | 1/1 | 4.1 | 3.6 - 4 | 3.8 |
| Potassium | 1/1 | 11,600 | 6,240 - 9,740 | 7,990 |
| Selenium | 1/1 | 2.8 | 2 - 3.1 | 3 |
| Sodium | 1/1 | 71,900 | 39,990 - 48,900 | 44,400 |
| Vanadium | 1/1 | 1.9J | 2.5 - 3.1 | 2.8 |
| Zinc | 1/1 | 117 | 86.1 - 125 | 105.55 |

Notes:

Clams were only obtained from one study area sampling station.

J - value reported is estimated

mg/kg - milligrams per kilogram

Baker

Baker Environmental, Inc.
ATTACHMENT C

Form I's

STONE BAY

SEDIMENT - INORGANICS

TOTAL METALS

1

EPA SAMPLE NO.

INORGANIC ANALYSIS DATA SHEET

RF-SD01-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC

Case No.: CTO 100 SAS No.:

SDG No.: Y0199A

Matrix (soil/water): SOIL

Lab Sample ID: 990034-19 S

Level (low/med): LOW

Date Received: 01/19/99

% Solids: 73.0

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| ICAS No. | Analyte | Concentration | CI | Q | IM |
|------------|-----------|---------------|----|---|------|
| 17429-90-5 | Aluminum | 3000 | | * | IP |
| 17440-36-0 | Antimony | 0.59 | UI | N | IP |
| 17440-38-2 | Arsenic | 2.4 | UI | | IP |
| 17440-39-3 | Barium | 3.5 | UI | | IP |
| 17440-41-7 | Beryllium | 0.23 | UI | | IP |
| 17440-43-9 | Cadmium | 0.09 | UI | | IP |
| 17440-70-2 | Calcium | 5320 | UI | | IP |
| 17440-47-3 | Chromium | 7.8 | UI | | IP |
| 17440-48-4 | Cobalt | 1.2 | UI | | IP |
| 17440-50-8 | Copper | 2.4 | UI | | IP |
| 17439-89-6 | Iron | 3620 | UI | * | IP |
| 17439-92-1 | Lead | 4.6 | UI | * | IP |
| 17439-95-4 | Magnesium | 1420 | UI | | IP |
| 17439-96-5 | Manganese | 17.7 | UI | | IP |
| 17439-97-6 | Mercury | 0.06 | UI | | IAVI |
| 17440-02-0 | Nickel | 2.3 | UI | | IP |
| 17440-09-7 | Potassium | 605 | UI | | IP |
| 17782-49-2 | Selenium | 0.73 | UI | N | IP |
| 17440-22-4 | Silver | 1.5 | UI | | IP |
| 17440-23-5 | Sodium | 3470 | UI | | IP |
| 17440-28-0 | Thallium | 0.46 | UI | | IP |
| 17440-62-2 | Vanadium | 7.7 | UI | | IP |
| 17440-66-6 | Zinc | 7.9 | UI | | IP |

Color Before: BROWN

Clarity Before:

Texture: MEDIUM

Color After: COLORLESS

Clarity After:

Artifacts:

Comments:

TOTAL METALS

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

RF-SD02-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC

Case No.: CTO 100

SAS No.:

SDG No.: Y0199A

Matrix (soil/water): SOIL

Lab Sample ID: 990034-20 S

Level (low/med): LOW

Date Received: 01/19/99

% Solids: 49.2

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| ICAS No. | Analyte | Concentration | CI | Q | IM |
|------------|-----------|---------------|----|---|------|
| 17429-90-5 | Aluminum | 8160 | | * | IP |
| 17440-36-0 | Antimony | 1.0 | UI | N | IP |
| 17440-38-2 | Arsenic | 6.0 | | | IP |
| 17440-39-3 | Barium | 8.9 | BI | | IP |
| 17440-41-7 | Beryllium | 0.39 | UI | | IP |
| 17440-43-9 | Cadmium | 0.15 | UI | | IP |
| 17440-70-2 | Calcium | 26900 | | | IP |
| 17440-47-3 | Chromium | 19.2 | | | IP |
| 17440-48-4 | Cobalt | 2.1 | BI | | IP |
| 17440-50-8 | Copper | 4.6 | BI | | IP |
| 17439-89-6 | Iron | 10100 | | * | IP |
| 17439-92-1 | Lead | 10.5 | | * | IP |
| 17439-95-4 | Magnesium | 4200 | | | IP |
| 17439-96-5 | Manganese | 58.7 | | | IP |
| 17439-97-6 | Mercury | 0.09 | UI | | IAVI |
| 17440-02-0 | Nickel | 4.7 | BI | | IP |
| 17440-09-7 | Potassium | 1500 | BI | | IP |
| 17782-49-2 | Selenium | 1.6 | BI | N | IP |
| 17440-22-4 | Silver | 3.2 | BI | | IP |
| 17440-23-5 | Sodium | 9470 | | | IP |
| 17440-28-0 | Thallium | 0.77 | UI | | IP |
| 17440-62-2 | Vanadium | 19.2 | BI | | IP |
| 17440-66-6 | Zinc | 23.8 | | | IP |

Color Before: BROWN

Clarity Before:

Texture: MEDIUM

Color After: COLORLESS

Clarity After:

Artifacts:

Comments:

130

TOTAL METALS

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

SB-SD01-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC

Case No.: CTO 100

SAS No.:

SDG No.: Y0199A

Matrix (soil/water): SOIL

Lab Sample ID: 990034-14 S

Level (low/med): LOW

Date Received: 01/19/99

% Solids: 76.4

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| ICAS No. | Analyte | Concentration | Q | IM |
|------------|-----------|---------------|---|-----|
| 17429-90-5 | Aluminum | 1750 | * | IP |
| 17440-36-0 | Antimony | 0.66 | N | IP |
| 17440-38-2 | Arsenic | 1.1 | | IP |
| 17440-39-3 | Barium | 2.9 | | IP |
| 17440-41-7 | Beryllium | 0.26 | | IP |
| 17440-43-9 | Cadmium | 0.10 | | IP |
| 17440-70-2 | Calcium | 484 | | IP |
| 17440-47-3 | Chromium | 5.6 | | IP |
| 17440-48-4 | Cobalt | 1.4 | | IP |
| 17440-50-8 | Copper | 2.7 | | IP |
| 17439-89-6 | Iron | 2380 | * | IP |
| 17439-92-1 | Lead | 6.0 | * | IP |
| 17439-95-4 | Magnesium | 843 | | IP |
| 17439-96-5 | Manganese | 8.9 | | IP |
| 17439-97-6 | Mercury | 0.05 | | AVI |
| 17440-02-0 | Nickel | 2.5 | | IP |
| 17440-09-7 | Potassium | 430 | | IP |
| 17782-49-2 | Selenium | 0.81 | N | IP |
| 17440-22-4 | Silver | 0.75 | | IP |
| 17440-23-5 | Sodium | 3980 | | IP |
| 17440-28-0 | Thallium | 0.51 | | IP |
| 17440-62-2 | Vanadium | 5.3 | | IP |
| 17440-66-6 | Zinc | 5.8 | | IP |

Color Before: BROWN

Clarity Before:

Texture: MEDIUM

Color After: COLORLESS

Clarity After:

Artifacts:

Comments:

TOTAL METALS

1

EPA SAMPLE NO.

INORGANIC ANALYSIS DATA SHEET

SB-SD02-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC

Case No.: CTO 100 SAS No.:

SDG No.: Y0199A

Matrix (soil/water): SOIL

Lab Sample ID: 990034-15 S

Level (low/med): LOW

Date Received: 01/19/99

* Solids: 74.6

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| ICAS No. | Analyte | Concentration | CI | Q | IM |
|------------|-----------|---------------|----|---|------|
| 17429-90-5 | Aluminum | 1780 | | * | IP |
| 17440-36-0 | Antimony | 0.61 | UI | N | IP |
| 17440-38-2 | Arsenic | 0.85 | IB | | IP |
| 17440-39-3 | Barium | 2.1 | IB | | IP |
| 17440-41-7 | Beryllium | 0.24 | UI | | IP |
| 17440-43-9 | Cadmium | 0.09 | UI | | IP |
| 17440-70-2 | Calcium | 161 | IB | | IP |
| 17440-47-3 | Chromium | 4.2 | II | | IP |
| 17440-48-4 | Cobalt | 1.3 | UI | | IP |
| 17440-50-8 | Copper | 16.6 | II | | IP |
| 17439-89-6 | Iron | 2120 | II | * | IP |
| 17439-92-1 | Lead | 5.7 | II | * | IP |
| 17439-95-4 | Magnesium | 592 | IB | | IP |
| 17439-96-5 | Manganese | 6.3 | II | | IP |
| 17439-97-6 | Mercury | 0.06 | UI | | IAVI |
| 17440-02-0 | Nickel | 2.3 | UI | | IP |
| 17440-09-7 | Potassium | 341 | IB | | IP |
| 17782-49-2 | Selenium | 0.75 | UI | N | IP |
| 17440-22-4 | Silver | 0.90 | IB | | IP |
| 17440-23-5 | Sodium | 2530 | II | | IP |
| 17440-28-0 | Thallium | 0.47 | UI | | IP |
| 17440-62-2 | Vanadium | 4.5 | IB | | IP |
| 17440-66-6 | Zinc | 5.6 | II | | IP |

Color Before: BROWN

Clarity Before:

Texture: MEDIUM

Color After: COLORLESS

Clarity After:

Artifacts:

Comments:

132

TOTAL METALS

1

EPA SAMPLE NO.

INORGANIC ANALYSIS DATA SHEET

SB-SD03-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC

Case No.: CTO 100

SAS No.:

SDG No.: Y0199A

Matrix (soil/water): SOIL

Lab Sample ID: 990034-16 S

Level (low/med): LOW

Date Received: 01/19/99

% Solids:

82.9

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| ICAS No. | Analyte | Concentration | Q | IM |
|------------|-----------|---------------|---|-----|
| 17429-90-5 | Aluminum | 1580 | * | IP |
| 17440-36-0 | Antimony | 0.571 | N | IP |
| 17440-38-2 | Arsenic | 0.681 | | IP |
| 17440-39-3 | Barium | 2.2 | | IP |
| 17440-41-7 | Beryllium | 0.221 | | IP |
| 17440-43-9 | Cadmium | 0.091 | | IP |
| 17440-70-2 | Calcium | 323 | | IP |
| 17440-47-3 | Chromium | 3.2 | | IP |
| 17440-48-4 | Cobalt | 1.2 | | IP |
| 17440-50-8 | Copper | 2.4 | | IP |
| 17439-89-6 | Iron | 1480 | * | IP |
| 17439-92-1 | Lead | 2.5 | * | IP |
| 17439-95-4 | Magnesium | 511 | | IP |
| 17439-96-5 | Manganese | 4.2 | | IP |
| 17439-97-6 | Mercury | 0.061 | | AVI |
| 17440-02-0 | Nickel | 2.2 | | IP |
| 17440-09-7 | Potassium | 285 | | IP |
| 17782-49-2 | Selenium | 0.701 | N | IP |
| 17440-22-4 | Silver | 0.461 | | IP |
| 17440-23-5 | Sodium | 2080 | | IP |
| 17440-28-0 | Thallium | 0.441 | | IP |
| 17440-62-2 | Vanadium | 3.7 | | IP |
| 17440-66-6 | Zinc | 2.9 | | IP |

Color Before: BROWN

Clarity Before:

Texture: MEDIUM

Color After: COLORLESS

Clarity After:

Artifacts:

Comments:

133

TOTAL METALS

1

EPA SAMPLE NO.

INORGANIC ANALYSIS DATA SHEET

SB-SD04-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC

Case No.: CTD 100

SAS No.:

SDG No.: Y0199A

Matrix (soil/water): SOIL

Lab Sample ID: 990034-17 S

Level (low/med): LOW

Date Received: 01/19/99

% Solids: 35.8

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| ICAS No. | Analyte | Concentration | CI | Q | IM |
|------------|-----------|---------------|----|---|------|
| 17429-90-5 | Aluminum | 14800 | | * | IP |
| 17440-36-0 | Antimony | 1.4 | UI | N | IP |
| 17440-38-2 | Arsenic | 6.8 | | | IP |
| 17440-39-3 | Barium | 14.4 | BI | | IP |
| 17440-41-7 | Beryllium | 0.53 | UI | | IP |
| 17440-43-9 | Cadmium | 0.21 | UI | | IP |
| 17440-70-2 | Calcium | 4930 | | | IP |
| 17440-47-3 | Chromium | 29.8 | | | IP |
| 17440-48-4 | Cobalt | 2.8 | UI | | IP |
| 17440-50-8 | Copper | 8.7 | BI | | IP |
| 17439-89-6 | Iron | 17400 | | * | IP |
| 17439-92-1 | Lead | 20.7 | | * | IP |
| 17439-95-4 | Magnesium | 5280 | | | IP |
| 17439-96-5 | Manganese | 88.0 | | | IP |
| 17439-97-6 | Mercury | 0.13 | UI | | IAVI |
| 17440-02-0 | Nickel | 5.8 | BI | | IP |
| 17440-09-7 | Potassium | 2510 | BI | | IP |
| 17782-49-2 | Selenium | 1.7 | UI | N | IP |
| 17440-22-4 | Silver | 4.6 | BI | | IP |
| 17440-23-5 | Sodium | 15000 | | | IP |
| 17440-28-0 | Thallium | 1.3 | BI | | IP |
| 17440-62-2 | Vanadium | 25.7 | BI | | IP |
| 17440-66-6 | Zinc | 38.8 | | | IP |

Color Before: BROWN

Clarity Before:

Texture: MEDIUM

Color After: COLORLESS

Clarity After:

Artifacts:

Comments:

134

TOTAL METALS

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

| |
|-------------|
| SB-SD05-99A |
|-------------|

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC

Case No.: CTO 100 SAS No.:

SDG No.: Y0199A

Matrix (soil/water): SOIL

Lab Sample ID: 990034-18 S

Level (low/med): LOW

Date Received: 01/19/99

% Solids: 79.1

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| ICAS No. | Analyte | Concentration | Cl | Q | IM |
|------------|-----------|---------------|----|---|-----|
| 17429-90-5 | Aluminum | 1580 | | * | IP |
| 17440-36-0 | Antimony | 0.58 | U | N | IP |
| 17440-38-2 | Arsenic | 0.84 | B | | IP |
| 17440-39-3 | Barium | 2.7 | B | | IP |
| 17440-41-7 | Beryllium | 0.23 | U | | IP |
| 17440-43-9 | Cadmium | 0.09 | U | | IP |
| 17440-70-2 | Calcium | 1320 | | | IP |
| 17440-47-3 | Chromium | 3.9 | | | IP |
| 17440-48-4 | Cobalt | 1.2 | U | | IP |
| 17440-50-8 | Copper | 2.3 | U | | IP |
| 17439-89-6 | Iron | 1660 | | * | IP |
| 17439-92-1 | Lead | 3.4 | | * | IP |
| 17439-95-4 | Magnesium | 475 | B | | IP |
| 17439-96-5 | Manganese | 8.7 | | | IP |
| 17439-97-6 | Mercury | 0.06 | U | | IAV |
| 17440-02-0 | Nickel | 2.2 | U | | IP |
| 17440-09-7 | Potassium | 273 | B | | IP |
| 17782-49-2 | Selenium | 0.72 | U | N | IP |
| 17440-22-4 | Silver | 0.60 | B | | IP |
| 17440-23-5 | Sodium | 1860 | | | IP |
| 17440-28-0 | Thallium | 0.45 | U | | IP |
| 17440-62-2 | Vanadium | 3.6 | B | | IP |
| 17440-66-6 | Zinc | 3.3 | B | | IP |

Color Before: BROWN

Clarity Before:

Texture: MEDIUM

Color After: COLORLESS

Clarity After:

Artifacts:

Comments:

135

STONE BAY

SEDIMENT - TOTAL ORGANIC CARBON

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL ORGANIC CARBON

EPA 415.1/9060

Client: Baker Environmental

Client Sample ID: RF-SD01-99A

Date Sampled: 01/16/99

Laboratory ID: 990034-19

Date Sample Received: 01/19/99

Percent Solids: 73.0

Matrix: Sediment

| Target Analyte | Result | Units | Method Reporting Limit | Date Prep'd | Date Analyzed |
|----------------------|--------|---------|------------------------|-------------|---------------|
| Total Organic Carbon | 4930 | mg/Kg + | 68.5 | 01/25/99 | 01/25/99 |

+ Dry weight basis.

Reported by: _____

Jeffrey D. Maymon

Approved by: _____

Donald Totozelli

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL ORGANIC CARBON

EPA 415.1/9060

Client: Baker Environmental

Client Sample ID: RF-SD02-99A

Date Sampled: 01/16/99

Laboratory ID: 990034-20

Date Sample Received: 01/19/99

Percent Solids: 49.2

Matrix: Sediment

| Target Analyte | Result | Units | Method Reporting Limit | Date Prep'd | Date Analyzed |
|----------------------|--------|--------|------------------------|-------------|---------------|
| Total Organic Carbon | 12400 | mg/Kg+ | 102 | 01/25/99 | 01/25/99 |

+ Dry weight basis.

Reported by: _____

Jeffrey D. Maymon

Approved by: _____

Donald Zatoelli

STONE BAY

**SEDIMENT - GEOTECHNICAL
LABORATORY TEST DATA**



GeoTesting Express

1145 MASSACHUSETTS AVE.
BOXBOROUGH, MA 01719
978-635-0424 (FAX) 978-635-0266

January 26, 1999

Ms. Peg Marple
Ceimic Corporation
10 Dean Knauss Drive
South Ferry Industrial Park
Narragansett, RI 02882

RE: Baker Environmental (GTX-2130)

Dear Ms. Marple:

Enclosed are the test results you requested for the above referenced project. We received the following three soil samples from you on January 22, 1999:

RF-SD01-99A
RF-SD02-99A
SB-SD01-99A

A copy of the chain of custody form for these samples is located at the back of this report. GeoTesting Express performed one sieve analysis (ASTM D 422) and one Atterberg limits (ASTM D 4318) on each of these samples. Two of the samples were determined to be non-plastic.


The results presented in this report apply only to the items tested. This report shall not be reproduced except in full, without written approval from GeoTesting Express. The remainder of these samples will be retained for a period of sixty days and will then be discarded unless otherwise notified by you. Please call me directly if you have any questions or require additional information. Thank you for allowing us this opportunity to once again provide your firm with physical properties testing of soils. We look forward to working with you again in the future.

Respectfully yours;

Gary T. Torosian
Laboratory Manager

Boring No. : ---
 Sample No.: RF-SD01-99A
 Test Method ASTM D 422
 Filename : RFSD0199

Project : Baker Environmental
 Project No.: GTX-2130
 Location: ---
 Date : Tue Jan 26 1999



| | | | | | | |
|---------|--------|------|--------|--------|------|--------------|
| COBBLES | GRAVEL | | SAND | | | SILT OR CLAY |
| | COARSE | FINE | COARSE | MEDIUM | FINE | |

Classification :
 (SM) Silty sand
 Visual Description :
 Wet, dark greenish gray silty sand

Remarks :
 Hydrometer not requested

Figure 1

GEOTECHNICAL LABORATORY TEST DATA

Project : Baker Environmental
 Project No. : GTX-2130
 Boring No. : ---
 Sample No. : RF-SD01-99A
 Location : ---
 Soil Description : Wet, dark greenish gray silty sand
 Remarks : Hydrometer not requested

Depth : ---
 Test Date : 01/25/99
 Test Method : ASTM D 422

Filename : RFSD0199
 Elevation : ---
 Tested by : tje
 Checked by : gtt

| Sieve Mesh | Sieve Openings | | FINE SIEVE SET | | Percent Finer (%) |
|------------|----------------|-------------|----------------------|---------------------------------|-------------------|
| | Inches | Millimeters | Weight Retained (gm) | Cumulative Weight Retained (gm) | |
| #4 | 0.187 | 4.75 | 0.00 | 0.00 | 100 |
| #10 | 0.079 | 2.00 | 0.17 | 0.17 | 100 |
| #20 | 0.033 | 0.84 | 0.31 | 0.48 | 100 |
| #40 | 0.017 | 0.42 | 0.25 | 0.73 | 99 |
| #60 | 0.010 | 0.25 | 4.59 | 5.32 | 96 |
| #100 | 0.006 | 0.15 | 80.33 | 85.65 | 31 |
| #200 | 0.003 | 0.07 | 21.24 | 106.89 | 14 |
| Pan | | | 16.89 | 123.78 | 0 |

Total Dry Weight of Sample = 133.01

- D85 : 0.2295 mm
- D60 : 0.1881 mm
- D50 : 0.1736 mm
- D30 : 0.1442 mm
- D15 : 0.0782 mm
- D10 : 0.0638 mm

Soil Classification

ASTM Group Symbol : SM
 ASTM Group Name : Silty sand
 AASHTO Group Symbol : A-2-4(0)
 AASHTO Group Name : Silty Gravel and Sand

ATTERBERG LIMITS

| | | | |
|--|----------------------------|-------------------------|------------------------------|
| PROJECT Baker Environmental | PROJECT NUMBER GTX-2130 | TESTED BY tje | BORING NUMBER --- |
| LOCATION --- | | CHECKED BY gtt | SAMPLE NUMBER RF-SD01-99A |
| SAMPLE DESCRIPTION Wet, dark greenish gray silty sand | | DATE Tue Jan 26 1999 | FILENAME RFSD0199 |

LIQUID LIMIT DETERMINATIONS

| | | | | |
|----------------------------|--|--|--|--|
| CONTAINER NUMBER | | | | |
| WT. WET SOIL + TARE | | | | |
| WT. DRY SOIL + TARE | | | | |
| WT. WATER | | | | |
| TARE WT. | | | | |
| WT. DRY SOIL | | | | |
| WATER CONTENT, W_N (%) | | | | |
| NUMBER OF BLOWS, N | | | | |
| ONE-POINT LIQUID LIMIT, LL | | | | |

PLASTIC LIMIT DETERMINATIONS

| | | | | |
|---------------------|--|--|--|--|
| CONTAINER NUMBER | | | | |
| WT. WET SOIL + TARE | | | | |
| WT. DRY SOIL + TARE | | | | |
| WT. WATER | | | | |
| TARE WT. | | | | |
| WT. DRY SOIL | | | | |
| WATER CONTENT (%) | | | | |

SUMMARY OF RESULTS

| | |
|------------------------------|------|
| NATURAL WATER CONTENT, W (%) | 34.2 |
| LIQUID LIMIT, LL | |
| PLASTIC LIMIT, PL | |
| PLASTICITY INDEX, PI | |
| LIQUIDITY INDEX, LI* | |

*LI = (W - PL)/PI

PLASTICITY CHART





Fig. 1.0

Boring No. : ---
 Sample No.: RF-SD02-99A
 Test Method ASTM D 422
 Filename : RFSD0299

Project : Baker Environmental
 Project No.: GTX-2130
 Location: ---
 Date : Tue Jan 26 1999



| | | | | | | |
|---------|--------|------|--------|--------|------|--------------|
| COBBLES | GRAVEL | | SAND | | | SILT OR CLAY |
| | COARSE | FINE | COARSE | MEDIUM | FINE | |

Classification :
 (CH) Sandy fat clay
 Visual Description :
 Very wet, dark greenish gray sandy clay w/ shells

Remarks :
 Hydrometer not requested

Figure 2

GEOTECHNICAL LABORATORY TEST DATA

Project : Baker Environmental
 Project No. : GTX-2130 Depth : ---
 Boring No. : --- Test Date : 01/25/99
 Sample No. : RF-SD02-99A Test Method : ASTM D 422
 Location : ---
 Soil Description : Very wet, dark greenish gray sandy clay w/ shells
 Remarks : Hydrometer not requested

Filename : RFSD0299
 Elevation : ---
 Tested by : tje
 Checked by : gtt

| Sieve Mesh | Sieve Openings | | FINE SIEVE SET | | Percent Finer (%) |
|------------|----------------|-------------|----------------------|---------------------------------|-------------------|
| | Inches | Millimeters | Weight Retained (gm) | Cumulative Weight Retained (gm) | |
| 0.375" | 0.374 | 9.51 | 0.00 | 0.00 | 100 |
| #4 | 0.187 | 4.75 | 0.11 | 0.11 | 100 |
| #10 | 0.079 | 2.00 | 0.39 | 0.50 | 99 |
| #20 | 0.033 | 0.84 | 0.34 | 0.84 | 98 |
| #40 | 0.017 | 0.42 | 0.34 | 1.18 | 98 |
| #60 | 0.010 | 0.25 | 1.82 | 3.00 | 94 |
| #100 | 0.006 | 0.15 | 7.87 | 10.87 | 78 |
| #200 | 0.003 | 0.07 | 12.10 | 22.97 | 54 |
| Pan | | | 27.25 | 50.22 | 0 |

Total Dry Weight of Sample = 59.62


D85 : 0.1856 mm
 D60 : 0.0874 mm
 D50 : N/A
 D30 : N/A
 D15 : N/A
 D10 : N/A

Soil Classification


ASTM Group Symbol : CH
 ASTM Group Name : Sandy fat clay
 AASHTO Group Symbol : A-7-6(24)
 AASHTO Group Name : Clayey Soils

ATTERBERG LIMITS

| PROJECT Baker Environmental | PROJECT NUMBER GTX-2130 | TESTED BY tje | BORING NUMBER --- |
|---|----------------------------|------------------------------|----------------------|
| LOCATION --- | CHECKED BY glt | SAMPLE NUMBER RF-SD02-99A | |
| SAMPLE DESCRIPTION Very wet, dark greenish gray sandy clay w/ shells | | DATE Tue Jan 26 1999 | FILENAME RFSD0299 |
| LIQUID LIMIT DETERMINATIONS | | | |
| CONTAINER NUMBER | bk17 | bk33 | bk137 |
| WT. WET SOIL + TARE | 37.14 | 39.16 | 37.22 |
| WT. DRY SOIL + TARE | 33.94 | 35.63 | 34.27 |
| WT. WATER | 3.2 | 3.53 | 2.95 |
| TARE WT. | 29.37 | 30.13 | 29.39 |
| WT. DRY SOIL | 4.57 | 5.5 | 4.88 |
| WATER CONTENT, W_N (%) | 70.02 | 64.18 | 60.45 |
| NUMBER OF BLOWS, N | 13 | 23 | 34 |
| ONE-POINT LIQUID LIMIT, LL | 64.69 | 63.54 | 62.74 |
| PLASTIC LIMIT DETERMINATIONS | | | |
| CONTAINER NUMBER | bk76 | bk144 | |
| WT. WET SOIL + TARE | 34.24 | 37.63 | |
| WT. DRY SOIL + TARE | 33.37 | 36.19 | |
| WT. WATER | 0.87 | 1.44 | |
| TARE WT. | 29.19 | 29.61 | |
| WT. DRY SOIL | 4.18 | 6.58 | |
| WATER CONTENT (%) | 20.81 | 21.88 | |




| SUMMARY OF RESULTS | |
|------------------------------|------|
| NATURAL WATER CONTENT, W (%) | 69.1 |
| LIQUID LIMIT, LL | 63.5 |
| PLASTIC LIMIT, PL | 21.3 |
| PLASTICITY INDEX, PI | 42.1 |
| LIQUIDITY INDEX, LI* | 1.13 |



Boring No. : ---
 Sample No.: SB-SD01-99A
 Test Method ASTM D 422
 Filename : SBS0199

Project : Baker Environmental
 Project No.: GTX-2130
 Location: ---
 Date : Tue Jan 26 1999



| | | | | | | |
|---------|--------|------|--------|--------|------|--------------|
| COBBLES | GRAVEL | | SAND | | | SILT OR CLAY |
| | COARSE | FINE | COARSE | MEDIUM | FINE | |

Classification :
 (SP-SM) Poorly graded sand with silt
 Visual Description :
 Wet, dark greenish gray sand with silt

Remarks :

Figure 3

GEOTECHNICAL LABORATORY TEST DATA

Project : Baker Environmental
 Project No. : GTX-2130 Depth : ---
 Boring No. : --- Test Date : 01/25/99
 Sample No. : SB-SD01-99A Test Method : ASTM D 422
 Location : ---
 Soil Description : Wet, dark greenish gray sand with silt
 Remarks : ---

Filename : SBS0199
 Elevation : ---
 Tested by : tje
 Checked by : gtt

| Sieve Mesh | Sieve Openings | | FINE SIEVE SET | | Percent Finer (%) |
|------------|----------------|-------------|----------------------|---------------------------------|-------------------|
| | Inches | Millimeters | Weight Retained (gm) | Cumulative Weight Retained (gm) | |
| #4 | 0.187 | 4.75 | 0.00 | 0.00 | 100 |
| #10 | 0.079 | 2.00 | 0.08 | 0.08 | 100 |
| #20 | 0.033 | 0.84 | 0.08 | 0.16 | 100 |
| #40 | 0.017 | 0.42 | 0.29 | 0.45 | 100 |
| #60 | 0.010 | 0.25 | 12.89 | 13.34 | 86 |
| #100 | 0.006 | 0.15 | 38.42 | 51.76 | 47 |
| #200 | 0.003 | 0.07 | 36.88 | 88.64 | 8 |
| Pan | | | 8.22 | 96.86 | 0 |

Total Dry Weight of Sample = 106.28


- D85 : 0.2460 mm
- D60 : 0.1776 mm
- D50 : 0.1558 mm
- D30 : 0.1099 mm
- D15 : 0.0834 mm
- D10 : 0.0761 mm

Soil Classification

ASTM Group Symbol : SP-SM
 ASTM Group Name : Poorly graded sand with silt
 AASHTO Group Symbol : A-3(0)
 AASHTO Group Name : Fine Sand

ATTERBERG LIMITS

| | | | |
|--|----------------------------|------------------------------|------------------------------|
| PROJECT Baker Environmental | PROJECT NUMBER GTX-2130 | TESTED BY tje | BORING NUMBER --- |
| LOCATION --- | | CHECKED BY gtt | SAMPLE NUMBER SB-SD01-99A |
| SAMPLE DESCRIPTION Wet, dark greenish gray sand with silt | | DATE Tue Jan 26 1999 | FILENAME SBSD0199 |
| LIQUID LIMIT DETERMINATIONS | | | |
| CONTAINER NUMBER | | | |
| WT. WET SOIL + TARE | | | |
| WT. DRY SOIL + TARE | | | |
| WT. WATER | | | |
| TARE WT. | | | |
| WT. DRY SOIL | | | |
| WATER CONTENT, W_N (%) | | | |
| NUMBER OF BLOWS, N | | | |
| ONE-POINT LIQUID LIMIT, LL | | | |
| PLASTIC LIMIT DETERMINATIONS | | | |
| CONTAINER NUMBER | | | |
| WT. WET SOIL + TARE | | | |
| WT. DRY SOIL + TARE | | | |
| WT. WATER | | | |
| TARE WT. | | | |
| WT. DRY SOIL | | | |
| WATER CONTENT (%) | | | |
| | | SUMMARY OF RESULTS | |
| | | NATURAL WATER CONTENT, W (%) | 27.9 |
| | | LIQUID LIMIT, LL | |
| | | PLASTIC LIMIT, PL | |
| | | PLASTICITY INDEX, PI | |
| | | LIQUIDITY INDEX, LI* | |
| | | *LI = (W - PL)/PI | |



STONE BAY

**CLAM, MUSSEL, AND
OYSTER TISSUE - INORGANICS**

TOTAL METALS TISSUE

1

EPA SAMPLE NO.

INORGANIC ANALYSIS DATA SHEET

RF-CL01-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC

Case No.: CTO 100 SAS No.:

SDG No.: TY0199A

Matrix (soil/water): SOIL

Lab Sample ID: T990034-08 S

Level (low/med): LOW

Date Received: 01/19/99

* Solids: 15.4

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| CAS No. | Analyte | Concentration | Q | IM |
|------------|-----------|---------------|-----|----|
| 17429-90-5 | Aluminum | 64.2 | | IP |
| 17440-36-0 | Antimony | 0.10 | | IP |
| 17440-38-2 | Arsenic | 11.1 | *E | IP |
| 17440-39-3 | Barium | 0.99 | | IP |
| 17440-41-7 | Beryllium | 0.04 | | IP |
| 17440-43-9 | Cadmium | 0.21 | | IP |
| 17440-70-2 | Calcium | 3840 | *E | IP |
| 17440-47-3 | Chromium | 0.62 | | IP |
| 17440-48-4 | Cobalt | 1.4 | | IP |
| 17440-50-8 | Copper | 9.5 | | IP |
| 17439-89-6 | Iron | 158 | E | IP |
| 17439-92-1 | Lead | 1.0 | N | IP |
| 17439-95-4 | Magnesium | 4950 | E | IP |
| 17439-96-5 | Manganese | 9.0 | N*E | IP |
| 17440-02-0 | Nickel | 3.6 | | IP |
| 17440-09-7 | Potassium | 6240 | E | IP |
| 17782-49-2 | Selenium | 2.0 | N | IP |
| 17440-22-4 | Silver | 0.03 | N | IP |
| 17440-23-5 | Sodium | 39900 | | IP |
| 17440-28-0 | Thallium | 0.08 | | IP |
| 17440-62-2 | Vanadium | 2.5 | | IP |
| 17440-66-6 | Zinc | 86.1 | NE | IP |

Color Before: GREY

Clarity Before:

Texture: MEDIUM

Color After: YELLOW

Clarity After:

Artifacts:

Comments:

TOTAL METALS TISSUE

1

EPA SAMPLE NO.

INORGANIC ANALYSIS DATA SHEET

| |
|-------------|
| RF-CL02-99A |
|-------------|

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC Case No.: CTO 100 SAS No.: SDG No.: TY0199A

Matrix (soil/water): SOIL Lab Sample ID: T990034-21 S

Level (low/med): LOW Date Received: 01/19/99

% Solids: 11.5

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| CAS No. | Analyte | Concentration | CI | Q | IM |
|------------|-----------|---------------|----|-----|----|
| 17429-90-5 | Aluminum | 249 | | | IP |
| 17440-36-0 | Antimony | 0.14 | UI | | IP |
| 17440-38-2 | Arsenic | 16.2 | | *E | IP |
| 17440-39-3 | Barium | 1.7 | BI | | IP |
| 17440-41-7 | Beryllium | 0.05 | UI | | IP |
| 17440-43-9 | Cadmium | 0.23 | BI | | IP |
| 17440-70-2 | Calcium | 5720 | | *E | IP |
| 17440-47-3 | Chromium | 1.2 | | | IP |
| 17440-48-4 | Cobalt | 2.1 | BI | | IP |
| 17440-50-8 | Copper | 10.1 | | | IP |
| 17439-89-6 | Iron | 370 | | E | IP |
| 17439-92-1 | Lead | 0.70 | | N | IP |
| 17439-95-4 | Magnesium | 6020 | | E | IP |
| 17439-96-5 | Manganese | 9.0 | | N*E | IP |
| 17440-02-0 | Nickel | 4.0 | | | IP |
| 17440-09-7 | Potassium | 9740 | | E | IP |
| 17782-49-2 | Selenium | 3.1 | | N | IP |
| 17440-22-4 | Silver | 0.04 | UI | N | IP |
| 17440-23-5 | Sodium | 48900 | | | IP |
| 17440-28-0 | Thallium | 0.11 | UI | | IP |
| 17440-62-2 | Vanadium | 3.1 | | | IP |
| 17440-66-6 | Zinc | 125 | | NE | IP |

Color Before: GREY Clarity Before: Texture: MEDIUM

Color After: YELLOW Clarity After: Artifacts:

Comments:

TOTAL METALS TISSUE

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

SB-CL04-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC

Case No.: CTO 100 SAS No.:

SDG No.: TY0199A

Matrix (soil/water): SOIL

Lab Sample ID: T990034-11 S

Level (low/med): LOW

Date Received: 01/19/99

% Solids: 10.7

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| CAS No. | Analyte | Concentration | CI | Q | IM |
|------------|-----------|---------------|----|-----|----|
| 17429-90-5 | Aluminum | 275 | | | IP |
| 17440-36-0 | Antimony | 0.14 | UI | | IP |
| 17440-38-2 | Arsenic | 12.4 | | *E | IP |
| 17440-39-3 | Barium | 1.7 | BI | | IP |
| 17440-41-7 | Beryllium | 0.05 | UI | | IP |
| 17440-43-9 | Cadmium | 0.47 | | | IP |
| 17440-70-2 | Calcium | 12400 | | *E | IP |
| 17440-47-3 | Chromium | 1.2 | | | IP |
| 17440-48-4 | Cobalt | 2.3 | BI | | IP |
| 17440-50-8 | Copper | 12.6 | | | IP |
| 17439-89-6 | Iron | 442 | | E | IP |
| 17439-92-1 | Lead | 1.8 | | N | IP |
| 17439-95-4 | Magnesium | 8370 | | E | IP |
| 17439-96-5 | Manganese | 33.5 | | N*E | IP |
| 17440-02-0 | Nickel | 4.1 | | | IP |
| 17440-09-7 | Potassium | 11600 | | E | IP |
| 17782-49-2 | Selenium | 2.8 | | N | IP |
| 17440-22-4 | Silver | 0.04 | UI | N | IP |
| 17440-23-5 | Sodium | 71900 | | | IP |
| 17440-28-0 | Thallium | 0.10 | UI | | IP |
| 17440-62-2 | Vanadium | 1.9 | BI | | IP |
| 17440-66-6 | Zinc | 117 | | NE | IP |

Color Before: GREY

Clarity Before:

Texture: MEDIUM

Color After: YELLOW

Clarity After:

Artifacts:

Comments:

TOTAL METALS TISSUE

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

RF-MU02-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC

Case No.: CTO 100 SAS No.:

SDG No.: TY0199A

Matrix (soil/water): SOIL

Lab Sample ID: T990034-09 S

Level (low/med): LOW

Date Received: 01/19/99

% Solids: 17.9

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| CAS No. | Analyte | Concentration | Q | IM |
|------------|-----------|---------------|-----|----|
| 17429-90-5 | Aluminum | 319 | | IP |
| 17440-36-0 | Antimony | 0.091U | | IP |
| 17440-38-2 | Arsenic | 15.2 | *E | IP |
| 17440-39-3 | Barium | 1.2 | | IP |
| 17440-41-7 | Beryllium | 0.031U | | IP |
| 17440-43-9 | Cadmium | 0.21 | | IP |
| 17440-70-2 | Calcium | 19800 | *E | IP |
| 17440-47-3 | Chromium | 1.1 | | IP |
| 17440-48-4 | Cobalt | 0.34 | | IP |
| 17440-50-8 | Copper | 4.1 | | IP |
| 17439-89-6 | Iron | 473 | E | IP |
| 17439-92-1 | Lead | 1.0 | N | IP |
| 17439-95-4 | Magnesium | 3410 | E | IP |
| 17439-96-5 | Manganese | 22.8 | N*E | IP |
| 17440-02-0 | Nickel | 1.1 | | IP |
| 17440-09-7 | Potassium | 6390 | E | IP |
| 17782-49-2 | Selenium | 3.5 | N | IP |
| 17440-22-4 | Silver | 0.031U | N | IP |
| 17440-23-5 | Sodium | 27200 | | IP |
| 17440-28-0 | Thallium | 0.071U | | IP |
| 17440-62-2 | Vanadium | 3.1 | | IP |
| 17440-66-6 | Zinc | 38.3 | NE | IP |

Color Before: GREY

Clarity Before:

Texture: MEDIUM

Color After: YELLOW

Clarity After:

Artifacts:

Comments:

TOTAL METALS TISSUE

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

SB-MU01/02-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC Case No.: CTO 100 SAS No.: SDG No.: TY0199A

Matrix (soil/water): SOIL Lab Sample ID: T990034-02 S

Level (low/med): LOW Date Received: 01/19/99

% Solids: 15.4

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| CAS No. | Analyte | Concentration | Q | IM |
|------------|-----------|---------------|-----|----|
| 17429-90-5 | Aluminum | 1010 | | IP |
| 17440-36-0 | Antimony | 0.091 | | IP |
| 17440-38-2 | Arsenic | 11.0 | *E | IP |
| 17440-39-3 | Barium | 1.5 | | IP |
| 17440-41-7 | Beryllium | 0.051 | | IP |
| 17440-43-9 | Cadmium | 0.241 | | IP |
| 17440-70-2 | Calcium | 19500 | *E | IP |
| 17440-47-3 | Chromium | 1.9 | | IP |
| 17440-48-4 | Cobalt | 0.551 | | IP |
| 17440-50-8 | Copper | 12.5 | | IP |
| 17439-89-6 | Iron | 1100 | E | IP |
| 17439-92-1 | Lead | 4.8 | N | IP |
| 17439-95-4 | Magnesium | 3000 | E | IP |
| 17439-96-5 | Manganese | 26.3 | N*E | IP |
| 17440-02-0 | Nickel | 3.0 | | IP |
| 17440-09-7 | Potassium | 7130 | E | IP |
| 17782-49-2 | Selenium | 3.4 | N | IP |
| 17440-22-4 | Silver | 0.031 | N | IP |
| 17440-23-5 | Sodium | 24100 | | IP |
| 17440-28-0 | Thallium | 0.071 | | IP |
| 17440-62-2 | Vanadium | 3.3 | | IP |
| 17440-66-6 | Zinc | 45.0 | NE | IP |

Color Before: GREY Clarity Before: Texture: MEDIUM

Color After: YELLOW Clarity After: Artifacts:

Comments:

TOTAL METALS TISSUE

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

Lab Name: CEIMIC CORPORATION

| |
|-------------|
| SB-MU04-99A |
|-------------|

Contract: BAKER ENV.

Lab Code: CEIMIC Case No.: CTO 100 SAS No.: SDG No.: TY0199A

Matrix (soil/water): SOIL Lab Sample ID: T990034-12 S

Level (low/med): LOW Date Received: 01/19/99

* Solids: 6.7

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| CAS No. | Analyte | Concentration | Q | IM |
|------------|-----------|---------------|------|----|
| 17429-90-5 | Aluminum | 455 | | IP |
| 17440-36-0 | Antimony | 0.23 | UI | IP |
| 17440-38-2 | Arsenic | 37.4 | EE | IP |
| 17440-39-3 | Barium | 2.3 | BI | IP |
| 17440-41-7 | Beryllium | 0.09 | UI | IP |
| 17440-43-9 | Cadmium | 0.52 | | IP |
| 17440-70-2 | Calcium | 24100 | EE | IP |
| 17440-47-3 | Chromium | 2.0 | | IP |
| 17440-48-4 | Cobalt | 0.64 | BI | IP |
| 17440-50-8 | Copper | 12.9 | | IP |
| 17439-89-6 | Iron | 643 | EE | IP |
| 17439-92-1 | Lead | 1.6 | NN | IP |
| 17439-95-4 | Magnesium | 7950 | EE | IP |
| 17439-96-5 | Manganese | 91.9 | NN*E | IP |
| 17440-02-0 | Nickel | 3.0 | BI | IP |
| 17440-09-7 | Potassium | 18900 | EE | IP |
| 17782-49-2 | Selenium | 9.4 | NN | IP |
| 17440-22-4 | Silver | 0.07 | UIN | IP |
| 17440-23-5 | Sodium | 69800 | | IP |
| 17440-28-0 | Thallium | 0.18 | UI | IP |
| 17440-62-2 | Vanadium | 4.9 | | IP |
| 17440-66-6 | Zinc | 101 | NE | IP |

Color Before: GREY Clarity Before: Texture: MEDIUM

Color After: YELLOW Clarity After: Artifacts:

Comments:

TOTAL METALS TISSUE

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

SB-MU05-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC Case No.: CTO 100 SAS No.: SDG No.: TY0199A

Matrix (soil/water): SOIL Lab Sample ID: T990034-05 S

Level (low/med): LOW Date Received: 01/19/99

% Solids: 18.6

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| CAS No. | Analyte | Concentration | CI | Q | IM |
|------------|-----------|---------------|----|-----|----|
| 17429-90-5 | Aluminum | 317 | | | IP |
| 17440-36-0 | Antimony | 0.091 | U | | IP |
| 17440-38-2 | Arsenic | 11.3 | | *E | IP |
| 17440-39-3 | Barium | 0.831 | B | | IP |
| 17440-41-7 | Beryllium | 0.031 | U | | IP |
| 17440-43-9 | Cadmium | 0.191 | | | IP |
| 17440-70-2 | Calcium | 8470 | | *E | IP |
| 17440-47-3 | Chromium | 0.821 | | | IP |
| 17440-48-4 | Cobalt | 0.351 | B | | IP |
| 17440-50-8 | Copper | 4.1 | | | IP |
| 17439-89-6 | Iron | 366 | | E | IP |
| 17439-92-1 | Lead | 1.3 | | N | IP |
| 17439-95-4 | Magnesium | 3010 | | E | IP |
| 17439-96-5 | Manganese | 18.1 | | N*E | IP |
| 17440-02-0 | Nickel | 0.731 | B | | IP |
| 17440-09-7 | Potassium | 6080 | | E | IP |
| 17782-49-2 | Selenium | 3.3 | | N | IP |
| 17440-22-4 | Silver | 0.031 | U | N | IP |
| 17440-23-5 | Sodium | 27600 | | | IP |
| 17440-28-0 | Thallium | 0.071 | U | | IP |
| 17440-62-2 | Vanadium | 1.9 | | | IP |
| 17440-66-6 | Zinc | 34.1 | | NE | IP |

Color Before: GREY Clarity Before: Texture: MEDIUM

Color After: YELLOW Clarity After: Artifacts:

Comments:

TOTAL METALS TISSUE

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

Lab Name: CEIMIC CORPORATION

| |
|-------------|
| RF-0Y01-99A |
|-------------|

Contract: BAKER ENV.

Lab Code: CEIMIC Case No.: CTO 100 SAS No.: SDG No.: TY0199A

Matrix (soil/water): SOIL Lab Sample ID: T990034-07 S

Level (low/med): LOW Date Received: 01/19/99

* Solids: 11.8

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| ICAS No. | Analyte | Concentration | CI | Q | IM |
|------------|-----------|---------------|----|-----|----|
| 17429-90-5 | Aluminum | 104 | | | IP |
| 17440-36-0 | Antimony | 0.13 | UI | | IP |
| 17440-38-2 | Arsenic | 18.8 | | *E | IP |
| 17440-39-3 | Barium | 1.2 | BI | | IP |
| 17440-41-7 | Beryllium | 0.05 | UI | | IP |
| 17440-43-9 | Cadmium | 0.20 | BI | | IP |
| 17440-70-2 | Calcium | 4390 | | *E | IP |
| 17440-47-3 | Chromium | 0.85 | | | IP |
| 17440-48-4 | Cobalt | 2.6 | | | IP |
| 17440-50-8 | Copper | 8.2 | | | IP |
| 17439-89-6 | Iron | 243 | | E | IP |
| 17439-92-1 | Lead | 2.0 | | N | IP |
| 17439-95-4 | Magnesium | 5900 | | E | IP |
| 17439-96-5 | Manganese | 10.3 | | N*E | IP |
| 17440-02-0 | Nickel | 4.5 | | | IP |
| 17440-09-7 | Potassium | 7500 | | E | IP |
| 17782-49-2 | Selenium | 2.3 | | N | IP |
| 17440-22-4 | Silver | 0.04 | UI | N | IP |
| 17440-23-5 | Sodium | 46800 | | | IP |
| 17440-28-0 | Thallium | 0.10 | UI | | IP |
| 17440-62-2 | Vanadium | 3.3 | | | IP |
| 17440-66-6 | Zinc | 88.9 | | NE | IP |

Color Before: GREY Clarity Before: Texture: MEDIUM

Color After: YELLOW Clarity After: Artifacts:

Comments:

TOTAL METALS TISSUE

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

RF-0Y02-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC Case No.: CTO 100 SAS No.: SDG No.: TY0199A

Matrix (soil/water): SOIL Lab Sample ID: T990034-13 S

Level (low/med): LOW Date Received: 01/19/99

* Solids: 9.8

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| ICAS No. | Analyte | Concentration | CI | Q | IM |
|------------|-----------|---------------|----|-----|----|
| 17429-90-5 | Aluminum | 320 | | | IP |
| 17440-36-0 | Antimony | 0.15 | U | | IP |
| 17440-38-2 | Arsenic | 32.2 | | *E | IP |
| 17440-39-3 | Barium | 1.2 | B | | IP |
| 17440-41-7 | Beryllium | 0.06 | U | | IP |
| 17440-43-9 | Cadmium | 1.8 | | | IP |
| 17440-70-2 | Calcium | 18200 | | E | IP |
| 17440-47-3 | Chromium | 1.5 | | | IP |
| 17440-48-4 | Cobalt | 0.84 | B | | IP |
| 17440-50-8 | Copper | 88.2 | | | IP |
| 17439-89-6 | Iron | 581 | | E | IP |
| 17439-92-1 | Lead | 1.2 | | N | IP |
| 17439-95-4 | Magnesium | 6150 | | E | IP |
| 17439-96-5 | Manganese | 19.2 | | N*E | IP |
| 17440-02-0 | Nickel | 2.1 | B | | IP |
| 17440-09-7 | Potassium | 15200 | | E | IP |
| 17782-49-2 | Selenium | 5.6 | | N | IP |
| 17440-22-4 | Silver | 1.4 | | N | IP |
| 17440-23-5 | Sodium | 51000 | | | IP |
| 17440-28-0 | Thallium | 0.12 | U | | IP |
| 17440-62-2 | Vanadium | 4.6 | | | IP |
| 17440-66-6 | Zinc | 2230 | | NE | IP |

Color Before: GREY Clarity Before: Texture: MEDIUM

Color After: YELLOW Clarity After: Artifacts:

Comments:

TOTAL METALS TISSUE

1

EPA SAMPLE NO.

INORGANIC ANALYSIS DATA SHEET

| |
|-------------|
| SB-0Y01-99A |
|-------------|

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC Case No.: CTO 100 SAS No.: SDG No.: TY0199A

Matrix (soil/water): SOIL Lab Sample ID: T990034-01 S

Level (low/med): LOW Date Received: 01/19/99

% Solids: 15.1

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| CAS No. | Analyte | Concentration | Q | IM |
|------------|-----------|---------------|-----|----|
| 17429-90-5 | Aluminum | 111 | | IP |
| 17440-36-0 | Antimony | 0.10 | U | IP |
| 17440-38-2 | Arsenic | 12.5 | *E | IP |
| 17440-39-3 | Barium | 0.83 | B | IP |
| 17440-41-7 | Beryllium | 0.04 | U | IP |
| 17440-43-9 | Cadmium | 1.6 | | IP |
| 17440-70-2 | Calcium | 2330 | E | IP |
| 17440-47-3 | Chromium | 0.65 | | IP |
| 17440-48-4 | Cobalt | 0.37 | B | IP |
| 17440-50-8 | Copper | 50.3 | | IP |
| 17439-89-6 | Iron | 225 | E | IP |
| 17439-92-1 | Lead | 1.6 | N | IP |
| 17439-95-4 | Magnesium | 2740 | E | IP |
| 17439-96-5 | Manganese | 9.8 | N*E | IP |
| 17440-02-0 | Nickel | 2.4 | | IP |
| 17440-09-7 | Potassium | 8940 | E | IP |
| 17782-49-2 | Selenium | 3.2 | N | IP |
| 17440-22-4 | Silver | 0.34 | B N | IP |
| 17440-23-5 | Sodium | 19700 | | IP |
| 17440-28-0 | Thallium | 0.08 | U | IP |
| 17440-62-2 | Vanadium | 1.4 | B | IP |
| 17440-66-6 | Zinc | 1280 | NE | IP |

Color Before: GREY Clarity Before: Texture: MEDIUM

Color After: YELLOW Clarity After: Artifacts:

Comments:

TOTAL METALS TISSUE

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

| |
|-------------|
| SB-0Y02-99A |
|-------------|

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC Case No.: CTO 100 SAS No.: SDG No.: TY0199A

Matrix (soil/water): SOIL Lab Sample ID: T990034-06 S

Level (low/med): LOW Date Received: 01/19/99

% Solids: 15.7

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| CAS No. | Analyte | Concentration | Q | IM |
|------------|-----------|---------------|-----|----|
| 17429-90-5 | Aluminum | 49.3 | | IP |
| 17440-36-0 | Antimony | 0.10 | U | IP |
| 17440-38-2 | Arsenic | 7.4 | E | IP |
| 17440-39-3 | Barium | 0.35 | R | IP |
| 17440-41-7 | Beryllium | 0.04 | U | IP |
| 17440-43-9 | Cadmium | 1.1 | | IP |
| 17440-70-2 | Calcium | 2830 | E | IP |
| 17440-47-3 | Chromium | 0.46 | | IP |
| 17440-48-4 | Cobalt | 0.34 | R | IP |
| 17440-50-8 | Copper | 45.1 | | IP |
| 17439-89-6 | Iron | 123 | E | IP |
| 17439-92-1 | Lead | 0.69 | N | IP |
| 17439-95-4 | Magnesium | 2450 | E | IP |
| 17439-96-5 | Manganese | 6.4 | N* | IP |
| 17440-02-0 | Nickel | 1.1 | R | IP |
| 17440-09-7 | Potassium | 8780 | E | IP |
| 17782-49-2 | Selenium | 2.8 | N | IP |
| 17440-22-4 | Silver | 0.03 | U N | IP |
| 17440-23-5 | Sodium | 18600 | | IP |
| 17440-28-0 | Thallium | 0.08 | U | IP |
| 17440-62-2 | Vanadium | 0.68 | R | IP |
| 17440-66-6 | Zinc | 793 | NE | IP |

Color Before: GREY Clarity Before: Texture: MEDIUM

Color After: YELLOW Clarity After: Artifacts:

Comments:

TOTAL METALS TISSUE

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

SR-OY03-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC Case No.: CTO 100 SAS No.: SDG No.: TY0199A

Matrix (soil/water): SOIL Lab Sample ID: T990034-03 S

Level (low/med): LOW Date Received: 01/19/99

% Solids: 12.0

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| ICAS No. | Analyte | Concentration | Q | M |
|------------|-----------|---------------|----|----|
| 17429-90-5 | Aluminum | 90.6 | | IP |
| 17440-36-0 | Antimony | 0.12 | U | IP |
| 17440-38-2 | Arsenic | 10.0 | E | IP |
| 17440-39-3 | Barium | 0.62 | B | IP |
| 17440-41-7 | Beryllium | 0.05 | U | IP |
| 17440-43-9 | Cadmium | 1.7 | | IP |
| 17440-70-2 | Calcium | 4040 | E | IP |
| 17440-47-3 | Chromium | 0.66 | | IP |
| 17440-48-4 | Cobalt | 0.25 | U | IP |
| 17440-50-8 | Copper | 46.0 | | IP |
| 17439-89-6 | Iron | 199 | E | IP |
| 17439-92-1 | Lead | 1.0 | N | IP |
| 17439-95-4 | Magnesium | 3960 | E | IP |
| 17439-96-5 | Manganese | 11.8 | N | IP |
| 17440-02-0 | Nickel | 2.3 | | IP |
| 17440-09-7 | Potassium | 9090 | E | IP |
| 17782-49-2 | Selenium | 3.1 | N | IP |
| 17440-22-4 | Silver | 0.04 | U | IP |
| 17440-23-5 | Sodium | 31700 | | IP |
| 17440-28-0 | Thallium | 0.09 | U | IP |
| 17440-62-2 | Vanadium | 0.56 | B | IP |
| 17440-66-6 | Zinc | 722 | NE | IP |

Color Before: GREY Clarity Before: Texture: MEDIUM

Color After: YELLOW Clarity After: Artifacts:

Comments:

TOTAL METALS TISSUE

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

SB-OY04-99A

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC Case No.: CTD 100 SAS No.: SDG No.: TY0199A

Matrix (soil/water): SOIL Lab Sample ID: T990034-10 S

Level (low/med): LOW Date Received: 01/19/99

% Solids: 11.3

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| CAS No. | Analyte | Concentration | Q | IM |
|------------|-----------|---------------|-----|----|
| 17429-90-5 | Aluminum | 69.5 | | IP |
| 17440-36-0 | Antimony | 0.13 | U | IP |
| 17440-38-2 | Arsenic | 11.9 | *E | IP |
| 17440-39-3 | Barium | 0.68 | B | IP |
| 17440-41-7 | Beryllium | 0.05 | U | IP |
| 17440-43-9 | Cadmium | 1.3 | | IP |
| 17440-70-2 | Calcium | 7840 | *E | IP |
| 17440-47-3 | Chromium | 0.72 | | IP |
| 17440-48-4 | Cobalt | 0.27 | U | IP |
| 17440-50-8 | Copper | 21.4 | | IP |
| 17439-89-6 | Iron | 228 | E | IP |
| 17439-92-1 | Lead | 0.82 | N | IP |
| 17439-95-4 | Magnesium | 5110 | E | IP |
| 17439-96-5 | Manganese | 7.0 | N*E | IP |
| 17440-02-0 | Nickel | 1.8 | B | IP |
| 17440-09-7 | Potassium | 11200 | E | IP |
| 17782-49-2 | Selenium | 2.8 | N | IP |
| 17440-22-4 | Silver | 0.04 | U N | IP |
| 17440-23-5 | Sodium | 43800 | | IP |
| 17440-28-0 | Thallium | 0.10 | U | IP |
| 17440-62-2 | Vanadium | 1.7 | B | IP |
| 17440-66-6 | Zinc | 728 | NE | IP |

Color Before: GREY Clarity Before: Texture: MEDIUM

Color After: YELLOW Clarity After: Artifacts:

Comments:

TOTAL METALS TISSUE

1

INORGANIC ANALYSIS DATA SHEET

EPA SAMPLE NO.

| |
|-------------|
| SB-0Y05-99A |
|-------------|

Lab Name: CEIMIC CORPORATION

Contract: BAKER ENV.

Lab Code: CEIMIC Case No.: CTO 100 SAS No.: SDG No.: TY0199A

Matrix (soil/water): SOIL Lab Sample ID: T990034-04 S

Level (low/med): LOW Date Received: 01/19/99

% Solids: 20.0

Concentration Units (ug/L or mg/Kg dry weight): MG/KG

| ICAS No. | Analyte | Concentration | CI | Q | IM |
|------------|-----------|---------------|----|-----|----|
| 17429-90-5 | Aluminum | 78.9 | | | IP |
| 17440-36-0 | Antimony | 0.08 | U | | IP |
| 17440-38-2 | Arsenic | 4.1 | | *E | IP |
| 17440-39-3 | Barium | 0.30 | B | | IP |
| 17440-41-7 | Beryllium | 0.03 | U | | IP |
| 17440-43-9 | Cadmium | 0.60 | | | IP |
| 17440-70-2 | Calcium | 5960 | | E | IP |
| 17440-47-3 | Chromium | 0.43 | | | IP |
| 17440-48-4 | Cobalt | 0.21 | B | | IP |
| 17440-50-8 | Copper | 17.1 | | | IP |
| 17439-89-6 | Iron | 137 | | E | IP |
| 17439-92-1 | Lead | 0.40 | | N | IP |
| 17439-95-4 | Magnesium | 2620 | | E | IP |
| 17439-96-5 | Manganese | 5.7 | | N*E | IP |
| 17440-02-0 | Nickel | 1.3 | | | IP |
| 17440-09-7 | Potassium | 5000 | | E | IP |
| 17782-49-2 | Selenium | 1.0 | | N | IP |
| 17440-22-4 | Silver | 0.02 | U | N | IP |
| 17440-23-5 | Sodium | 22400 | | | IP |
| 17440-28-0 | Thallium | 0.06 | U | | IP |
| 17440-62-2 | Vanadium | 0.35 | B | | IP |
| 17440-66-6 | Zinc | 894 | | NE | IP |

Color Before: GREY Clarity Before: Texture: MEDIUM

Color After: YELLOW Clarity After: Artifacts:

Comments:

STONE BAY

**CLAM, MUSSEL, AND
OYSTER TISSUE - MERCURY**

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: PBO

Date Sampled:

Laboratory ID: PBO

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm) +

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | ND | 0.003 |

ND = Not Detected

Reported by: _____

Donald Totocelli

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

**LABORATORY CONTROL SAMPLE SUMMARY
TOTAL METALS
CLP METHOD ILM04.0**

Client: Baker Environmental

Ceimic Project: 990034

Laboratory Control Spike ID: SRM-2

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm)

| Target Analyte | Preparation Batch | True Value | Lab Control Spike Result | Lab Control Spike Recovery(%) | QC Limits(%) |
|----------------|-------------------|------------|--------------------------|-------------------------------|--------------|
| Mercury | 0201 | 0.064 | 0.0570 | 89.1 | 80 - 120 |

Reported by: _____

Donald Totorelli

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: RF-CL01-99A

Date Sampled: 01/16/99

Laboratory ID: 990034-08

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm) +

Percent Solids: 15

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.050 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Totolli

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

DUPLICATE SAMPLE SUMMARY
TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: RF-CL01-99A

Date Sampled: 01/16/99

Laboratory ID: 990034-08Dup

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Tissue

Concentration in: mg/Kg (ppm)+

Duplicate Percent Solids: 15

| Target Analyte | Sample Result | Duplicate Result | RPD(%) | QC Limit(%) |
|----------------|---------------|------------------|--------|-------------|
| Mercury | 0.0500 | 0.0620 | 21 | 20 |

+ Dry weight basis.

RPD = Relative Percent Difference

Reported by: _____

Donald Tortorelli

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

**SPIKE SAMPLE SUMMARY
TOTAL METALS
CLP METHOD ILM04.0**

Client: Baker Environmental

Client Sample ID: RF-CL01-99A

Date Sampled: 01/16/99

Laboratory ID: 990034-08Spk

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Tissue

Concentration in: mg/Kg (ppm)+

Percent Solids: 15

| Target Analyte | Sample Result | Predigest Spike Added | Spiked Sample Result | Recovery(%) | | |
|----------------|---------------|-----------------------|----------------------|-----------------|-----------|-------------------|
| | | | | Predigest Spike | QC Limits | Post Digest Spike |
| Mercury | 0.0500 | 0.150 | 0.167 | 78 | 75 - 125 | NR |

NR = Not Required
+ Dry weight basis.

Reported by: Donald Zatoalli

Approved by: MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: RF-CL02-99A

Date Sampled: 01/16/99

Laboratory ID: 990034-21

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm) +

Percent Solids: 12

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.054 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Tortorelli

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: SB-CL04-99A

Date Sampled: 01/15/99

Laboratory ID: 990034-11

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm) +

Percent Solids: 11

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.048 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Tortorelli

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: SB-MU01/02-99A

Date Sampled: 01/16/99

Laboratory ID: 990034-02

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm) +

Percent Solids: 15

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.088 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Tatorelli

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

**TOTAL METALS
CLP METHOD ILM04.0**

Client: Baker Environmental

Client Sample ID: RF-MU02-99A

Date Sampled: 01/16/99

Laboratory ID: 990034-09

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm)+

Percent Solids: 18

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.082 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Fatoelli

Approved by: _____

ZIP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: SB-MU04-99A

Date Sampled: 01/16/99

Laboratory ID: 990034-12

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm)+

Percent Solids: 7

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.157 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Tortelli

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: SB-MU05-99A

Date Sampled: 01/15/99

Laboratory ID: 990034-05

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm) +

Percent Solids: 19

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.086 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Tortorelli

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: RF-OY01-99A

Date Sampled: 01/16/99

Laboratory ID: 990034-07

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm) +

Percent Solids: 12

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.094 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Fortsch

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: SB-OY01-99A

Date Sampled: 01/14/99

Laboratory ID: 990034-01

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm)+

Percent Solids: 15

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.063 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Fattouhi

Approved by: _____

NP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: SB-OY02-99A

Date Sampled: 01/15/99

Laboratory ID: 990034-06

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm) +

Percent Solids: 16

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.069 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Fortelli

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: RF-OY02-99A

Date Sampled: 01/16/99

Laboratory ID: 990034-13

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm) +

Percent Solids: 10

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.156 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Totelli

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: SB-OY03-99A

Date Sampled: 01/15/99

Laboratory ID: 990034-03

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm)+

Percent Solids: 12

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.252 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Tortorelli

Approved by: _____

JP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

TOTAL METALS
CLP METHOD ILM04.0

Client: Baker Environmental

Client Sample ID: SB-OY04-99A

Date Sampled: 01/15/99

Laboratory ID: 990034-10

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm)+

Percent Solids: 11

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.070 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Tortorelli

Approved by: _____

MP

**CEIMIC
Corporation**
"Analytical Chemistry for Environmental Management"

**TOTAL METALS
CLP METHOD ILM04.0**

Client: Baker Environmental

Client Sample ID: SB-OY05-99A

Date Sampled: 01/15/99

Laboratory ID: 990034-04

Date Sample Received: 01/19/99

Date Analysis Completed: 02/01/99

Matrix: Soil

Concentration in: mg/Kg (ppm)+

Percent Solids: 20

| Target Analyte | Preparation Batch | Sample Concentration | Reporting Limit |
|----------------|-------------------|----------------------|-----------------|
| Mercury | 0201 | 0.053 | 0.003 |

+ Dry weight basis.

Reported by: _____

Donald Tatorelli

Approved by: _____

MP

Baker

Baker Environmental, Inc.

ATTACHMENT D

Analytical Quality Assurance

ENVIRONMENTAL
RESOURCE ASSOCIATES
Arvada, Colorado 80002 1-800-ERA-0122

930621C → 930621C



Certification

F21 → F24

PriorityPollutnt™/CLP Quality Control Standards

Inorganics in Soil

Lot 216

| Parameter | Lot Number 216 Certified Value mg/kg | Advisory Range mg/kg | |
|---------------------|--|-------------------------|-------------|
| TRACE METALS | | | |
| aluminum | 6000 | 3800 - 8400 | 60 - 140% |
| antimony | 27.8 | 14 - 117 | 50 - 150 |
| arsenic | 67.7 | 41 - 105 | 61 - 150 |
| barium | 187 | 131 - 243 | 70 - 130 |
| beryllium | 57.5 | 35 - 81 | 61 - 141 |
| cadmium | 110 | 55 - 186 | 50 - 150 |
| calcium | 2040 | 1220 - 2880 | 60 - 140 |
| chromium | 189 | 95 - 265 | 50 - 140 |
| cobalt | 87.0 | 43 - 130 | 30 49 - 149 |
| copper | 141 | 84 - 200 | 60 - 142 |
| iron | 10800 | 7020 - 15100 | 65 - 140 |
| lead | 100 | 55 - 140 | 55 - 140 |
| magnesium | 2050 | 1200 - 3080 | 59 - 150 |
| manganese | 294 | 206 - 383 | 70 - 130 |
| mercury | 2.36 | 1.3 - 3.8 | 55 - 161 |
| molybdenum | 124 | 93 - 167 | 75 - 135 |
| nickel | 79.6 | 40 - 112 | 50 - 140 |
| potassium | 2130 | 1280 - 2770 | 60 - 130 |
| selenium | 99.1 | 54 - 149 | 54 - 150 |
| silver | 124 | 62 - 186 | 50 - 150 |
| sodium | 527 | 316 - 738 | 60 - 140 |
| thallium | 67.9 | 34 - 102 | 50 - 150 |
| vanadium | 84.8 | 59 - 115 | 70 - 135 |
| zinc | 197 | 98 - 280 | 50 - 142 |

1. The Trace Metals Certified Values are equal to the mean recoveries for each parameter as determined in an interlaboratory round robin study (3 laboratories, 10 to 24 data points per parameter). The standard was digested using Method 3050, SW-846 and the digest analyzed by ICP and atomic absorption spectroscopy.

2. The Advisory Ranges are listed as guidelines for acceptable recoveries given the limitations of the EPA methodologies commonly used to determine these parameters. The range closely approximates the 95% confidence interval for these parameters based upon experimental data from this lot, previous ERA lots and published USEPA data.



FBO
94632B1

National Institute of Standards & Technology

Certificate of Analysis

Standard Reference Material 1566a

Oyster Tissue

This Standard Reference Material (SRM) is intended primarily for use in calibrating instrumentation and validating methodology for the chemical analysis of marine bivalve tissue. A unit of SRM 1566a contains approximately 25 grams of oyster tissue.

Certified Concentrations of Constituent Elements: The certified elemental concentrations are shown in Table 1. Certified values are based on results obtained by reference methods of known accuracy; or alternatively, from results obtained by two or more independent and reliable analytical methods. Noncertified values are given, for information only, in Table 2. All values are based on minimum sample size of 250 mg of the dried material.

NOTICE AND WARNINGS TO USERS

Expiration of Certification: This certification is invalid after 5 years from the date of shipping. Please return the attached registration card to register your SRM.

Storage: The material should be kept in its tightly closed, original bottle and stored in a desiccator over $Mg(ClO_4)_2$ at temperatures between 10-30 °C. It should not be exposed to intense sources of radiation, including ultraviolet lamps or sunlight.

Use: A minimum sample weight of 250 mg of the dried material (see Instructions for Drying) is necessary for any certified value in Table 1 to be valid within the stated uncertainty. The bottle should be shaken well before each use, closed tightly immediately after use, and stored as described above.

The statistical analysis of the data was performed by S.B. Schiller and K.R. Eberhardt of the Statistical Engineering Division.

The overall direction and coordination of the analytical chemistry measurements leading to this certificate were performed in the NIST Center for Analytical Chemistry by R. Zeisler.

The technical and support aspects involved in the preparation, certification, and issuance of this Standard Reference Material were coordinated through the Office of Standard Reference Materials by R. Alvarez.

Gaithersburg, MD 20899
October 2, 1989

Stanley D. Rasberry, Chief
Office of Standard Reference Materials

(over)

Instructions for Drying: Before weighing, samples of SRM 1566a should be dried to constant weight by one of the following procedures:

1. Reduced-pressure drying at room temperature for 48 hours over $Mg(ClO_4)_2$ in a vacuum desiccator at approximately 1.3×10^4 Pa (100 mm Hg).
2. Vacuum drying at room temperature for 24 hours at a pressure of approximately 30 Pa (0.2 mm Hg) using a cold trap.
3. Freeze drying for 20 hours at a pressure of approximately 3 Pa (0.02 mm Hg).

Source and Preparation of Material: The oysters for this reference material were obtained from a commercial source. They had been shucked, frozen, and packaged in sealed plastic bags. The oyster material was ground, freeze-dried, and powdered at Leon Laboratories, Fort Lauderdale, FL. At NIST, the oyster tissue was jet-milled to pass a 355- μ m screen, radiation-sterilized, and bottled.

Homogeneity Assessment: Samples from randomly selected bottles of SRM 1566a were analyzed for homogeneity by x-ray fluorescence and neutron activation methods. In addition, results by other analytical methods were examined for evidence of inhomogeneity. The uncertainties in Table 1 include estimates of inhomogeneity.

Table 1. Certified Concentrations of Constituent Elements

| Element ¹ | Concentration, ² Wt. Percent | Element ¹ | Concentration, ² Wt. Percent |
|---------------------------------|--|------------------------------|--|
| Calcium ^{d,e,o} | 0.196 ± 0.019 | Potassium ^{f,k} | 0.790 ± 0.047 |
| Chlorine ^{g,k,o} | 0.829 ± 0.014 | Sodium ^{d,k} | 0.417 ± 0.013 |
| Magnesium ^{d,e,k} | 0.118 ± 0.017 | Sulfur ^{d,l} | 0.862 ± 0.019 |
| Phosphorus ^{d,n} | 0.623 ± 0.018 | | |
| Element ¹ | Concentration, ² μ g/g | Element ¹ | Concentration, ² μ g/g |
| Aluminum ^{f,k} | 202.5 ± 12.5 | Manganese ^{d,e,k,o} | 12.3 ± 1.5 |
| Arsenic ^{c,k,l} | 14.0 ± 1.2 | Mercury ^{d,l} | 0.0642 ± .0067 |
| Cadmium ^{b,h,k,l,m} | 4.15 ± 0.38 | Nickel ^{b,h,l,o} | 2.25 ± 0.44 |
| Chromium ^{b,k,l} | 1.43 ± 0.46 | Selenium ^{k,l,o} | 2.21 ± 0.24 |
| Cobalt ^{k,l} | 0.57 ± 0.11 | Silver ^{b,k,l} | 1.68 ± 0.15 |
| Copper ^{b,e,h,k,l,m,o} | 66.3 ± 4.3 | Strontium ^{e,h,o} | 11.1 ± 1.0 |
| Iodine ^{i,k,l} | 4.46 ± 0.42 | Uranium ^{d,l} | 0.132 ± 0.012 |
| Iron ^{d,k,o} | 539 ± 15 | Vanadium ^{d,k,l} | 4.68 ± 0.15 |
| Lead ^h | 0.371 ± 0.014 | Zinc ^{d,e,k,l,m,o} | 830 ± 57 |

^o Lead was determined by isotope dilution mass spectrometry, inductively coupled plasma, at NIST and at another laboratory.

1. Analytical Methods:

- ^aAtomic absorption spectrometry, cold vapor
- ^bAtomic absorption spectrometry, electrothermal
- ^cAtomic absorption spectrometry, hydride generation
- ^dAtomic emission spectrometry, direct current plasma
- ^eAtomic emission spectrometry, inductively coupled plasma
- ^fAtomic emission spectroscopy, flame
- ^gIon chromatography
- ^hIsotope dilution mass spectrometry, inductively coupled plasma
- ⁱIsotope dilution mass spectrometry, resonance ionization
- ^jIsotope dilution mass spectrometry, thermal ionization
- ^kNeutron activation, instrumental
- ^lNeutron activation, radiochemical
- ^mPolarography
- ⁿSpectrophotometry
- ^oX-ray fluorescence spectrometry

2. Based on dry weight. (For drying instructions, please refer to the section of this certificate on Instructions for Drying.)

The certified concentrations are weighted means of results from two or more analytical techniques. The weights for the weighted means were computed according to the iterative procedure of Paule and Mandel [1]. Each uncertainty is obtained from a 95% prediction interval plus an allowance for systematic error among the methods used. The allowance for systematic error is equal to the greatest difference between the weighted mean (certified value) and the component means for the analytical methods used. In the absence of systematic error, the resulting uncertainty limits will cover the concentration of approximately 95% of samples of this SRM having a minimum sample size of 250 mg.

Table 2. Noncertified Concentrations of Constituent Elements

| Element | Concentration, Percent by Weight |
|---------------------|-------------------------------------|
| Nitrogen (Kjeldahl) | (6.81) |

Method Reference. Official Methods of Analysis of the Association of Official Analytical Chemists, Arlington, VA, 14th Ed., 1984, p. 16, Nitrogen (Total) in Fertilizers, Kjeldahl Method (Final Action): Method 2.057, Improved Method for Nitrate Free Samples. Mercuric Oxide was used as a catalyst. Samples were dried as described in procedure 2 under "Instructions for Drying".

| Element | Concentration, µg/g | Element | Concentration, µg/g |
|-----------|------------------------|----------|------------------------|
| Antimony | (0.01) | Rubidium | (3) |
| Cerium | (0.4) | Samarium | (0.06) |
| Cesium | (0.02) | Scandium | (0.06) |
| Europium | (0.01) | Tantalum | (0.003) |
| Fluorine | (240) | Terbium | (0.007) |
| Gold | (0.01) | Thorium | (0.04) |
| Hafnium | (0.04) | *Tin | (3) |
| Lanthanum | (0.3) | | |

Note: There is evidence that tin is inhomogeneously distributed in the SRM.

Analysts:

Inorganic Analytical Research Division, National Institute of Standards & Technology.

- | | |
|-------------------------|-----------------------|
| 1. E.S. Beary | 17. P.J. Paulsen |
| 2. R.W. Burke | 18. P.A. Pella* |
| 3. T.A. Butler | 19. K.W. Pratt |
| 4. Z. Chun** | 20. T.C. Rains |
| 5. M.S. Epstein | 21. T.A. Rush |
| 6. J.D. Fassett | 22. D.S. Simons* |
| 7. K. Gilliland-Garrity | 23. G.A. Sleater* |
| 8. R. R. Greenberg | 24. M.V. Smith |
| 9. K.E. Hehn | 25. S.F. Stone |
| 10. L.A. Holland | 26. T.M. Sullivan |
| 11. W.F. Koch | 27. R.L. Watters, Jr. |
| 12. W.R. Kelly | 28. L.J. Wood |
| 13. H.M. Kingston | 29. W.Z. Yao** |
| 14. A. Marlow* | 30. X. Yingkai** |
| 15. J.R. Moody | 31. R. Zeisler |
| 16. T.J. Murphy | |

- * Gas and Particulate Science Division
- ** Guest Scientists

Cooperating Analysts & Organizations

Analytical Chemistry Division, National Research Council of Canada, Ottawa, Canada K1A 0R6 -- S. Berman (Coordinator), V. Boyko, J. McLaren, M. Miedema, S. Willie

Battelle, Pacific Northwest Laboratories, Richland, WA 99352 -- R.W. Sanders

Institute "Josef Stefan," Ljubljana, 61111 Yugoslavia, -- A.R. Byrne, M. Dermelj, A. Vakselj and J. Smrke

Reference

- [1] R.C. Paule and J. Mandel, Consensus Values and Weighting Factors, NBS J. Research 87, 377-385 (1982).

Baker

Baker Environmental, Inc.
ATTACHMENT E
Ecological Information

E.1
Life History Information

LIFE HISTORY OF THE GREAT BLUE HERON (*Ardea herodias*)

The great blue heron is the largest and most widely distributed American heron, inhabiting lakes, ponds, rivers, and marshes but is occasionally found in newly-plowed fields and meadows (Bull and Farrand 1977; Eckert 1987). With the exception of the breeding season, this species is mostly solitary in its habits. This heron feeds either by standing motionless in the water waiting for prey, or searching in a stealthy manner with a very slow and careful walk. The primary food item is fish, although frogs, small turtles, crustaceans, mice, voles, shrews, snakes, and ground-nesting birds are also taken. Almost without exception, the great blue heron will shake its bill in the water immediately after swallowing prey, perhaps to wash debris off. Although the digestive fluids of the heron are acidic enough to dissolve bone rapidly, occasionally an undigested pellet of feathers and fur is regurgitated (Eckert 1987).

Courtship occurs soon after the spring migration, with copulation usually occurring on the ground. Colonial nests are placed on the uppermost branches of trees or shrubs. Occasionally a ground nest will be built if it can be placed in a secluded area. Successful nesting areas are usually returned to year after year. Three to four eggs are incubated equally by both sexes for about twenty-eight days (Eckert 1987).

The great blue heron is migratory in the northernmost portion of its range. Lingered birds usually fall prey to severe weather (Bull and Farrand 1977). Southward migration in autumn begins in early October; northward migration in spring begins in March or early April (Eckert 1987).

EXPOSURE PROFILE OF THE GREAT BLUE HERON (*Ardea herodias*)

Adult great blue heron (*Ardea herodias*) range in weight from 2,204 to 2,576 g (U.S. EPA 1993). A food ingestion rate of 0.18 g/g BW/day and a water ingestion rate of 0.045 g/g BW/day are reported for this species (U.S. EPA 1993). Based on these values a 2,204 g heron will consume 396.7 g food/day and 99.2 g water/day. Fall feeding territory size is reported as 1.5 acres, with summer foraging distances from nesting colonies ranging from 2 to 5 miles (U.S. EPA 1993).

An incidental sediment ingestion rate could not be identified for the great blue heron. Therefore, in order to evaluate this exposure pathway, a model was developed which predicted the amount of sediment which may be entrained in the digestive system of a fish, the bluegill (*Lepomis machrochirus*) and crayfish (*Orconectes* sp.). Fish and crayfish are assumed to be the only food source for the great blue heron in order to complete this derivation.

Bluegills commonly reach a size of 12 ounces (Pflieger 1975). From this, the amount of sediment entrained in fish 12 ounces (340 g) in weight was predicted. A study evaluating the stomach contents of 153 bluegills reported an average content of detritus and sediment to be 9.6 percent of the total diet (Kolehmainen 1974). A daily food ingestion rate of 1.75 percent of the body weight per day has been reported for the bluegill (Kolehmainen 1974). This provides a predicted intake rate of 5.95 g of food per day for a 340 g fish. If a conservative assumption is made that 9.6 percent of the food ingested is entirely sediment, it can be predicted that a fish of this size may contain 0.5712 g of sediment in its digestive system.

For the purpose of this model, it was assumed that the level of sediment contained in the digestive system of a fish remains constant over time. This value (0.5712 g) was divided by the predicted fish body weight (340 g) in order to express sediment entrained in fish digestive systems in units of grams of sediment per gram of fish body weight. This provided a value of 0.00168 g sediment/g body weight. When this value is multiplied by the food ingestion rate of the great blue heron (396.7 g/day), the predicted sediment ingestion rate for the heron via consumption of fish is approximately 0.7 g/day.

As with the bluegill, life history information for the crayfish *Orconectes* sp. was used in predicting the incidental sediment ingestion rate for the great blue heron via consumption of freshwater invertebrates. Adult *O. virilis* weigh from 5 to over 20 g and consume 0.3 to 1 percent of its total body weight per day (Kim 1994; Tack 1941; Vannote 1963). In order to express the food ingestion rate in units of g/day, the highest reported food ingestion rate of 1 percent of the total body weight per day was multiplied by the lowest reported body weight of 5 g to

yield a food ingestion rate of 0.05 g/day. *Orconectes* spp. detritus ingestion rates range from 10 percent of the total diet per day in young-of-the-year *Orconectes immunis* (Vannote 1963) to 11 percent of the total diet per day in *O. virilis* (Tack 1941). For this risk assessment, it will be assumed that these values represent the percentage of sediment in the diet of a crayfish. The food ingestion rate of 0.05 g/day was multiplied by the incidental sediment ingestion rate of 11 percent of the total diet per day to yield an incidental sediment ingestion rate of 0.0055 g/day. For the purpose of this model, it was assumed that the level of sediment contained in the digestive system of crayfish remains constant over time. Therefore, in order to express the amount of sediment entrained in a crayfish's digestive system in units of gram of sediment per gram of crayfish body weight, the sediment ingestion rate of 0.0055 g/day was divided by the lowest adult crayfish body weight of 5 g to yield a sediment ingestion rate of 0.0011 g sediment/g BW of crayfish/day. When this value is multiplied by the food ingestion rate of the great blue heron (396.7 g/day), the predicted incidental sediment ingestion rate for the great blue heron via consumption of crayfish is 0.44 g/day.

[The user of this file should then decide what the percent composition of fish versus crayfish the diet is assumed to be for the particular risk assessment, then apply the percentages to these calculated sediment ingestion rates]

REFERENCES

Bull, J. And J. Farrand, Jr. 1977. *The Audubon Society Field Guide to North American Birds, Eastern Region*. Alfred A. Knopf, New York.

Kim, P. 1994. Ecological Risk Assessment, Cannelton Industries Site. Roy F. Weston, Inc. Final Report to the U.S. EPA/ERT. Contract No. 68-C4-0022.

Pflieger, W.L. 1975. *The Fishes of Missouri*. Missouri Dept. Conserv., MO.

Tack, P.I. 1941. "The Life History and Ecology of the Crayfish, *Cambarus immunis* Hagen." *Am. Midl. Nat.* 25:420-466.

U.S. EPA. 1993. *Wildlife Exposure Factors Handbook, Volume I of II*. United States Environmental Protection Agency, Office of Research and Development, Washington, D.C. EPA/600/R-93/187a.

Vannote, R.L. 1963. "Community Productivity and Energy Flow in an Enriched Warm Water Stream." Ph.D. Thesis, Michigan State Univer., E. Lansing. 156pp. In: Momot, W.T., H. Gowing, and P.D. Jones. 1978. "The Dynamics of Crayfish and Their Role in Ecosystems." *Am. Midl. Nat.* 99:10-35.

LIFE HISTORY INFORMATION FOR GREAT BLUE HERON
(Ardea herodias)

| | |
|-------------------------|---|
| Body Weight: | 3.0 kg (Newell et al. 1987) |
| Ingestion Rate: | 0.6 kg/day (Newell et al. 1987) |
| Home Range: | Based on a 10 km radius feeding range, the home range of great blue heron is 30,000 ha (Erwin and Spindelov 1991). This equates to 74,130 acres. |
| Water Ingestion Rate: | 0.12 L/day estimated based on the allometric equation (Water Ingestion Rate = $0.059 Wt^{0.67}$, where Wt is the average body weight of the species in kg) |
| Diet: | Almost all aquatic (Erwin and Spindelov 1991) |
| Sediment Ingestion Rate | 54 g/day (based on 9% of the dietary ingestion rate) calculated based on soil ingestion rates reported for shore birds and Canada gees (Beyer et al. 1991) |

LIFE HISTORY OF THE MINK (*Mustela vison*)

Adult mink (*Mustela vison*) weigh from 520 to 1,730 g (Merritt 1987; U.S. EPA 1993). Home ranges vary from 19 to 1,900 acres (U.S. EPA 1993). The lowest reported body weight of 520 g and the smallest reported home range of 19 acres was assumed for this risk assessment.

Mink are opportunistic carnivores that hunt principally along shorelines and emergent vegetation (U.S. EPA 1993). Seasonal availability and regional preferences govern the primary constituent of the mink's diet. Mammals and crayfish usually result as the most abundant prey items, but fish, amphibians, and young birds are also taken (Merritt 1987; Linscombe *et al.* 1982; U.S. EPA 1993). A year-round food ingestion rate of 0.22 g/g BW/day has been estimated for both male and female mink (U.S. EPA 1993). In order to express this value in units of g/day, the food ingestion rate was multiplied by the lowest reported body weight (550 g) to yield a food ingestion rate of 121 g/day. The highest reported estimated water ingestion rate of 0.11 g/g BW/day for farm-raised females was used in this risk assessment (U.S. EPA 1993). In order to express this value in units of g/day, this water ingestion rate was multiplied by the lowest reported body weight of 550 g to yield a water ingestion rate of 60.5 g/day (60.5 ml/day).

An incidental soil or sediment ingestion rate was not available from the literature, therefore, a predicted incidental ingestion rate for soil and sediment rate that may be entrained in the digestive system of prey items (fish, aquatic invertebrates, and small mammals) was used for this risk assessment. Consumption of these prey items was assumed to be the primary mechanism by which a carnivorous mammal such as the mink may incidentally ingest soil or sediment. The derivation of these predicted levels of incidental soil and sediment ingestion via consumption of fish, aquatic invertebrates, and small mammals is described below.

Life history information for the bluegill (*Lepomis macrochirus*) was used to predict the amount of sediment that may be ingested by mink via consumption of fish. Adult bluegills range in size from 100 to 230 mm (Pflieger 1975; Smith 1985). In keeping with the conservative approach of this risk assessment, the amount of sediment entrained in the lowest body size of 100 mm in length was predicted. The weight of a 100 mm bluegill was calculated to be 18.11 g based on the following algorithm relating length to weight (Hillman 1982):

$$\log \text{Weight (g)} = -5.374 + 3.316 \log \text{Length (mm)}$$

A daily food ingestion rate of 1.75 percent of the body weight per day has been reported for the bluegill (Kolehmainen 1974). This provides a predicted intake rate of 0.32 g of food per day for a 18.11 g fish. A study evaluating the stomach contents of 153 bluegills reported an average content of detritus and sediment to be 9.6 percent of the total diet (Kolehmainen 1974). If a conservative assumption is made that 9.6 percent of the food ingested is entirely sediment, it can be predicted that a fish of this size may contain 0.03 g of sediment in its digestive system.

For the purpose of this model, it was assumed that the level of sediment contained in the digestive system of a fish remains constant over time. This value (0.03 g) was divided by the predicted fish body weight (18.11 g) in order to express sediment entrained in fish digestive systems in units of grams of sediment per gram of fish body weight. This provided a value of 0.0017 g sediment/g body weight. When this value is multiplied by the food ingestion rate of the mink (121 g/day), the predicted sediment ingestion rate for the mink via consumption of fish is 0.2 g/day.

Life history information for the crayfish *Orconectes* sp. was used in predicting the incidental sediment ingestion rate for the mink via consumption of freshwater invertebrates. Adult *O. virilis* weigh from 5 to 20+ g and consume 0.3 to 1 percent of its total body weight per day (Kim 1994; Tack 1941; Vannote 1963). In order to express the food ingestion rate in units of g/day, the highest reported food ingestion rate of 1 percent of the total body weight per day was multiplied by the lowest reported body weight of 5 g to yield a food ingestion rate of 0.05 g/day.

Orconectes spp. detritus ingestion rates range from 10 percent of the total diet per day in young-of-the-year *Orconectes immunis* (Vannote 1963) to 11 percent of the total diet per day in *O. virilis* (Tack 1941). For this risk assessment, it will be assumed that these values represent the percentage of sediment in the diet of a crayfish. The food ingestion rate of 0.05 g/day was multiplied by the incidental sediment ingestion rate of 11 percent of the total diet per day to yield an incidental sediment ingestion rate of 0.0055 g/day.

For the purpose of this model, it was assumed that the level of sediment contained in the digestive system of crayfish remains constant over time. Therefore, in order to express the amount of sediment entrained in a crayfish's digestive system in units of gram of sediment per gram of crayfish body weight, the sediment ingestion rate of 0.0055 g/day was divided by the lowest adult crayfish body weight of 5 g to yield a sediment ingestion rate of 0.0011 g sediment/g BW of crayfish. When this value is multiplied by the food ingestion rate of the mink (121 g/day), the predicted incidental sediment ingestion rate for the mink via consumption of crayfish is 0.133 g/day.

Life history information for the white-footed mouse was used in predicting the incidental soil ingestion rate for the mink via consumption of small mammals. A soil ingestion rate of less than 2 percent of the total diet has been reported for the white-footed mouse (Beyer *et al.* 1994). A conservative soil ingestion rate of 1.9 percent of the total diet was assumed for this risk assessment. In order to express this value in units of g/day, this soil ingestion rate was multiplied by the food ingestion rate of the white footed mouse (4.50 g/day; U.S. EPA 1993) to yield a soil ingestion rate of 0.0855 g/day.

For this risk assessment, it was assumed that the level of soil contained in the digestive system of a white-footed mouse remains constant over time. In order to express this value in units of g soil/g mouse BW, this value of 0.0855 g was divided by the lowest reported body weight of the white-footed mouse (13 g; Merritt 1987) to yield a value of 0.0065 g soil/g BW. When this value is multiplied by the food ingestion rate for the mink (121 g/day; U.S. EPA 1993), the predicted incidental soil ingestion rate for the mink via consumption of white-footed mice is 0.795 g/day.

Assuming that fish, crayfish, and mice comprise equal proportions of the mink's diet, the incidental soil and sediment ingestion rate via consumption of these prey items is approximately 0.4 g/day.

REFERENCES

- Beyer, W.N., E.E. Conner, and S. Gerould. 1994. "Estimates of Soil Ingestion by Wildlife." *J. Wildl. Manage.*, 58(2):375-382.
- Hillman, W.P. 1982. "Structure and Dynamics of Unique Bluegill Populations." Masters thesis. University of Missouri, Columbia, MO.
- Kim, P. 1994. Ecological Risk Assessment, Cannelton Industries Site. Roy F. Weston, Inc. Final Report to the U.S. EPA/ERT. Contract No. 68-C4-0022.
- Kolehmainen, S.E. 1974. Daily Feeding Rates of Bluegill (*Lepomis macrochirus*) Determined by a Refined Radioisotope Method. *J. Fish. Res. Bd. Can.*, 31:67-74.
- Linscombe, G., N. Kinler, and R.J. Aulerich. 1982. Mink. In: Chapman, J.A., G.A. Feldhamer, eds. *Wild Mammals of North America*. Baltimore, MD: John Hopkins University Press; pp.329-643.
- Merritt, J.F. 1987. "Guide to the Mammals of Pennsylvania." Univ. Pittsburgh Press, PA. 408p.
- Pflieger, W.L. 1975. "The Fishes of Missouri." Missouri Dept. Conserv.
- Smith, C.L. 1985. "The Inland Fishes of New York State." New York State Dept. Environ. Conserv.
- Tack, P.I. 1941. "The Life History and Ecology of the Crayfish, *Cambarus immunis* Hagen. *Am. Midl. Nat.* 25:420-466.
- U.S. EPA. 1993. "Wildlife Exposure Factors Handbook, Volume I of II." United States Environmental Protection Agency, Office of Research and Development, Washington, D.C. EPA/600/R-93/187a.
- Vannote, R.L. 1963. "Community Productivity and Energy Flow in an Enriched Warm Water Stream." Ph.D. Thesis, Michigan State Univer., E. Lansing. 156pp. In: Momot, W.T., H. Gowing, and P.D. Jones. 1978. "The Dynamics of

TOXICITY BENCHMARK VALUES

Copper

One study (OHMD, 1987) conducted showed that an oral dose of 100 mg/kg/day fed to a dog caused death. For this report, a NOAEL value of 1 mg/kg/day (100mg/kg/day divided by 10) was used in the mink model.

A dose of 350 mg/kg (61.3 mg/kg/day) caused a significant decrease in the growth and consumption by chickens (Smith, 1969). Another study on chickens, found a dose of 326 mg/kg (23.5 mg/kg/day) caused respiratory problems (Hatch, 1978). Assuming that respiratory problems are acute effects, a NOAEL (23.5 mg/kg/day divided by 100) of 0.235 mg/kg/day was developed for the heron model (Hatch, 1978).

Lead

A study conducted on mice determined that 1.5 mg/kg/day of lead caused a reduction in success of implanted ova (Clark, 1979). Another study (Clark, 1979) found that 2.2 mg/kg/day of lead produced a reduction in the frequency of pregnancy when the dose was administered for 3 to 5 days. For this report, a value of 0.15 mg/kg/day was used as a NOAEL (1.5 mg/kg/day divided by 10) was used in the mink model.

In a single dose study (Lawler et al., 1991), the gastric motility of adult male and female red-tailed hawks (*Buteo jamaicensis*) fed 0.82 and 1.64 mg/kg BW/day of lead was evaluated through the use of surgically implanted transducers for a period of 3 weeks. Neither concentration had any effect on the gastric contractions or egestion of undigested material pellets.

Another study conducted on red-tailed hawks (Reiser and Temple, 1981) found that 3 mg/kg/day of lead caused the clinical symptoms of lead poisoning. A similar study (Osborne et al., 1983) found that 3 mg/kg/day fed to starlings caused a reduction in muscle condition and altered their feeding activity. For this assessment, a value of 0.3 mg/kg/day NOAEL developed from red-tailed hawk and sterling studies was used in the heron model.

Reference

Clark, D.R., Jr. 1979. "Lead Concentrations: Bats vs. Terrestrial Mammals Collected near a Major Highway." Environ. Sci. Tech. 13:338-341.

Hatch, R.C. 1978. "Poisons Causing Respiratory Insufficiency." In: Veterinary Pharmacology and Therapeutics. L.M. Jones, N.H. Booth, and L.E. McDonald (eds.) Ames Press, Iowa State University. Ames, Iowa.

Lawler, E.M., G.E. Duke, and P.T. Redig. 1991. "Effect of sublethal lead exposure on gastric motility of red-tailed hawks." Arch. Environ. Contam. Toxicol. 21:78-83.

OHMD. Oil and Hazard Materials Technical Assistance Data Systems Database. OHMTAD. 1987. Developed by the Office of Water and Waste Management of the USEPA.

Osborne, D., W.J. Every, and K.R. Bull. 1983. "The Toxicity of Trialkyl Lead Compounds to Birds." Environmental Pollution (Series A). 31:261-275.

Reiser, and Temple, 1981

Smith, M.S. 1969. "Responses of chicks to dietary supplements of copper sulphate." British Poultry Science 10:97-108.

Stone Bay - Study Area
Aquatic Receptor Models
MCB, Camp Lejeune

| Ecological Contaminant of Concern | Sediment Mean (mg/kg) | Sediment Maximum (mg/kg) | Surface Water Mean (mg/L) | Surface Water Maximum (mg/L) | Tissue Mean (mg/L) | Tissue Maximum (mg/L) | Species with the Maxi. Concentration |
|-----------------------------------|-----------------------|--------------------------|---------------------------|------------------------------|--------------------|-----------------------|--------------------------------------|
| Copper | 9.23 | 16.6 | 0.000996 | 0.00153 | 35.98 | 50.3 | oyster |
| Lead | 7.66 | 20.7 | 0.0005278 | 0.000926 | 2.57 | 4.8 | mussel |

Stone Bay Aquatic Assessment
 MCB, Camp Lejeune

Great Blue Heron

Body Weight 3.000000 kg
 Food Ingestion Rate 0.600000 kg/day
 Water Ingestion Rate 0.120000 L/day
 Sediment Ingestion Rate 0.054000 kg/day

Maximum Concentrations - Conservative Model

| ECOC | Sediment Concentration (mg/kg) | Water Concentration (mg/L) | Fish Concentration (mg/kg) | Dose (mg/kg/day) | NOAEL (mg/kg/day) | LOAEL (mg/kg/day) | NOAEL HQ _n | LOAEL HQ _l |
|--------|--------------------------------|----------------------------|----------------------------|------------------|-------------------|-------------------|-----------------------|-----------------------|
| Copper | 16.6 | 0.00153 | 50.3 | 10.3588612 | 0.235 | 2.35 | 4.41E+01 | 4.41E+00 |
| Lead | 20.7 | 0.000926 | 4.8 | 1.33263704 | 0.3 | 3 | 4.44E+00 | 4.44E-01 |

NA - Not Available

HQ_n - Hazard Quotient based on the NOAEL

HQ_l - Hazard Quotient based on the LOAEL

NOAEL - No Observed Adverse Effects Level

LOAEL - Lowest Observed Adverse Effects Level

Stone Bay Aquatic Assessment
 MCB, Camp Lejeune

Great Blue Heron

Body Weight 3.0000000 kg
 Food Ingestion Rate 0.6000000 kg/day
 Water Ingestion Rate 0.1200000 L/day
 Sediment Ingestion Rate 0.0540000 kg/day

Mean Concentrations - Less Conservative Model

| | Sediment Concentration (mg/kg) | Water Concentration (mg/L) | Fish Concentration (mg/kg) | Dose (mg/kg/day) | NOAEL (mg/kg/day) | LOAEL (mg/kg/day) | NOAEL HQ _n | LOAEL HQ ₁ |
|--------|--------------------------------------|----------------------------------|----------------------------------|---------------------|----------------------|----------------------|--------------------------|--------------------------|
| ECOC | | | | | | | | |
| Copper | 9.23 | 0.000996 | 35.98 | 7.36217984 | 0.235 | 2.35 | 3.13E+01 | 3.13E+00 |
| Lead | 7.66 | 0.0005278 | 2.57 | 0.651901112 | 0.3 | 3 | 2.17E+00 | 2.17E-01 |

NA - Not Available

HQ_n - Hazard Quotient based on the NOAEL

HQ₁ - Hazard Quotient based on the LOAEL

NOAEL - No Observed Adverse Effects Level

LOAEL - Lowest Observed Adverse Effects Level

Stone Bay Aquatic Assessment
 MCB, Camp Lejeune

Mink

Body Weight 0.5200000 kg
 Food Ingestion Rate 0.1210000 kg/day
 Water Ingestion Rate 0.0605000 L/day
 Soil Ingestion Rate 0.0003330 kg/day

Maximum Concentrations - Conservative Model

| | Soil Concentration (mg/kg) | Water Concentration (mg/L) | Vegetation Concentration (mg/kg) | Invetebrate Concentration (mg/kg) | Dose (mg/kg/day) | NOAEL (mg/kg/day) | LOAEL (mg/kg/day) | NOAEL HQ _n | LOAEL HQ ₁ |
|--------|----------------------------------|----------------------------------|--|---|---------------------|----------------------|----------------------|--------------------------|--------------------------|
| ECOC | | | | | | | | | |
| Copper | 16.6 | 0.00153 | 6.64 | 50.3 | 13.2603084 | 1 | 10 | 1.33E+01 | 1.33E+00 |
| Lead | 20.7 | 0.000926 | 8.28 | 4.8 | 3.05697908 | 0.15 | 1.5 | 2.04E+01 | 2.04E+00 |

NA - Not Available

HQ_n - Hazard Quotient based on the NOAEL

HQ₁ - Hazard Quotient based on the LOAEL

NOAEL - No Observed Adverse Effects Level

LOAEL - Lowest Observed Adverse Effects Level

Stone Bay Aquatic Assessment
 MCB, Camp Lejeune

Mink

Body Weight 0.5200000 kg
 Food Ingestion Rate 0.1210000 kg/day
 Water Ingestion Rate 0.0605000 L/day
 Soil Ingestion Rate 0.0003330 kg/day

Mean Concentrations - Less Conservative Model

| | Soil Concentration (mg/kg) | Water Concentration (mg/L) | Vegetation Concentration (mg/kg) | Invertebrate Concentration (mg/kg) | Dose (mg/kg/day) | NOAEL (mg/kg/day) | LOAEL (mg/kg/day) | NOAEL HQ _n | LOAEL HQ _l |
|--------|----------------------------------|----------------------------------|--|--|---------------------|----------------------|----------------------|--------------------------|--------------------------|
| ECOC | | | | | | | | | |
| Copper | 9.23 | 0.000996 | 3.692 | 35.98 | 9.23739586 | 1 | 10 | 9.24E+00 | 9.24E-01 |
| Lead | 7.66 | 0.0005278 | 3.064 | 2.57 | 1.31595522 | 0.15 | 1.5 | 8.77E+00 | 8.77E-01 |

NA - Not Available

HQ_n - Hazard Quotient based on the NOAEL

HQ_l - Hazard Quotient based on the LOAEL

NOAEL - No Observed Adverse Effects Level

LOAEL - Lowest Observed Adverse Effects Level

Stone Bay - Reference Areas
 Aquatic Receptor Models
 MCB, Camp Lejeune

| Ecological Contaminant of Concern | Sediment Mean (mg/kg) | Sediment Maximum (mg/kg) | Surface Water Mean (mg/L) | Surface Water Maximum (mg/L) | Tissue Mean (mg/kg) | Tissue Maximum (mg/kg) | Species with the Maxi. Concentration |
|-----------------------------------|-----------------------|--------------------------|---------------------------|------------------------------|---------------------|------------------------|--------------------------------------|
| Copper | 4.6 | 4.6 | | | 48.2 | 88.2 | oyster |
| Lead | 7.55 | 10.5 | | | 1.6 | 2 | oyster |

Stone Bay Aquatic Assessment - Reference Areas
 MCB, Camp Lejeune

Mink

Body Weight 0.5200000 kg
 Food Ingestion Rate 0.1210000 kg/day
 Water Ingestion Rate 0.0605000 L/day
 Soil Ingestion Rate 0.0003330 kg/day

Maximum Concentrations - Conservative Model

| | Soil Concentration (mg/kg) | Water Concentration (mg/L) | Vegetation Concentration (mg/kg) | Invetebrate Concentration (mg/kg) | Dose (mg/kg/day) | NOAEL (mg/kg/day) | LOAEL (mg/kg/day) | NOAEL HQ _n | LOAEL HQ ₁ |
|--------|----------------------------------|----------------------------------|--|---|---------------------|----------------------|----------------------|--------------------------|--------------------------|
| ECOC | | | | | | | | | |
| Copper | 4.6 | 0 | 1.84 | 88.2 | 20.9545612 | 1 | 10 | 2.10E+01 | 2.10E+00 |
| Lead | 10.5 | 0 | 4.2 | 2 | 1.44941635 | 0.15 | 1.5 | 9.66E+00 | 9.66E-01 |

NA - Not Available
 HQ_n - Hazard Quotient based on the NOAEL
 HQ₁ - Hazard Quotient based on the LOAEL

NOAEL - No Observed Adverse Effects Level
 LOAEL - Lowest Observed Adverse Effects Level

Stone Bay Aquatic Assessment - Reference Areas
 MCB, Camp Lejeune

Mink

Body Weight 0.5200000 kg
 Food Ingestion Rate 0.1210000 kg/day
 Water Ingestion Rate 0.0605000 L/day
 Soil Ingestion Rate 0.0003330 kg/day

Mean Concentrations - Less Conservative Model

| ECOC | Soil Concentration (mg/kg) | Water Concentration (mg/L) | Vegetation Concentration (mg/kg) | Invertebrate Concentration (mg/kg) | Dose (mg/kg/day) | NOAEL (mg/kg/day) | LOAEL (mg/kg/day) | NOAEL HQ _n | LOAEL HQ ₁ |
|--------|----------------------------|----------------------------|----------------------------------|------------------------------------|------------------|-------------------|-------------------|-----------------------|-----------------------|
| Copper | 4.6 | 0 | 1.84 | 48.2 | 11.6468688 | 1 | 10 | 1.16E+01 | 1.16E+00 |
| Lead | 7.55 | 0 | 3.02 | 1.6 | 1.07987337 | 0.15 | 1.5 | 7.20E+00 | 7.20E-01 |

NA - Not Available

HQ_n - Hazard Quotient based on the NOAEL

HQ₁ - Hazard Quotient based on the LOAEL

NOAEL - No Observed Adverse Effects Level

LOAEL - Lowest Observed Adverse Effects Level